


Universität Konstanz



The permanence of R-boundedness and property(α) under
interpolation and applications to parabolic systems

Jürgen Saal
Mario Kaip

Konstanzer Schriften in Mathematik

Nr. 290, November 2011

ISSN 1430-3558

The permanence of \mathcal{R} -boundedness and property(α) under interpolation and applications to parabolic systems

Jürgen Saal

Center of Smart Interfaces
TU Darmstadt, Germany
saal@csi.tu-darmstadt.de

Mario Kaip

Department of mathematics
University of Konstanz, Germany
Mario.Kaip@uni-konstanz.de

MSC 2010: Primary 46B70, 47F05; Secondary 35K41, 35K46

Keywords: Interpolation, \mathcal{R} -boundedness, maximal regularity, parabolic systems

Abstract

This note consists of two parts. In the first part we consider the behavior of \mathcal{R} -boundedness, \mathcal{R} -sectoriality, and property(α) under the interpolation of Banach spaces. In a general setting we prove that for interpolation functors of type h the \mathcal{R} -boundedness, the \mathcal{R} -sectoriality, and the property(α) preserve under interpolation. In particular, this is true for the standard real and complex interpolation methods.¹ The second part represents an application of the first part. We prove \mathcal{R} -sectoriality, or equivalently, maximal L^p -regularity for a general class of parabolic systems on interpolation spaces including scales of Besov- and Bessel-potential spaces over \mathbb{R}^n .

1 Introduction

The concept of \mathcal{R} -bounded operator families nowadays plays an important role in the treatment of linear and nonlinear problems. By the celebrated result of L. Weis [19], it is known that \mathcal{R} -boundedness of the resolvent family $\lambda(\lambda + A)^{-1}$ for λ in a complex sector with opening angle greater than $\pi/2$ implies maximal regularity for a linear operator A . The maximal regularity, in turn, is fundamental in the treatment of linear and nonlinear PDEs for various reasons: the construction of local-in-time strong solutions, of global weak solutions, of real analytic solutions, uniqueness proofs, a.s.o.

Also, in combination with a holomorphic functional calculus, a so-called H^∞ -calculus, the concept of \mathcal{R} -boundedness turned out to be very valuable. It allows for the introduction of a joint H^∞ -calculus of two closed linear operators A, B , cf. [13]. In particular, it gives an answer to the question under what circumstances $f(A, B)$ gives rise to a bounded operator for bounded real analytic functions $f(\lambda, z)$. In the simplest case we might have $f(\lambda, z) = (\lambda + z)^{-1}$, $A = d/dt$, and $B = -\Delta$, for instance. The full strength of such a joint H^∞ -calculus reveals in the treatment of free boundary value problems, see e.g. [6], [15]. This type problems often can be reduced to the boundary, on which, however, one is faced to a mixed order system. The associated matrix symbols (Lopatinskii matrix) not seldom have a complicated structure, but still are real analytic functions and therefore fit into the framework of the joint H^∞ -calculus developed in [13] or [10].

In this context also a geometric property of a Banach space (besides the property 'of class \mathcal{HT} ') comes into play: the so-called 'property(α)'. In many situations it represents the crucial ingredient for the step from uniform boundedness to \mathcal{R} -boundedness. For instance, if property(α) for a Banach space X is assumed, the standard multiplier results, if applicable, yield \mathcal{R} -boundedness of an operator family $(M_\lambda)_\lambda$ in a parameter λ , instead of uniform boundedness only. For $M_\lambda = \lambda(\lambda + A)^{-1}$ this leads directly to the \mathcal{R} -sectoriality of A or, equivalently, to the maximal L^p -regularity. We refer to [14, Theorem 5.2 b)] for a multiplier result of this type. If an operator has a bounded H^∞ -calculus on X , it automatically admits the

¹Partly, these results were indicated in [12], however, with just a very brief outline of their proofs.

stronger property of an \mathcal{R} -bounded H^∞ -calculus, provided X has property(α). This is another significant consequence of property(α). This fact particularly matters for the initialization and the application of a joint H^∞ -calculus, cf. [13].

For all these reasons, it is important to know about the behavior of the notions of \mathcal{R} -boundedness and property(α) with respect to other functional analytic operations, such as the interpolation of Banach spaces, for instance. In the first part of this paper we clarify this behavior. In fact, we will show that both properties preserve under interpolation. These results are indicated in [12]. However, their proofs are just outlined and for readers not so experienced in this topic it might be hard to follow the very brief argumentation given in [12]. It also seems that a rigorous proof so far is not contained anywhere else in the available literature, although the results are not seldom used in other works. With the aim to apply them in order to prove \mathcal{R} -sectoriality for a class of parameter-elliptic systems on Besov and Bessel-potential spaces, here we give a rigorous proof of the results on the interpolation of \mathcal{R} -boundedness and property(α) indicated in [12]. Indeed, we prove the preservation of these two properties in a very general setting for exact interpolation functors of type θ . This covers the cases of real and complex interpolation and also generalizes the results indicated in [12].

The proof of the permanence results under discussion is based on the characterization of \mathcal{R} -boundedness in terms of boundedness in Rademacher spaces. The Rademacher spaces are complementary in $L^p([0, 1], X)$ if X is of class \mathcal{HT} . By these facts the interpolation of \mathcal{R} -boundedness is reduced to the interpolation of X -valued L^p -spaces and to general facts concerning the interpolation of complementary subspaces. The preservation of property(α) under interpolation then can be reduced to the obtained results on the interpolation of \mathcal{R} -boundedness. Indeed, property(α) can be regarded as a special form of \mathcal{R} -boundedness on Rademacher spaces. Therefore the results on \mathcal{R} -boundedness apply.

This is also the reason, why we give the proof of the preservation of property(α) under interpolation here, although this result is not directly applied in this note. In a forthcoming work it will be applied in order to prove an \mathcal{R} -bounded H^∞ -calculus for a certain class of elliptic operators on scales of Besov and Bessel-potential spaces.

In the second part of this note we will apply the results obtained in the first part in order to prove \mathcal{R} -sectoriality for a class of parameter-elliptic systems realized on interpolation spaces. For classical works on parameter-elliptic systems we refer to [2] and [17]. Particularly, we will apply the obtained permanence results to real and complex interpolation functors. This yields the \mathcal{R} -sectoriality on scales of Besov and Bessel-potential spaces. For this purpose, we first establish the corresponding result for the model problem (i.e., constant coefficients) in Sobolev spaces $W^{k,p}(\mathbb{R}^n, \mathbb{C}^n)$ (see Proposition 5.12). This will be based on a multiplier theorem. Employing a localization procedure and perturbation arguments, the result generalizes to a class of variable coefficients (see Theorem 5.28). This result can be found in [14] for the special case $k = 0$. Interpolation and the outcome of the first part of this note then imply the \mathcal{R} -sectoriality on scales of interpolation spaces (see Theorem 5.29).

The paper is organized as follows. The first part includes Section 2 to 4. In Section 2 we clarify the notation. Section 3 includes the introduction of \mathcal{R} -bounded families, the characterization via Rademacher spaces, and the result on the permanence of \mathcal{R} -boundedness under interpolation (Theorem 3.18). The introduction, the characterization in terms of \mathcal{R} -boundedness, and the corresponding results on the interpolation of property(α) are the content of Section 4. In the second part, i.e. in Section 5, we prove the mentioned \mathcal{R} -sectoriality for a class of parameter-elliptic systems realized on scales of interpolation spaces. The main results here are Theorem 5.29 and Corollary 5.31.

2 Notation

Definition 2.1. *In the sequel we use the following notation:*

- The set $\{X_0, X_1\}$ is said to be a interpolation couple, if X_0 and X_1 are Banach spaces, which are embedded in a Hausdorff topological vector space \mathcal{X} . On $X_0 + X_1$ we define the norm $\|x\|_{X_0+X_1} := \inf_{x_k \in X_k: x_0+x_1=x} (\|x_0\|_{X_0} + \|x_1\|_{X_1})$ for $x \in X_0 + X_1$.
- Let $\{X_0, X_1\}, \{Y_0, Y_1\}$ be interpolation couples, then we define

$L(\{X_0, X_1\}, \{Y_0, Y_1\}) := \{T : X_0 + X_1 \rightarrow Y_0 + Y_1 \mid T \text{ linear and } T|_{X_k} \in L(X_k, Y_k), k = 0, 1\}$ and $L(\{X_0, X_1\}) := L(\{X_0, X_1\}, \{X_0, X_1\})$.

- For normed spaces X and Y we denote the existence of an injective continuous linear mapping from Y to X by $Y \hookrightarrow X$. By $L(Y, X)$ we denote the space of all linear and bounded operators from Y into X .
- By $Y \hookrightarrow_d X$ we denote the existence of an injective continuous linear mapping from Y to X with dense image in X .
- Let X and Y be normed spaces. The equality $X = Y$ is used with the meaning that there exists an isomorphism between X and Y . In particular we have equivalence of the norms in this case.
- By virtue of the appearance of various spaces we often want to make clear in which space an integral or a series converges. Hence we write “ $\int \dots dx_{[X]}$ ” and “ ${}_{[X]} \sum \dots$ ” to indicate the convergence in X .

For the definition of an interpolation functor \mathcal{F} we follow [18, 1.2.2]. Particularly we make use of interpolation functors of *type h*, which means that we have an estimate as

$$\|T|_{\mathcal{F}(\{X_0, X_1\})}\|_{L(\mathcal{F}(\{X_0, X_1\}), \mathcal{F}(\{Y_0, Y_1\}))} \leq C_h \cdot h(\|T|_{X_0}\|_{L(X_0, Y_0)}, \|T|_{X_1}\|_{L(X_1, Y_1)})$$

with a constant $C_h > 0$. An interpolation functor is said to be *exact of type θ* , if it is of type $h(t_0, t_1) := t_0^{1-\theta} t_1^\theta$ with $C_h = 1$. Important examples of exact interpolation functors of type θ are given by the *real and complex interpolation methods*, which are defined as in [1, 7.9/7.51]. The proof of the exactness for these two methods can be found e.g. in [18, 1.3.3 (a)/1.9.3 (a)]. As usually, we denote the real interpolation space by $(X_0, X_1)_{\theta, p}$ and the complex interpolation space by $[X_0, X_1]_\theta$. We want to mention that in this note we only use the *K-method* for real interpolation.

Let $(\Omega, \mathcal{B}, \mu)$ be a measure space with a σ -finite measure μ on \mathcal{B} . For a Banach space X we denote the Banach space valued L^p -space $L^p(\Omega, \mathcal{B}, \mu, X)$ by $L^p(X)$. For an domain $\Omega \subset \mathbb{R}^n$ we denote the Sobolev space of order $m \in \mathbb{N}_0$ by $W^{m,p}(\Omega, X)$ or $W^{m,p}(X)$. Here and in the following we always consider the case $p \in (1, \infty)$. An interpolation functor \mathcal{F} is said to be *L^p -compatible*, if we have $\mathcal{F}(\{L^p(X_0), L^p(X_1)\}) = L^p(\mathcal{F}(\{X_0, X_1\}))$ for all interpolation couples $\{X_0, X_1\}$. In [18] it is proved that real and complex interpolation methods are *L^p -compatible* interpolation functors such that

$$C_p^{(1)} \|\cdot\|_{(L^p(X_0), L^p(X_1))_{\theta, p}} \leq \|\cdot\|_{L^p((X_0, X_1)_{\theta, p})} \leq C_p^{(2)} \|\cdot\|_{(L^p(X_0), L^p(X_1))_{\theta, p}}, \quad (1)$$

$$\|\cdot\|_{[L^p(X_0), L^p(X_1)]_\theta} = \|\cdot\|_{L^p([X_0, X_1]_\theta)} \quad (2)$$

where the constants $C_p^{(1)} > 0$ and $C_p^{(2)} > 0$ are independent of the spaces X_0 and X_1 .

Another important property of a Banach space X is the continuity of the Hilbert transform $H = \mathcal{F}^{-1}[i\xi/|\xi|]\mathcal{F}$ on $L^p(\mathbb{R}, X)$. If this is satisfied, X is said to be of class \mathcal{HT} (or equivalently UMD), cf. [3, 4.4.1 Theorem].

Remark 2.2. *If two Banach spaces X and Y are of class \mathcal{HT} , then the real and complex interpolation spaces in between X and Y are also of class \mathcal{HT} . By analyzing the proof in [3, 4.5.2 Theorem (vii)] one obtains that this is also true for all L^p -compatible interpolation functors. With [3, Theorem 4.5.2] it can be easily seen that $W^{m,p}(\Omega, X)$ ($m \in \mathbb{N}_0$) is also of class \mathcal{HT} . Therefore all closed subspaces of $W^{m,p}(\Omega, X)$ are of class \mathcal{HT} , due to [3, Theorem 4.5.2 (viii)].*

To avoid confusion with different definitions of the resolvent set, we give it here. Under the resolvent set $\rho(A)$ of a linear and densely defined operator $A : D(A) \subset X \rightarrow X$ we understand the set of all $\lambda \in \mathbb{C}$ such that $(\lambda - A) : D(A) \rightarrow X$ is bijective. If A is closed we obviously have $\{(\lambda - A)^{-1} : \lambda \in \rho(A)\} \subset L(X)$.

3 \mathcal{R} -boundedness and Rademacher spaces

3.1 Basic properties and definitions

The following definitions and basic consequences can be found in detail in [14, Section 2] or [8, Section 11].

Definition 3.1 (\mathcal{R} -boundedness). *Let X, Y be Banach spaces, $\mathcal{T} \subset L(X, Y)$, and $p \in [1, \infty)$. Then \mathcal{T} is said to be \mathcal{R} -bounded, if there exists a constant $C > 0$ such that for all $m \in \mathbb{N}$, $(T_k)_{k=1, \dots, m} \subset \mathcal{T}$, and all $(x_k)_{k=1, \dots, m} \subset X$ we have*

$$\left\| \sum_{k=1}^m r_k T_k x_k \right\|_{L^p([0,1], Y)} \leq C \left\| \sum_{k=1}^m r_k x_k \right\|_{L^p([0,1], X)}. \quad (3)$$

Then $\mathcal{R}_p(\mathcal{T}) := \min\{C > 0 : (\mathcal{B}) \text{ is satisfied}\}$ is said to be the \mathcal{R} -bound of \mathcal{T} . For $k \in \mathbb{N}$ the functions $r_k : [0, 1] \rightarrow \{-1, 1\}$, $t \mapsto \text{sign}(\sin(2^k \pi t))$ are called Rademacher-functions.

Next we introduce the Rademacher spaces. With their help the intricate definition of \mathcal{R} -boundedness can be characterized in a convenient way.

Definition 3.2 ($\text{Rad}_p(X)$ and $\text{Rad}_p^m(X)$). *For a Banach space X , $p \in [1, \infty)$, and $m \in \mathbb{N}$ the spaces*

$$\begin{aligned} \text{Rad}_p(X) &:= \left\{ (x_k)_{k \in \mathbb{N}} \subset X : \sum_{k=1}^{\infty} r_k x_k \text{ convergent in } L^p([0, 1], X) \right\} \\ \text{Rad}_p^m(X) &:= \{(x_k)_{k=1, \dots, m} \subset X\}, \end{aligned}$$

equipped with the norms

$$\begin{aligned} \|(x_k)_k\|_{\text{Rad}_p(X)} &:= \left\| \sum_{k=1}^{\infty} r_k x_k \right\|_{L^p([0,1], X)}, \\ \|(x_k)_{k=1, \dots, m}\|_{\text{Rad}_p^m(X)} &:= \left\| \sum_{k=1}^m r_k x_k \right\|_{L^p([0,1], X)} \end{aligned}$$

respectively, are called Rademacher spaces.

Remark 3.3. (i) *The spaces $\text{Rad}_p(X)$ and $\text{Rad}_p^m(X)$ are Banach spaces and $\bigcup_{m=1}^{\infty} \text{Rad}_p^m(X)$ is dense in $\text{Rad}_p(X)$.*

(ii) *Let X be a Banach space, $p \in [1, \infty)$, and $m \in \mathbb{N}$. Then we have the following norm preserving embeddings*

$$\begin{aligned} \text{Rad}_p(X) &\hookrightarrow L^p([0, 1], X), & (x_k)_k &\mapsto \sum_{k=1}^{\infty} r_k x_k, \\ \text{Rad}_p^m(X) &\hookrightarrow L^p([0, 1], X), & (x_k)_{k=1, \dots, m} &\mapsto \sum_{k=1}^m r_k x_k. \end{aligned}$$

(iii) *If X is of class \mathcal{HT} , Remark 2.2 implies that $\text{Rad}_p^m(X)$ and $\text{Rad}_p(X)$ are also of class \mathcal{HT} , since they are closed subspaces of $L^p([0, 1], X)$.*

Theorem 3.4 (Inequality of Kahane). *Let X be a Banach space and $p \in [1, \infty)$. Then there exists a constant $C_p^{(K)} > 0$, such that for all $(x_k)_{k \in \mathbb{N}} \subset X$ we have*

$$\frac{1}{C_p^{(K)}} \left\| \sum_{k=1}^{\infty} r_k x_k \right\|_{L^1([0,1], X)} \leq \left\| \sum_{k=1}^{\infty} r_k x_k \right\|_{L^p([0,1], X)} \leq C_p^{(K)} \left\| \sum_{k=1}^{\infty} r_k x_k \right\|_{L^1([0,1], X)}.$$

Theorem 3.5 (Kahane's contraction principle). *Let X be a Banach space and $p \in [1, \infty)$. Then we have*

$$\left\| \sum_{j=1}^n r_j a_j x_j \right\|_{L^p([0,1], X)} \leq 2 \left\| \sum_{j=1}^n r_j b_j x_j \right\|_{L^p([0,1], X)}$$

for all $n \in \mathbb{N}$, all $(a_j)_j, (b_j)_j \subset \mathbb{C}$ with $|a_j| \leq |b_j|$, and all $(x_j)_j \subset X$.

Remark 3.6 (Some well-known facts about \mathcal{R} -boundedness). *The proof of the following results can be found in [5].*

- (i) *If $\mathcal{T} \subset L(X, Y)$ is \mathcal{R} -bounded for one $p \in [1, \infty)$, then we have an estimate as (3) for all $p \in [1, \infty)$. The \mathcal{R} -bounds can be estimated as $[C_p^{(K)}]^{-2} \mathcal{R}_1(\mathcal{T}) \leq \mathcal{R}_p(\mathcal{T}) \leq [C_p^{(K)}]^2 \mathcal{R}_1(\mathcal{T})$.*
- (ii) *If $\mathcal{T}, \mathcal{S} \subset L(X, Y)$ are \mathcal{R} -bounded, then $\mathcal{T} + \mathcal{S} := \{T + S : T \in \mathcal{T}, S \in \mathcal{S}\}$ is also \mathcal{R} -bounded with $\mathcal{R}_p(\mathcal{T} + \mathcal{S}) \leq \mathcal{R}_p(\mathcal{T}) + \mathcal{R}_p(\mathcal{S})$.*
- (iii) *For two given \mathcal{R} -bounded families $\mathcal{T}_1 \subset L(Z, Y)$ and $\mathcal{T}_2 \subset L(X, Z)$ we obtain the \mathcal{R} -boundedness of $\mathcal{T}_1 \mathcal{T}_2 := \{T_1 T_2 : T_k \in \mathcal{T}_k, k = 1, 2\} \subset L(X, Y)$ with $\mathcal{R}_p(\mathcal{T}_1 \mathcal{T}_2) \leq \mathcal{R}_p(\mathcal{T}_1) \cdot \mathcal{R}_p(\mathcal{T}_2)$.*
- (iv) *If $\mathcal{T} \subset L(X, Y)$ is \mathcal{R} -bounded, then \mathcal{T} is also uniformly bounded. The converse, in general, is only true if X and Y are both Hilbert spaces.*

Remark 3.7. *For a given $C > 0$ we have the equivalence of the following three statements:*

- (i) *For all $m \in \mathbb{N}$ and $(T_k)_{k=1, \dots, m} \subset \mathcal{T}$ we have that $\|\mathbb{T}_m\|_{L(\text{Rad}_p^m(X), \text{Rad}_p^m(Y))} \leq C$, where \mathbb{T}_m is defined by*

$$\mathbb{T}_m : \text{Rad}_p^m(X) \rightarrow \text{Rad}_p^m(Y), \quad \sum_{k=1}^m r_k x_k \mapsto \sum_{k=1}^m r_k T_k x_k.$$

- (ii) *For all $(T_k)_{k \in \mathbb{N}} \subset \mathcal{T}$ the operator*

$$\mathbb{T} : \text{Rad}_p(X) \longrightarrow \text{Rad}_p(Y), \quad \sum_{k=1}^{\infty} r_k x_k \mapsto \sum_{k=1}^{\infty} r_k T_k x_k$$

is well-defined and $\|\mathbb{T}\|_{L(\text{Rad}_p(X), \text{Rad}_p(Y))} \leq C$.

- (iii) *$\mathcal{T} \subset L(X, Y)$ is \mathcal{R} -bounded with $\mathcal{R}_p(\mathcal{T}) \leq C$.*

Proof. This is obtained as an easy consequence of Remark 3.3. □

Thanks to Remark 3.7 the behavior of \mathcal{R} -boundedness under interpolation is completely reduced to the investigation of the interpolation of the Rademacher spaces. Having this in mind, we next analyze the corresponding properties of these spaces. To this end, the existence of projections onto $\text{Rad}_p^m(X)$ and $\text{Rad}_p(X)$ will turn out to be helpful. For the space $\text{Rad}_p^m(X)$ we easily obtain the following result:

Theorem 3.8. *The operator*

$$R_m^X : L^p([0, 1], X) \longrightarrow L^p([0, 1], X), f \mapsto \sum_{k=1}^m r_k \int_0^1 r_k(u) f(u) du$$

is continuous and even a projection onto $\text{Rad}_p^m(X)$.

The existence of a projection onto $\text{Rad}_p(X)$ is a more involved issue. The study of this problem requires some knowledge on the geometry of Banach spaces. Here we just cite the following result which is obtained as a corollary of [16, Remark 3.1.] and [8, Section 13].

Theorem 3.9 (Projection onto $\text{Rad}_p(X)$). *Let X be a Banach space of class \mathcal{HT} and $p \in (1, \infty)$. Then*

$$R^X : L^p([0, 1], X) \rightarrow L^p([0, 1], X); \quad f \mapsto \sum_{k=1}^{\infty} r_k \left(\int_0^1 r_k(u) f(u) du \right) \quad (4)$$

defines a bounded operator which is a projection onto $\text{Rad}_p(X)$.

Remark 3.10 (K-Convexity). *In our case it is sufficient to consider spaces of class \mathcal{HT} . Some of the following results (in particular Corollary 3.16 and 3.19) can be generalized to the more general class of 'K-convex' spaces. The space X is called 'K-convex' if the Rademacher projection in (4) exists. One can show that this is equivalent to 'B-convexity' and 'non-trivial type', see for example in [8].*

The results of the Corollaries 3.16, 3.19, and 4.6 remain valid if we only assume that X_0 and X_1 are K-convex. The proofs are exactly the same but we need the permanence of K-convexity under real and complex interpolation which can be found in [11, Proposition 5.1] for example.

3.2 Interpolation of $\text{Rad}_p(X)$ and $\text{Rad}_p^m(X)$

In order to obtain a suitable characterization of the interpolation spaces $(\text{Rad}_p(X), \text{Rad}_p(Y))_{\theta, p}$ and $[\text{Rad}_p(X), \text{Rad}_p(Y)]_{\theta}$ we will apply the abstract isomorphism result derived in [18, 1.2.4]. The idea to use this isomorphism result is taken from [12, Prop. 3.7].

Definition 3.11. *Let X and Y be Banach spaces. The operator $R \in L(X, Y)$ is said to be a retraction if there exists an $S \in L(Y, X)$ with $RS = \text{id}_Y$. In this case S is said to be the coretraction belonging to R .*

Remark 3.12. *Let X be a Banach space and $U \subset X$ be a closed subspace with the standard subspace topology. If there exists a projection $P \in L(X)$ with $\text{range}(P) = U$ then it is easy to see, that P is a retraction with coretraction $S : U \rightarrow X, x \mapsto x$.*

Theorem 3.13 (see [18], Theorem 1.2.4). *Let $\{A_0, A_1\}, \{B_0, B_1\}$ be two interpolation couples and let*

$$R \in L(\{A_0, A_1\}, \{B_0, B_1\}), \quad S \in L(\{B_0, B_1\}, \{A_0, A_1\})$$

such that $R|_{A_k} \in L(A_k, B_k)$ and $S|_{B_k} \in L(B_k, A_k)$ are retraction and coretraction ($k = 0, 1$). Then for an arbitrary interpolation functor \mathcal{F} we have that $(SR)|_{\mathcal{F}(\{A_0, A_1\})} \in L(\mathcal{F}(\{A_0, A_1\}))$ is a projection onto $W := \text{range}((SR)|_{\mathcal{F}(\{A_0, A_1\})}) \subset \mathcal{F}(\{A_0, A_1\})$, where the topology on W is given by the subspace topology relative to $\mathcal{F}(\{A_0, A_1\})$. In particular, the mapping $S|_{\mathcal{F}(\{B_0, B_1\})}$ yields an isomorphism between $\mathcal{F}(\{B_0, B_1\})$ and W .

Proposition 3.14 ($\mathcal{F}(\{\text{Rad}_p(X_0), \text{Rad}_p(X_1)\})$). *Let $\{X_0, X_1\}$ be an interpolation couple of Banach spaces, which are of class \mathcal{HT} . For $p \in (1, \infty)$ and an L^p -compatible interpolation functor \mathcal{F} of type h with $C_h \geq 1$ we have*

$$\mathcal{F}(\{\text{Rad}_p(X_0), \text{Rad}_p(X_1)\}) = \text{Rad}_p(\mathcal{F}(\{X_0, X_1\})),$$

where the equivalence of the norms is given by

$$\frac{1}{C_h C_2} \|f\|_{\text{Rad}_p(\mathcal{F}(\{X_0, X_1\}))} \leq \|f\|_{\mathcal{F}(\{\text{Rad}_p(X_0), \text{Rad}_p(X_1)\})} \leq \frac{C_h}{C_1} \cdot h(\|R^{X_0}\|, \|R^{X_1}\|) \|f\|_{\text{Rad}_p(\mathcal{F}(\{X_0, X_1\}))},$$

The constants $C_1, C_2 > 0$ come from the assumed equivalence

$$C_1 \| \cdot \|_{\mathcal{F}(\{L^p([0,1], X_0), L^p([0,1], X_1)\})} \leq \| \cdot \|_{L^p([0,1], \mathcal{F}(\{X_0, X_1\}))} \leq C_2 \| \cdot \|_{\mathcal{F}(\{L^p([0,1], X_0), L^p([0,1], X_1)\})}.$$

Proof. The aim is, of course, to apply Theorem 3.13. Therefore we define the spaces $A_k := L^p([0, 1], X_k)$ and $B_k := \text{Rad}_p(X_k)$. Let $R_k := R^{X_k} \in L(L^p([0, 1], X_k), \text{Rad}_p(X_k))$ be the projection given through Theorem 3.9 and let $S_k \in L(\text{Rad}_p(X_k), L^p([0, 1], X_k))$ be the embedding that exists according to Remark 3.12

and Remark 3.3 (ii). On $\mathbb{E} := L^p([0, 1], X_0 + X_1)$ we define the operators

$$\begin{aligned} R : (A_0 + A_1) &\rightarrow (B_0 + B_1) \quad , \quad f \mapsto \sum_{k=1}^{\infty} r_k \int_0^1 r_k(u) f(u) du_{[X_0+X_1]} \\ S : (B_0 + B_1) &\rightarrow (A_0 + A_1) \quad , \quad g \mapsto g. \end{aligned}$$

It can be easily seen that R and S are well-defined and that we have $R|_{L^p([0,1], X_k)} = R_k$ and $S|_{\text{Rad}_p(X_k)} = S_k$. It is also clear that S is norm preserving. Thus we can apply Theorem 3.13. At first this implies that

$$W := \text{range}((SR)|_{\mathcal{F}(\{A_0, A_1\})}) = \text{range}(R|_{\mathcal{F}(\{L^p([0,1], X_0), L^p([0,1], X_1)\})})$$

is well-defined. Due to Remark 2.2 we see that $\mathcal{F}(\{X_0, X_1\})$ is also of class \mathcal{HT} . Therefore we obtain the existence of the projection $R^{\mathcal{F}(\{X_0, X_1\})}$ onto $\text{Rad}_p(\mathcal{F}(\{X_0, X_1\}))$. Employing $L^p([0, 1], \mathcal{F}(\{X_0, X_1\})) \hookrightarrow \mathbb{E}$ and the L^p -compatibility of \mathcal{F} we obtain

$$W = \text{range}(R|_{L^p([0,1], \mathcal{F}(\{X_0, X_1\})})) = \text{range}(R^{\mathcal{F}(\{X_0, X_1\})}) = \text{Rad}_p(\mathcal{F}(\{X_0, X_1\})).$$

So far these equalities are only equalities of sets by the fact that on W we have the relative topology with respect to $\mathcal{F}(\{L^p([0, 1], X_0), L^p([0, 1], X_1)\})$. However, the L^p -compatibility of \mathcal{F} yields the topological equality of W and $\text{Rad}_p(\mathcal{F}(\{X_0, X_1\}))$. So we have $\mathcal{F}(\{\text{Rad}_p(X_0), \text{Rad}_p(X_1)\}) = W$ thanks to Theorem 3.13.

It remains to determine the constants which are involved in the equivalence of the norms. The L^p -compatibility of \mathcal{F} and the definition of $\|\cdot\|_{\text{Rad}_p(\mathcal{F}(\{X_0, X_1\}))}$ yield

$$\begin{aligned} \|f\|_{\text{Rad}_p(\mathcal{F}(\{X_0, X_1\}))} &= \|f\|_{L^p([0,1], \mathcal{F}(\{X_0, X_1\}))} \leq C_2 \|f\|_{\mathcal{F}(\{L^p([0,1], X_0), L^p([0,1], X_1)\})} \\ &\leq C_2 C_h \|f\|_{\mathcal{F}(\{\text{Rad}_p(X_0), \text{Rad}_p(X_1)\})} \quad (f \in W). \end{aligned}$$

In the last estimate we used the fact that the S_k 's are norm preserving. In view of $R|_{\mathcal{F}(\{A_0, A_1\})} f = f$ for all $f \in W$ and again by the L^p -compatibility we obtain

$$\begin{aligned} \|f\|_{\mathcal{F}(\{\text{Rad}_p(X_0), \text{Rad}_p(X_1)\})} &\leq C_h \cdot h(\|R_0\|, \|R_1\|) \cdot \|f\|_{\mathcal{F}(\{L^p([0,1], X_0), L^p([0,1], X_1)\})} \\ &\leq \frac{C_h}{C_1} h(\|R_0\|, \|R_1\|) \cdot \|f\|_{L^p([0,1], \mathcal{F}(\{X_0, X_1\}))} \quad (f \in W). \end{aligned}$$

This implies the claimed equivalence of the norms and therefore the assertion is proved. \square

Completely analogous we can obtain the following interpolation result for the space $\text{Rad}_p^m(X)$.

Proposition 3.15 ($\mathcal{F}(\{\text{Rad}_p^m(X_0), \text{Rad}_p^m(X_1)\})$). *Let $\{X_0, X_1\}$ be an interpolation couple, $p \in (1, \infty)$, and $m \in \mathbb{N}$. If \mathcal{F} is an L^p -compatible interpolation functor of type h with $C_h \geq 1$, then we have*

$$\mathcal{F}(\{\text{Rad}_p^m(X_0), \text{Rad}_p^m(X_1)\}) = \text{Rad}_p^m(\mathcal{F}(\{X_0, X_1\})),$$

where the equivalence of the norms is given by

$$\frac{1}{C_h C_2} \|f\|_{\text{Rad}_p^m(\mathcal{F}(\{X_0, X_1\}))} \leq \|f\|_{\mathcal{F}(\{\text{Rad}_p^m(X_0), \text{Rad}_p^m(X_1)\})} \leq \frac{C_h}{C_1} h(\|R_m^{X_0}\|, \|R_m^{X_1}\|) \|f\|_{\text{Rad}_p^m(\mathcal{F}(\{X_0, X_1\}))}.$$

The constants $C_1, C_2 > 0$ are the same as in Proposition 3.14.

Corollary 3.16. *The results of Proposition 3.14 and Theorem 3.15 in particular hold for the real and the complex interpolation functors. Let $\{X_0, X_1\}$ be an interpolation couple such that X_0 and X_1 are of class \mathcal{HT} . Then we have*

$$\begin{aligned} (\text{Rad}_p(X_0), \text{Rad}_p(X_1))_{\theta, p} &= \text{Rad}_p((X_0, X_1)_{\theta, p}), \\ [\text{Rad}_p(X_0), \text{Rad}_p(X_1)]_{\theta} &= \text{Rad}_p([X_0, X_1]_{\theta}) \end{aligned}$$

for $p \in (1, \infty)$ and $0 < \theta < 1$.

3.3 \mathcal{R} -boundedness and interpolation

Definition 3.17. Let $\{X_0, X_1\}$ and $\{Y_0, Y_1\}$ be interpolation couples and $\mathcal{T} \subset L(\{X_0, X_1\}, \{Y_0, Y_1\})$. Then we define $\mathcal{T}_{|X_k} := \{T_{|X_k} : T \in \mathcal{T}\} \subset L(X_k, Y_k)$ for $k = 0, 1$.

Notation: In the following we set $L^p(X) := L^p([0, 1], X)$.

Theorem 3.18 (\mathcal{R} -boundedness and interpolation). Let $\{X_0, X_1\}$ and $\{Y_0, Y_1\}$ be interpolation couples of Banach spaces, which are all of class \mathcal{HT} . Assume that $\mathcal{T} \subset L(\{X_0, X_1\}, \{Y_0, Y_1\})$ and that \mathcal{F} is an L^p -compatible interpolation functor of type h with $C_h \geq 1$ for a $p \in (1, \infty)$. If $\mathcal{T}_{|X_k} \subset L(X_k, Y_k)$ is \mathcal{R} -bounded with \mathcal{R} -bound $\mathcal{R}_p(\mathcal{T}_{|X_k})$ for $k = 0, 1$ then

$$\mathcal{T}_{|\mathcal{F}(\{X_0, X_1\})} \subset L(\mathcal{F}(\{X_0, X_1\}), \mathcal{F}(\{Y_0, Y_1\}))$$

is also \mathcal{R} -bounded with

$$\mathcal{R}_p(\mathcal{T}_{|\mathcal{F}(\{X_0, X_1\})}) \leq C_0 \cdot h(\|R^{X_0}\|, \|R^{X_1}\|) \cdot h(\mathcal{R}_p(\mathcal{T}_{|X_0}), \mathcal{R}_p(\mathcal{T}_{|X_1})),$$

where $C_0 := \frac{C_2 C_h^3}{C_1}$ and where $C_1, C_2 > 0$ come from Proposition 3.14.

Proof. We use the characterization of \mathcal{R} -boundedness given in Remark 3.7 (ii). Let $(T_j)_{j \in \mathbb{N}} \subset \mathcal{T}$ be an arbitrary series of operators. We have to show that

$$\begin{aligned} \mathbb{T} : \text{Rad}_p(\mathcal{F}(\{X_0, X_1\})) &\longrightarrow \text{Rad}_p(\mathcal{F}(\{Y_0, Y_1\})), \\ [L^p(\mathcal{F}(\{X_0, X_1\}))] \sum_{j=1}^{\infty} r_j x_j &\mapsto [L^p(\mathcal{F}(\{Y_0, Y_1\}))] \sum_{j=1}^{\infty} r_j T_j x_j \end{aligned}$$

is a well-defined operator satisfying

$$\|\mathbb{T}\|_{L(\text{Rad}_p(\mathcal{F}(\{X_0, X_1\})), \text{Rad}_p(\mathcal{F}(\{Y_0, Y_1\})))} \leq C_0 \cdot h(\|R^{X_0}\|, \|R^{X_1}\|) \cdot h(\mathcal{R}_p(\mathcal{T}_{|X_0}), \mathcal{R}_p(\mathcal{T}_{|X_1}))$$

with C_0 as given in the statement of the theorem. We define the operator

$$\begin{aligned} \mathbb{S} : \text{Rad}_p(X_0) + \text{Rad}_p(X_1) &\longrightarrow \text{Rad}_p(Y_0) + \text{Rad}_p(Y_1) \\ f = f_0 + f_1 &\mapsto \mathbb{T}_0 f_0 + \mathbb{T}_1 f_1, \end{aligned}$$

with $\mathbb{T}_k : \text{Rad}_p(X_k) \longrightarrow \text{Rad}_p(Y_k)$, $[L^p(X_k)] \sum_{j=1}^{\infty} r_j x_j \mapsto [L^p(Y_k)] \sum_{j=1}^{\infty} r_j T_j x_j$. So we get

$$\tilde{\mathbb{S}} := \mathbb{S}|_{\mathcal{F}(\{\text{Rad}_p(X_0), \text{Rad}_p(X_1)\})} \in L(\mathcal{F}(\{\text{Rad}_p(X_0), \text{Rad}_p(X_1)\}), \mathcal{F}(\{\text{Rad}_p(Y_0), \text{Rad}_p(Y_1)\}))$$

with

$$\begin{aligned} \|\tilde{\mathbb{S}}\|_{L(\mathcal{F}(\{\text{Rad}_p(X_0), \text{Rad}_p(X_1)\}), \mathcal{F}(\{\text{Rad}_p(Y_0), \text{Rad}_p(Y_1)\}))} &\leq C_h \cdot h(\|\mathbb{T}_0\|, \|\mathbb{T}_1\|) \\ &\leq C_h \cdot h(\mathcal{R}_p(\mathcal{T}_{|X_0}), \mathcal{R}_p(\mathcal{T}_{|X_1})). \end{aligned}$$

In the last estimate we already used that $\|\mathbb{T}_k\|_{L(\text{Rad}_p(X_k), \text{Rad}_p(Y_k))} \leq \mathcal{R}_p(\mathcal{T}_{|X_k})$ for $k = 0, 1$. Hence we have $\|\tilde{\mathbb{S}}\|_{L(\text{Rad}_p(\mathcal{F}(\{X_0, X_1\})), \text{Rad}_p(\mathcal{F}(\{Y_0, Y_1\})))} \leq \left[\frac{C_2 C_h^3}{C_1} h(\|R^{X_0}\|, \|R^{X_1}\|) \right] \cdot h(\mathcal{R}_p(\mathcal{T}_{|X_0}), \mathcal{R}_p(\mathcal{T}_{|X_1}))$ by Proposition 3.14. Pick $f := [L^p(\mathcal{F}(X_0, X_1))] \sum_{j=1}^{\infty} r_j x_j \in \text{Rad}_p(\mathcal{F}(\{X_0, X_1\}))$. Then we have

$$f = [L^p(X_0 + X_1)] \sum_{j=1}^{\infty} r_j x_j \in \text{Rad}_p(X_0 + X_1).$$

In view of $\text{Rad}_p(\mathcal{F}(\{X_0, X_1\})) = \mathcal{F}(\{\text{Rad}_p(X_0), \text{Rad}_p(X_1)\}) \subset \text{Rad}_p(X_0) + \text{Rad}_p(X_1)$ we also have

$$\begin{aligned} f = f_0 + f_1 &= [L^p(X_0)] \sum_{j=1}^{\infty} r_j (x_j)_0 + [L^p(X_1)] \sum_{j=1}^{\infty} r_j (x_j)_1 \\ &= [L^p(X_0 + X_1)] \sum_{j=1}^{\infty} r_j [(x_j)_0 + (x_j)_1] \in \text{Rad}_p(X_0 + X_1) \end{aligned}$$

with $x_j = (x_j)_0 + (x_j)_1$ and $(x_j)_k \in X_k$ ($k = 0, 1$). Furthermore, we obtain

$$\begin{aligned} \tilde{\mathbb{S}}f = \mathbb{T}_0 f_0 + \mathbb{T}_1 f_1 &= [L^p(Y_0)] \sum_{j=1}^{\infty} r_j T_j(x_j)_0 + [L^p(Y_1)] \sum_{j=1}^{\infty} r_j T_j(x_j)_1 \\ &= [L^p(\mathcal{F}(\{Y_0, Y_1\}))] \sum_{j=1}^{\infty} r_j T_j \underbrace{[(x_j)_0 + (x_j)_1]}_{=x_j} \\ &= \mathbb{T}f. \end{aligned}$$

This yields $\mathbb{T} = \tilde{\mathbb{S}}$ which completes the proof. \square

Corollary 3.19. *Let $\{X_0, X_1\}$ and $\{Y_0, Y_1\}$ be interpolation couples of Banach spaces of class \mathcal{HT} . For given $\mathcal{T} \subset L(\{X_0, X_1\}, \{Y_0, Y_1\})$, $p \in (1, \infty)$, and $0 < \theta < 1$ we have:
If $\mathcal{T}|_{X_k} \subset L(X_k, Y_k)$ is \mathcal{R} -bounded with \mathcal{R} -bound $\mathcal{R}_p(\mathcal{T}|_{X_k})$, $k = 0, 1$, then*

$$\begin{aligned} \mathcal{T}|_{(X_0, X_1)_{\theta, p}} &\subset L((X_0, X_1)_{\theta, p}, (Y_0, Y_1)_{\theta, p}), \\ \mathcal{T}|_{[X_0, X_1]_{\theta}} &\subset L([X_0, X_1]_{\theta}, [Y_0, Y_1]_{\theta}) \end{aligned}$$

are also \mathcal{R} -bounded with

$$\begin{aligned} \mathcal{R}_p(\mathcal{T}|_{(X_0, X_1)_{\theta, p}}) &\leq C \cdot [\mathcal{R}_p(\mathcal{T}|_{X_0})]^{1-\theta} [\mathcal{R}_p(\mathcal{T}|_{X_1})]_{\theta}, \\ \mathcal{R}_p(\mathcal{T}|_{[X_0, X_1]_{\theta}}) &\leq C' \cdot [\mathcal{R}_p(\mathcal{T}|_{X_0})]^{1-\theta} [\mathcal{R}_p(\mathcal{T}|_{X_1})]_{\theta}, \end{aligned}$$

and with $C := \frac{C_p^{(2)}}{C_p^{(1)}} \|R^{X_0}\|^{1-\theta} \|R^{X_1}\|_{\theta}$ and $C' := \|R^{X_0}\|^{1-\theta} \|R^{X_1}\|_{\theta}$. The constants $C_p^{(k)}$ are the same as in (1).

3.4 Maximal L^p -regularity, \mathcal{R} -sectoriality, and interpolation

Definition 3.20 (Sectoriality). *A linear, densely defined operator $A : D(A) \subset X \rightarrow X$ is called sectorial, if there exists a $\theta \in (0, \pi]$ such that $\Sigma_{\theta} \subset \rho(A)$ and*

$$\sup_{\lambda \in \Sigma_{\theta}} \|\lambda(\lambda - A)^{-1}\|_{L(X)} < \infty.$$

Here we define $\Sigma_{\theta} := \{z \in \mathbb{C} \setminus \{0\} : |\arg(z)| < \theta\}$ as an open sector.

The number $\varphi(A) := \sup \left\{ \theta \in (0, \pi] : \Sigma_{\theta} \subset \rho(A) \wedge \sup_{\lambda \in \Sigma_{\theta}} \|\lambda(\lambda - A)^{-1}\|_{L(X)} < \infty \right\}$ is called spectral angle of A .

Definition 3.21 (\mathcal{R} -sectoriality). *A linear, densely defined operator $A : D(A) \subset X \rightarrow X$ is called \mathcal{R} -sectorial, if there exists a $\theta \in (0, \pi]$ such that $\Sigma_{\theta} \subset \rho(A)$ and such that*

$$\{\lambda(\lambda - A)^{-1} : \lambda \in \Sigma_{\theta}\} \subset L(X)$$

is \mathcal{R} -bounded. The number $\varphi_{\mathcal{R}}(A) := \sup\{\theta \in (0, \pi] : \Sigma_{\theta} \subset \rho(A) \wedge \{\lambda(\lambda - A)^{-1} : \lambda \in \Sigma_{\theta}\} \mathcal{R}\text{-bounded}\}$ is called \mathcal{R} -angle of A .

Observe that in view of Remark 3.6 \mathcal{R} -sectoriality implies sectoriality and we always have $\varphi_{\mathcal{R}}(A) \geq \varphi(A)$. We can now apply the results obtained in the previous sections to conclude that \mathcal{R} -sectoriality is preserved under interpolation.

Theorem 3.22 (\mathcal{R} -sectoriality and interpolation). *Let $p \in (1, \infty)$ and \mathcal{F} be an arbitrary L^p -compatible interpolation functor of type h . Let X_0, X_1 be Banach spaces of class \mathcal{HT} such that $X_0 \cap X_1 \hookrightarrow_d \mathcal{F}(\{X_0, X_1\})$. Furthermore, let*

$$\begin{aligned} A_0 &: D(A_1) \subset X_0 \rightarrow X_0, \\ A_1 &: D(A_2) \subset X_1 \rightarrow X_1, \end{aligned}$$

be two given operators such that $D(A_0) \cap D(A_1) \hookrightarrow_d X_0 \cap X_1$ and $A_0 u = A_1 u$ for all $u \in D(A_0) \cap D(A_1)$. If A_0 and A_1 are \mathcal{R} -sectorial, then the operator

$$B : D(B) \subset \mathcal{F}(\{X_0, X_1\}) \rightarrow \mathcal{F}(\{X_0, X_1\}), \quad D(B) := \mathcal{F}(\{D(A_0), D(A_1)\})$$

with $Bu := A_0 u_0 + A_1 u_1$ for $u = u_0 + u_1 \in D(B) \hookrightarrow D(A_0) + D(A_1)$ is also \mathcal{R} -sectorial. Note, that $D(A_k)$ is equipped with the graph norm $\|\cdot\|_{A_k}$. Moreover, we have $\varphi_{\mathcal{R}}(B) \geq \min_{k=0,1} \varphi_{\mathcal{R}}(A_k)$.

Proof. The operator B is densely defined because of $D(A_0) \cap D(A_1) \hookrightarrow_d X_0 \cap X_1 \hookrightarrow_d \mathcal{F}(\{X_0, X_1\})$ and $D(A_0) \cap D(A_1) \hookrightarrow \mathcal{F}(\{D(A_0), D(A_1)\}) \hookrightarrow \mathcal{F}(\{X_0, X_1\})$.

First we consider the relation of the resolvents of A_0, A_1 , and B . Let $\lambda \in \rho(A_0) \cap \rho(A_1)$ then we can define

$$R_\lambda : X_0 + X_1 \rightarrow D(A_0) + D(A_1), \quad x_0 + x_1 \mapsto (\lambda - A_0)^{-1} x_0 + (\lambda - A_1)^{-1} x_1$$

and get

$$[R_\lambda]_{X_k} = (\lambda - A_k)^{-1}, \quad [R_\lambda]_{\mathcal{F}(\{X_0, X_1\})} \in L(\mathcal{F}(\{X_0, X_1\}), \mathcal{F}(\{D(A_0), D(A_1)\})).$$

With this we can prove $\lambda \in \rho(B)$ and $(\lambda - B)^{-1} = [R_\lambda]_{\mathcal{F}(\{X_0, X_1\})}$. For $0 < \theta < \min_{k=1,2} \varphi_{\mathcal{R}}(A_k)$ we have $\Sigma_\theta \subset \rho(A_1) \cap \rho(A_2)$ and therefore also $\Sigma_\theta \subset \rho(B)$.

By assumption the families

$$\mathcal{T}_0 := \{[R_\lambda]_{X_0} : \lambda \in \Sigma_\theta\} \subset L(X_0), \quad \mathcal{T}_1 := \{[R_\lambda]_{X_1} : \lambda \in \Sigma_\theta\} \subset L(X_1)$$

are \mathcal{R} -bounded. Thus Theorem 3.18 yields the \mathcal{R} -boundedness of

$$\{\lambda(\lambda - B)^{-1} : \lambda \in \Sigma_\theta\} = \{[R_\lambda]_{\mathcal{F}(\{X_0, X_1\})} : \lambda \in \Sigma_\theta\} \subset L(\mathcal{F}(\{X, Y\})).$$

This proves the \mathcal{R} -sectoriality of B with $\varphi_{\mathcal{R}}(B) \geq \min_{k=1,2} \varphi_{\mathcal{R}}(A_k)$. \square

Remark 3.23. *The characterization of maximal L^p -regularity by \mathcal{R} -sectoriality with \mathcal{R} -angle bigger than $\frac{\pi}{2}$ allows for corresponding results on maximal L^p -regularity. The characterization mentioned above can e.g. be found in [14].*

Remark 3.24. *The results of Theorem 3.22 hold for interpolation functors of the real and the complex method, by the fact that they are L^p -compatible and since we always have $X_0 \cap X_1 \hookrightarrow_d \mathcal{F}(\{X_0, X_1\})$. A proof of the density of the last embedding can be found in [18, 1.6.2, 1.9.3], for example.*

4 Property (α)

Our first aim in this section is to interpret property (α) as a special form of \mathcal{R} -boundedness. With the help of this interpretation we will show that also property (α) carries over to interpolation spaces provided that the interpolated Banach spaces are of class \mathcal{HT} .

4.1 Fundamental facts about property (α)

First we recall the definition of property (α) from [14, Section 4.9]. This property is important in the context of an operator valued Fourier-multiplier theorem proved by L.Weis. cf. [14, Section 5.2 or Theorem 4.13]. Another application can be found in [14, 12.8 Theorem], where the authors proved, that the bounded H^∞ -calculus is equivalent to the a priori stronger property of an \mathcal{R} -bounded H^∞ -calculus, if the underlying Banach space has property (α) . The H^∞ -calculus is a powerful tool in the treatment of parabolic and elliptic partial differential equations. For more information on this topic we refer to [5] and [9], for instance.

Definition 4.1 (Property (α)). A Banach space X has property (α) if there exists a constant $C > 0$ such that for all $n \in \mathbb{N}$, $(\alpha_{ij})_{i,j=1,\dots,n} \subset \mathbb{C}$, $|\alpha_{ij}| \leq 1$, and all $(x_{ij})_{i,j=1,\dots,n} \subset X$ we have that

$$\int_0^1 \int_0^1 \left\| \sum_{i,j=1}^n r_i(u)r_j(v)\alpha_{ij}x_{ij} \right\|_X dudv \leq C \int_0^1 \int_0^1 \left\| \sum_{i,j=1}^n r_i(u)r_j(v)x_{ij} \right\|_X dudv. \quad (5)$$

In this case we set $C_\alpha := \min\{C > 0 : \text{estimate (5) holds}\}$.

By virtue of the following Lemma 4.3 (ii) with $p = 2$ and the orthogonality of the Rademacher functions we see that Hilbert spaces have property(α). Let X be a Banach space with property(α) then every closed subspace $Y \subset X$ has property(α). The cartesian product of Banach spaces with property(α) has also property(α). These results can be used to show that the Sobolev space $W^{m,p}(\Omega, X)$ has property(α) for $1 \leq p < \infty$ and $m \in \mathbb{N}_0$ if X possess property(α). This follows easily from the theorem of Tonelli for $m = 0$ and the fact that $W^{m,p}(\Omega, X)$ ($m > 0$) is isometric isomorphic to a closed subspace of $(L^p(\Omega, X))^N$.

Definition 4.2. We set

$$\begin{aligned} T_\alpha^m : \text{Rad}_p^m(X) &\rightarrow \text{Rad}_p^m(X) \\ (x_i)_{i=1,\dots,m} &\mapsto (\alpha_i x_i)_{i=1,\dots,m} \end{aligned}$$

for $m \in \mathbb{N}$ and $\alpha \in \mathbb{C}^m$. Additionally, we define the family

$$\mathcal{T}^m := \{T_\alpha^m : \alpha \in \mathbb{C}^m, |\alpha_i| \leq 1, i = 1, \dots, m\}.$$

Lemma 4.3 (Equivalent characterization of property (α)). Let X be a Banach space and $p \in [1, \infty)$. We have the following equivalences:

(i) X has property (α).

(ii) (p -independence) There exists a constant $\tilde{C} > 0$ such that for all $n \in \mathbb{N}$, $(\alpha_{ij})_{i,j=1,\dots,n} \subset \mathbb{C}$, $|\alpha_{ij}| \leq 1$ and all $(x_{ij})_{i,j=1,\dots,n} \subset X$ we have

$$\left(\int_0^1 \int_0^1 \left\| \sum_{i,j=1}^n r_i(u)r_j(v)\alpha_{ij}x_{ij} \right\|_X^p dudv \right)^{1/p} \leq \tilde{C} \left(\int_0^1 \int_0^1 \left\| \sum_{i,j=1}^n r_i(u)r_j(v)x_{ij} \right\|_X^p dudv \right)^{1/p}. \quad (6)$$

(iii) There exists a $C > 0$ such that we have $\mathcal{R}_p(\mathcal{T}^m) \leq C$ in $L(\text{Rad}_p^m(X))$ for all $m \in \mathbb{N}$.

Proof. The equivalence of (i) and (ii) is an immediate consequence of the inequality of Kahane (Theorem 3.4). In fact, it yields

$$\left[C_p^{(K)} \right]^{-2} \|(\xi_k)_k\|_{\text{Rad}_1^m(\text{Rad}_1^m(X))} \leq \|(\xi_k)_k\|_{\text{Rad}_p^m(\text{Rad}_p^m(X))} \leq \left[C_p^{(K)} \right]^2 \|(\xi_k)_k\|_{\text{Rad}_1^m(\text{Rad}_1^m(X))}$$

for $(\xi_k)_k \in \text{Rad}_1^m(\text{Rad}_1^m(X))$.

“(ii) \Rightarrow (iii)”: Here we can use the characterization of \mathcal{R} -boundedness by Remark 3.7 (i). For this purpose, we set $Y_m := \text{Rad}_p^m(X)$ ($m \in \mathbb{N}$), choose arbitrary $(T_{\alpha^{(j)}}^m)_{j \in \mathbb{N}} \subset \mathcal{T}^m$, and define the operator

$$\mathbb{T}_n : \text{Rad}_p^n(Y_m) \rightarrow \text{Rad}_p^n(Y_m), \quad (x_j)_{j=1,\dots,n} \mapsto (T_{\alpha^{(j)}}^m x_j)_{j=1,\dots,n}$$

with $\alpha^{(j)} = (\alpha_{ij})_{i=1,\dots,m} \in \mathbb{C}^m$, $x_j = (x_{ij})_{i=1,\dots,m} \in Y_m$. Under use of (ii) we get

$$\|\mathbb{T}_n(x_j)_j\|_{\text{Rad}_p^n(Y_m)} \leq \tilde{C} \|(x_j)_j\|_{\text{Rad}_p^n(Y_m)}$$

for all $n, m \in \mathbb{N}$ and $(x_j)_j \in \text{Rad}_p^n(Y_m)$. So, we have $\|\mathbb{T}_n\|_{L(\text{Rad}_p^n(Y_m))} \leq \tilde{C}$ for all $n, m \in \mathbb{N}$. Now Remark 3.7 implies (iii).

“(iii) \Rightarrow (ii)”: Can be done in an analogous way. \square

4.2 Property (α) and interpolation

Lemma 4.4. *Let X be a Banach space of class \mathcal{HT} and $Y \subset X$ be a Banach space such that $\|\cdot\|_Y = \|\cdot\|_X$ on Y . Then we have*

$$(i) \quad \|R_m^X\|_{L(L^p([0,1],X))} \leq 2\|R^X\|_{L(L^p([0,1],X))}, \quad m \in \mathbb{N},$$

$$(ii) \quad \|R^Y\|_{L(L^p([0,1],Y))} \leq \|R^X\|_{L(L^p([0,1],X))}$$

for all $p \in (1, \infty)$.

Proof. This follows easily by the contraction principle of Kahane (Theorem 3.5). \square

Theorem 4.5 (Property (α) and interpolation). *Let $\{X, Y\}$ be an interpolation couple of Banach spaces of class \mathcal{HT} and $p \in (1, \infty)$ and let \mathcal{F} be an L^p -compatible interpolation functor of type h . If X and Y have property (α) with constants $C_\alpha^X > 0$ and $C_\alpha^Y > 0$, then also the interpolation space $\mathcal{F}(\{X, Y\})$ has property (α) with*

$$C_\alpha^{\mathcal{F}(\{X, Y\})} \leq M_0 \cdot h(M_1 C_\alpha^X, M_1 C_\alpha^Y)$$

for some constants $M_0, M_1 > 0$.

Proof. The family $\mathcal{T}^m := \{T_\alpha^m : \alpha \in \mathbb{C}^m, |\alpha_i| \leq 1, i = 1, \dots, m\}$ can easily be interpreted as an subset of $L(\{\text{Rad}_p^m(X), \text{Rad}_p^m(Y)\})$. Then we define $\mathcal{T}_Z^m := \mathcal{T}_{\text{Rad}_p^m(Z)}^m$ for $Z \in \{X, Y, \mathcal{F}(\{X, Y\})\}$. Thanks to Lemma 4.3 we already know, that the families $\mathcal{T}_X^m \subset L(\text{Rad}_p^m(X))$ and $\mathcal{T}_Y^m \subset L(\text{Rad}_p^m(Y))$ are \mathcal{R} -bounded uniformly in $m \in \mathbb{N}$ with

$$\mathcal{R}_p(\mathcal{T}_X^m) \leq \left[C_p^{(K)}\right]^4 C_\alpha^X \quad \text{and} \quad \mathcal{R}_p(\mathcal{T}_Y^m) \leq \left[C_p^{(K)}\right]^4 C_\alpha^Y.$$

Therefore we obtain the \mathcal{R} -boundedness of

$$[\mathcal{T}^m]_{|\mathcal{F}(\{\text{Rad}_p^m(X), \text{Rad}_p^m(Y)\})} \subset L(\mathcal{F}(\{\text{Rad}_p^m(X), \text{Rad}_p^m(Y)\}))$$

by Theorem 3.18 and Remark 3.3. Additionally, this leads to an estimate of the \mathcal{R} -bound

$$\mathcal{R}_p\left([\mathcal{T}^m]_{|\mathcal{F}(\{\text{Rad}_p^m(X), \text{Rad}_p^m(Y)\})}\right) \leq C^{(m)} \cdot h(\mathcal{R}_p(\mathcal{T}_X^m), \mathcal{R}_p(\mathcal{T}_Y^m))$$

with $C^{(m)} := \frac{C_2 C_h^3}{C_1} \cdot h\left(\|R^{\text{Rad}_p^m(X)}\|, \|R^{\text{Rad}_p^m(Y)}\|\right)$. Proposition 3.15 yields

$$[\mathcal{T}^m]_{|\mathcal{F}(\{\text{Rad}_p^m(X), \text{Rad}_p^m(Y)\})} = [\mathcal{T}^m]_{|\text{Rad}_p^m(\mathcal{F}(\{X, Y\}))} \subset L(\text{Rad}_p^m(\mathcal{F}(\{X, Y\}))).$$

Thus we can consider the \mathcal{R} -bound of \mathcal{T}^m in $L(\text{Rad}_p^m(\mathcal{F}(\{X, Y\})))$ and obtain

$$\begin{aligned} \mathcal{R}_p\left([\mathcal{T}^m]_{|\text{Rad}_p^m(\mathcal{F}(\{X, Y\}))}\right) &\leq \left[\frac{C_2 C_h^2}{C_1} h(\|R_m^X\|, \|R_m^Y\|)\right] \cdot \mathcal{R}_p\left([\mathcal{T}^m]_{|\mathcal{F}(\{\text{Rad}_p^m(X), \text{Rad}_p^m(Y)\})}\right) \\ &\leq \underbrace{\left[\frac{C_2 C_h^2}{C_1} h(2\|R^X\|, 2\|R^Y\|)\right]}_{=: C'} \cdot \mathcal{R}_p\left([\mathcal{T}^m]_{|\mathcal{F}(\{\text{Rad}_p^m(X), \text{Rad}_p^m(Y)\})}\right) \end{aligned}$$

by Lemma 4.4 (i). By virtue of $\text{Rad}_p^m(X) \subset \text{Rad}_p(X)$, $\text{Rad}_p^m(Y) \subset \text{Rad}_p(Y)$, Remark 3.3 and Lemma 4.4 (ii) we have

$$\|R^{\text{Rad}_p^m(Z)}\|_{L(L^p([0,1], \text{Rad}_p^m(Z)))} \leq \|R^{\text{Rad}_p(Z)}\|_{L(L^p([0,1], \text{Rad}_p(Z)))} < \infty \quad (m \in \mathbb{N})$$

for $Z \in \{X, Y\}$. Hence there exists an upper bound for $(C^{(m)})_{m \in \mathbb{N}}$:

$$C^{(m)} = \frac{C_2 C_h^3}{C_1} h\left(\|R^{\text{Rad}_p^m(X)}\|, \|R^{\text{Rad}_p^m(Y)}\|\right) \leq \frac{C_2 C_h^3}{C_1} h\left(\|R^{\text{Rad}_p(X)}\|, \|R^{\text{Rad}_p(Y)}\|\right) =: C'' \quad (m \in \mathbb{N}).$$

Summarizing results in

$$\mathcal{R}_p([\mathcal{T}^m]_{|\text{Rad}_p^m(\mathcal{F}(\{X,Y\}))}) \leq M_0 \cdot h(M_1 C_\alpha^X, M_1 C_\alpha^Y)$$

with $M_0 := C' C'' [C_p^{(K)}]^4$ and $M_1 := [C_p^{(K)}]^4$. Due to Lemma 4.3 the assertion follows by the proved \mathcal{R} -boundedness in $\text{Rad}_p^m(\mathcal{F}(\{X, Y\}))$. \square

Corollary 4.6. *Let $\{X, Y\}$ be an interpolation couple of Banach spaces of class \mathcal{HT} . If X and Y have property (α) with $C_\alpha^X > 0$ and $C_\alpha^Y > 0$, then the real and complex interpolation spaces $(X, Y)_{\theta, p}$ and $[X, Y]_\theta$ also have property (α) for $p \in (1, \infty)$, $\theta \in (0, 1)$. In this case we have*

$$\begin{aligned} C_\alpha^{(X,Y)_{\theta,p}} &\leq M' [C_\alpha^X]^{1-\theta} [C_\alpha^Y]^\theta, \\ C_\alpha^{[X,Y]_\theta} &\leq M'' [C_\alpha^X]^{1-\theta} [C_\alpha^Y]^\theta \end{aligned}$$

for some constants $M', M'' > 0$.

5 Application to parabolic systems

In this chapter we consider realizations of parabolic differential equation systems in higher order spaces over \mathbb{R}^n . For example we define the Laplace operator on Sobolev spaces as

$$A_{k,p} : D(A_{k,p}) \subset W^{k,p}(\mathbb{R}^n) \rightarrow W^{k,p}(\mathbb{R}^n), \quad f \mapsto \Delta f,$$

with $D(A_{k,p}) := W^{k+2,p}(\mathbb{R}^n)$ ($k \in \mathbb{N}_0$). Similarly, we can define realizations on interpolation spaces such as Besov and Bessel-potential spaces. In the following we show that the realizations of parabolic systems on Sobolev spaces are \mathcal{R} -sectorial. The case $k = 0$ is proved in [14]. Here we generalize this result to the case $k \in \mathbb{N}$. Basically we follow the proof given in [14] for the case $k = 0$. However, due to the lack of differentiability of some cut-off functions used in [14], here we are forced to adapt the localization procedure at some places suitably. Applying the interpolation results of the previous chapters, we then will obtain \mathcal{R} -sectoriality for realizations on certain scales of interpolation spaces. To handle the parabolic problems under consideration we make use of Fourier multiplier methods. For this purpose and for the definition of Bessel-potential-spaces, here we recall the notion of a Fourier multiplier.

Definition 5.1 (Operator-valued Fourier-multiplier). *Let X, Y be Banach spaces, $1 < p < \infty$, and $m \in L^\infty(\mathbb{R}^n, L(X, Y))$. Then we define*

$$\begin{aligned} T_m : \mathcal{S}(\mathbb{R}^n, X) &\rightarrow L^\infty(\mathbb{R}^n, Y) \\ f &\mapsto \mathcal{F}^{-1} m \mathcal{F} f. \end{aligned}$$

The symbol m is said to be an L^p -Fourier-multiplier, if there exists a $C_p > 0$ such that

- (i) $T_m f \in L^p(\mathbb{R}^n, Y)$ for all $f \in \mathcal{S}(\mathbb{R}^n, X)$,
- (ii) $\|T_m f\|_{L^p(\mathbb{R}^n, Y)} \leq C_p \|f\|_{L^p(\mathbb{R}^n, X)}$ for all $f \in \mathcal{S}(\mathbb{R}^n, X)$.

In this case there exists a unique continuous extension of T_m from $L^p(\mathbb{R}^n, X)$ to $L^p(\mathbb{R}^n, Y)$ which, for simplicity, is also denoted by T_m .

Remark 5.2. *Let X be a Banach space of class \mathcal{HT} and $k \in \mathbb{N}$. We define $\langle \xi \rangle := (1 + |\xi|^2)^{1/2}$ and*

$$\begin{aligned} \Lambda_{-k} : L^p(\mathbb{R}^n, X) &\rightarrow W^{k,p}(\mathbb{R}^n, X); & f &\mapsto T_{\langle \xi \rangle^{-k} \text{id}_X} f \\ \Lambda_k : W^{k,p}(\mathbb{R}^n, X) &\rightarrow L^p(\mathbb{R}^n, X); & f &\mapsto \mathcal{F}^{-1} \left[\langle \xi \rangle^k \text{id}_X \right] \mathcal{F} f. \end{aligned}$$

By standard arguments we get $[\Lambda_{-k}]_{|W^{j,p}(\mathbb{R}^n, X)} \in L(W^{j,p}(\mathbb{R}^n, X), W^{j+k,p}(\mathbb{R}^n, X))$ for $j \in \mathbb{N}_0$ and $(\Lambda_{-k})^{-1} = \Lambda_k$.

5.1 Besov- and Bessel-potential-spaces

Next we recall some basic facts on Besov- and Bessel-potential spaces, which can be found e.g. in [1, 7.30-7.34] and [18, 2.3-2.4]. The Banach space-valued case can be found in [4].

Definition and Remark 5.3 (Results on Besov and Bessel-potential spaces). *Let X be a Banach space and $1 < p, q < \infty$. Then*

- (i) $B_{p,q}^s(\mathbb{R}^n, X) := (L^p(\mathbb{R}^n, X), W^{k,p}(\mathbb{R}^n, X))_{\frac{s}{k}, q}$ with $s \in (0, \infty) \cap [k-1, k)$ and equipped with the interpolation norm is called Besov space,
- (ii) $H^{s,p}(\mathbb{R}^n, X) := \{u \in \mathcal{S}'(\mathbb{R}^n, X) : \mathcal{F}^{-1} \langle \xi \rangle^s \mathcal{F}u \in L^p(\mathbb{R}^n, X)\}$ with $s \in \mathbb{R}_{\geq 0}$ is called Bessel-potential space, where the norm is given by $\|u\|_{H^{s,p}(\mathbb{R}^n, X)} := \|\mathcal{F}^{-1} \langle \xi \rangle^s \mathcal{F}u\|_{L^p(\mathbb{R}^n, X)}$.

Note that we also have the following representations.

- (iii) Let $m, s, j \in \mathbb{N}_0$ with $0 \leq m < s < j$ and $\lambda \in (0, 1)$ such that $s = (1 - \lambda)m + \lambda j$. Then we have

$$B_{p,q}^s(\mathbb{R}^n, X) = (W^{m,p}(\mathbb{R}^n, X), W^{j,p}(\mathbb{R}^n, X))_{\lambda, q}.$$

- (iv) Let X be of class \mathcal{HT} and $m, j \in \mathbb{N}_0$ with $0 \leq m < s < j$ and $s = (1 - \theta)m + \theta j$ for a $\theta \in (0, 1)$. Then we have:

$$\begin{aligned} H^{s,p}(\mathbb{R}^n, X) &= [W^{m,p}(\mathbb{R}^n, X), W^{j,p}(\mathbb{R}^n, X)]_{\theta} \\ W^{k,p}(\mathbb{R}^n, X) &= H^{k,p}(\mathbb{R}^n, X), \quad (k \in \mathbb{N}_0). \end{aligned}$$

Corollary 5.4. *Let $s \in \mathbb{R}_{>0}$, $1 < p, q < \infty$, and X be a Banach space of class \mathcal{HT} . By an application of Remark 2.2 and Theorem 4.5 we readily obtain the following results:*

- (i) The spaces $B_{p,q}^s(\mathbb{R}^n, X)$ and $H^{s,p}(\mathbb{R}^n, X)$ are of class \mathcal{HT} , too.
- (ii) If X has property (α) , then the spaces $B_{p,q}^s(\mathbb{R}^n, X)$ and $H^{s,p}(\mathbb{R}^n, X)$ have property (α) , too.

5.2 Parabolic systems of differential equations

In the following we always assume that $1 < p < \infty$. Furthermore, we set

$$A(x, D) := \sum_{|\alpha| \leq m} a_{\alpha}(x) D^{\alpha},$$

for $a_{\alpha} : \mathbb{R}^n \rightarrow \mathbb{C}^{N \times N}$, $m, N \in \mathbb{N}$, and for $D^{\alpha} := (-i)^{|\alpha|} \partial^{\alpha}$. We define the “ $W^{k,p}$ -realization” of the formal differential operator $A(x, D)$ by

$$A_{k,p} : D(A_{k,p}) \subset W^{k,p}(\mathbb{R}^n, \mathbb{C}^N) \rightarrow W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)$$

with $D(A_{k,p}) := W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N)$ and $A_{k,p}f := A(x, D)f$ for all $f \in W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N)$. For an interpolation functor \mathcal{F} we define the “ (\mathcal{F}, k, p) -realization” of $A(x, D)$ by

$$A_{\mathcal{F},k,p} : D(A_{\mathcal{F},k,p}) \subset \mathcal{F}(L^p, W^{k,p}) \rightarrow \mathcal{F}(L^p, W^{k,p}) \quad (7)$$

with

$$\begin{aligned} \mathcal{F}(L^p, W^{k,p}) &:= \mathcal{F}(\{L^p(\mathbb{R}^n, \mathbb{C}^N), W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)\}), \\ D(A_{\mathcal{F},k,p}) &:= \mathcal{F}(\{W^{m,p}(\mathbb{R}^n, \mathbb{C}^N), W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N)\}) \end{aligned}$$

and $A_{\mathcal{F},k,p}g := A(x, D)g$ for all $g \in D(A_{\mathcal{F},k,p})$. In particular, we define the “ L^p -realization” of $A(x, D)$ by $A_p := A_{0,p}$.

In order to obtain well-defined operators we need to assume some regularity for the coefficients. For the well-definedness of the $W^{k,p}$ -realization it is sufficient to assume that

$$a_\alpha \in W^{k,p_\infty}(\mathbb{R}^n, \mathbb{C}^{N \times N}), \quad p_\infty := \begin{cases} p, & kp > n \\ \infty, & \text{otherwise} \end{cases} \quad (8)$$

for all $\alpha \in \mathbb{N}_0^n$ with $|\alpha| \leq m$. Here we define $W^{k,\infty}(\mathbb{R}^n, \mathbb{C}^{N \times N}) := \{f \in L^\infty(\mathbb{R}^n, \mathbb{C}^{N \times N}) : D^\beta f \in L^\infty(\mathbb{R}^n, \mathbb{C}^{N \times N}), |\beta| \leq k\}$. As in [1, 4.39 Theorem] the Sobolev space $W^{k,p}$ has the algebra property in case of $kp > n$. Let $K^* > 0$ such that $\|uv\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} \leq K^* \|u\|_{W^{k,p_\infty}(\mathbb{R}^n, \mathbb{C}^N)} \|v\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)}$ for all $u \in W^{k,p_\infty}(\mathbb{R}^n, \mathbb{C}^N)$ and $v \in W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)$.

Remark 5.5. Let $s \in (0, \infty) \cap [k-1, k)$, $k \in \mathbb{N}$, and $1 < q < \infty$. If \mathcal{F} is the real or the complex interpolation functor, $A_{\mathcal{F},k,p}$ represents the Besov or Bessel-potential realization of $A(x, D)$:

(i) For “ $\mathcal{F} = (\cdot, \cdot)_{\frac{s}{k}, q}$ ” we have

$$A_{s,p,q}^{\mathcal{B}} : D(A_{s,p,q}^{\mathcal{B}}) \subset B_{p,q}^s(\mathbb{R}^n, \mathbb{C}^N) \rightarrow B_{p,q}^s(\mathbb{R}^n, \mathbb{C}^N), \quad D(A_{s,p,q}^{\mathcal{B}}) = B_{p,q}^{s+m}(\mathbb{R}^n, \mathbb{C}^N).$$

for $A_{s,p,q}^{\mathcal{B}} := A_{\mathcal{F},k,p}$.

(ii) For “ $\mathcal{F} = [\cdot, \cdot]_{\frac{s}{k}}$ ” we have

$$A_{s,p}^{\mathcal{H}} : D(A_{s,p}^{\mathcal{H}}) \subset H^{s,p}(\mathbb{R}^n, \mathbb{C}^N) \rightarrow H^{s,p}(\mathbb{R}^n, \mathbb{C}^N), \quad D(A_{s,p}^{\mathcal{H}}) = H^{s+m,p}(\mathbb{R}^n, \mathbb{C}^N).$$

$A_{s,p}^{\mathcal{H}} := A_{\mathcal{F},k,p}$.

Remark 5.6. Our approach to get \mathcal{R} -sectoriality for Besov- and Bessel-potential space realizations is a sort of ‘decent method’. The regularity assumption W^{k,p_∞} for the coefficients, of course, is not optimal for the interpolated operators. On the other hand, notice that by a standard perturbation argument the regularity for the coefficients of the interpolated operators can always be reduced to close to optimal. In [7, Section 5], for instance, this argument is used for a reduction from smooth to Hölder continuous coefficients. However, optimal conditions on the coefficients is none of our purposes here. Therefore we will not carry out this argument in what follows.

Definition 5.7 (Symbol). The symbol of $A(x, D)$ is defined by $a(x, \xi) := \sum_{|\alpha| \leq m} a_\alpha(x) \xi^\alpha$, $(x, \xi \in \mathbb{R}^n)$. The principal part of $A(x, D)$ is defined as $A_0(x, D) := \sum_{|\alpha|=m} a_\alpha(x) D^\alpha$. The principal symbol of $A(x, D)$ is then given by the symbol of the principal part, i.e. by $a_0(x, \xi) := \sum_{|\alpha|=m} a_\alpha(x) \xi^\alpha$ for $(x, \xi) \in \mathbb{R}^n \times \mathbb{R}^n$.

Definition 5.8 (Parameter-ellipticity and parabolicity). Let $A(x, D)$ be the formal differential operator given above.

(i) The operator $A(x, D)$ is said to be parameter-elliptic in $\overline{\Sigma}_\theta$, if there exists a constant $C_P > 0$ such that we have

$$|\det(a_0(x, \xi) - \lambda)| \geq C_P (|\xi|^m + |\lambda|)^N$$

for all $x \in \mathbb{R}^n$ and $(\xi, \lambda) \in (\mathbb{R}^n \times \overline{\Sigma}_\theta) \setminus \{0\}$.

(ii) The operator $A(x, D)$ is called parabolic, if $A(x, D)$ is parameter-elliptic in $\overline{\Sigma}_{\pi/2}$.

Remark 5.9. If we consider a parabolic $A(x, D)$ with bounded coefficients a_α ($|\alpha| = m$), then there exists a $\theta \in (\frac{\pi}{2}, \pi)$ such that $A(x, D)$ is even parameter-elliptic in $\overline{\Sigma}_\theta$.

5.2.1 The model-problem

In this section we consider the model problem, i.e., we assume the matrix-valued coefficients of $A(x, D)$ to be constant and that $A(x, D)$ is just a principal part, that is $A(x, D) = A_0(x, D)$. Hence the formal differential operator is of the form

$$A(D) := \sum_{|\alpha|=m} a_\alpha D^\alpha, \text{ with } a_\alpha \in \mathbb{C}^{N \times N}. \quad (9)$$

Lemma 5.10. *Let $v : \mathbb{R}^n \rightarrow \mathbb{C}$ be a function such that $\frac{1}{v} \in C^{|\alpha|}(\mathbb{R}^n)$ and $\alpha \in \{0, 1\}^n$. We set*

$$M(\alpha) := \left\{ (\beta_1, \dots, \beta_{|\alpha|}) \in (\mathbb{N}_0^n)^{|\alpha|} : \sum_{k=1}^{|\alpha|} \beta_k = \alpha \right\}.$$

Then it is easily seen, that there are constants $C(\beta_1, \dots, \beta_{|\alpha|}) \in \mathbb{Z}$ for $(\beta_1, \dots, \beta_{|\alpha|}) \in M(\alpha)$ satisfying $|C(\beta_1, \dots, \beta_{|\alpha|})| \leq |\alpha|!$ and such that we have

$$D^\alpha \frac{1}{v} = \frac{1}{v^{|\alpha|+1}} \sum_{(\beta_1, \dots, \beta_{|\alpha|}) \in M(\alpha)} C(\beta_1, \dots, \beta_{|\alpha|}) \prod_{k=1}^{|\alpha|} (D^{\beta_k} v).$$

Proof. This result follows by induction over $|\alpha|$. □

The next result is proved in [14, 6.2 Theorem] for the case $|\beta| = m$. We need the following extension in order to handle diagonal operators in a suitable way.

Lemma 5.11 (Fourier-multipliers and parameter-elliptic symbols). *Let $A(D)$ be given as in (9) and assume that it is parameter-elliptic in $\overline{\Sigma_\theta}$ for a $\theta \in (0, \pi)$ and a constant $C_P > 0$. Furthermore, let $M > 0$ such that $\sum_{|\alpha|=m} \|a_\alpha\|_{\mathbb{C}^{N \times N}} \leq M$, and for $|\beta| \leq m$ let*

$$m_\beta : (\mathbb{R}^n \times \overline{\Sigma_\theta}) \setminus \{0\} \rightarrow \mathbb{C}^{N \times N}, \quad (\xi, \lambda) \mapsto \xi^\beta \lambda^{\frac{m-|\beta|}{m}} (\lambda - a_0(\xi))^{-1}.$$

Then we have

- (i) *that $m_\beta(\cdot, \lambda)$ is a Fourier multiplier for all $\lambda \in \overline{\Sigma_\theta} \setminus \{0\}$,*
- (ii) *that the families*

$$\begin{aligned} \mathcal{T}_\beta &:= \{T_{m_\beta(\cdot, \lambda)} : \lambda \in \overline{\Sigma_\theta} \setminus \{0\}\} \subset L(L^p(\mathbb{R}^n, \mathbb{C}^N)) \\ \mathcal{T}_{\beta, 0} &:= \{T_{m_\beta(\cdot, \lambda)} : \lambda \in \overline{\Sigma_\theta}\} \subset L(L^p(\mathbb{R}^n, \mathbb{C}^N)) \end{aligned}$$

are \mathcal{R} -bounded. Moreover, the \mathcal{R}_p -bounds of \mathcal{T}_β and $\mathcal{T}_{\beta, 0}$ can be estimated from above by a constant only depending on p, n, m, N, M and C_P .

- (iii) *For every $k \in \mathbb{N}$ we also have the \mathcal{R} -boundedness of*

$$\mathcal{T}_\beta^k := \{[T_{m_\beta(\cdot, \lambda)}]_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} : \lambda \in \overline{\Sigma_\theta} \setminus \{0\}\} \subset L(W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)),$$

and again $\mathcal{R}_p(\mathcal{T}_\beta^k)$ is bounded from above by a constant only depending on k, p, n, m, N, M and C_P .

Proof. Let $\beta \in \mathbb{N}_0^n$, $|\beta| \leq m$, and define the function

$$\tilde{m}_\beta : (\mathbb{R}^n \times \overline{\Sigma_{\theta/m}}) \setminus \{0\} \rightarrow \mathbb{C}^{N \times N}, \quad (\xi, q) \mapsto \xi^\beta q^{m-|\beta|} (q^m - a_0(\xi))^{-1}.$$

By the homogeneity of \tilde{m}_β and the version of Michlins multiplier theorem given by [14, Theorem 5.2 b)], we obtain the \mathcal{R} -boundedness of $\{T_{\tilde{m}_\beta(\cdot, q)} : q \in \overline{\Sigma_{\theta/m}}\} \subset L(L^p(\mathbb{R}^n, \mathbb{C}^N))$. Thus, it remains to prove

an estimate for the \mathcal{R} -bound as asserted. To this end, we derive explicit estimates for $\xi^\alpha D^\alpha m_{\beta,\lambda}(\xi)$, $\alpha \in \{0,1\}^n$, and where $m_{\beta,\lambda} := \widehat{m}_\beta(\cdot, \lambda^{1/m})$. Here we use the representation of the inverse matrix $(\lambda - a_0(\xi))^{-1}$ by the adjugate $(\lambda - a_0(\xi))^\# := ((-1)^{i+j} \cdot \det(\lambda - a_0(\xi))_{j,i}^+)$, where $(\lambda - a_0(\xi))_{j,i}^+$ is the $(N-1) \times (N-1)$ -matrix that results from deleting row j and column i in $(\lambda - a_0(\xi))$. Then we obtain

$$\xi^\alpha D^\alpha m_{\beta,\lambda}(\xi) = \lambda^{\frac{m-|\beta|}{m}} \sum_{\gamma \leq \alpha} \left(\xi^{\alpha-\gamma} D^{\alpha-\gamma} \frac{\xi^\beta}{\det(\lambda - a_0(\xi))} \right) (\xi^\gamma D^\gamma (\lambda - a_0(\xi))^\#) \quad (10)$$

by the Leibniz rule. The equivalence of norms in $\mathbb{C}^{N \times N}$ yields

$$\|\xi^\gamma D^\gamma (\lambda - a_0(\xi))^\#\|_{\mathbb{C}^{N \times N}} \leq C(N) \cdot \sum_{i,j=1}^N |\xi^\gamma D^\gamma \det(\lambda - a_0(\xi))_{j,i}^+|.$$

For every $1 \leq i, j \leq N$ the Leibniz formula for the determinant implies

$$|\xi^\gamma D^\gamma \det(\lambda - a_0(\xi))_{j,i}^+| \leq \sum_{\sigma \in \mathcal{S}_{N-1}} \left| \xi^\gamma D^\gamma \prod_{k=1}^{N-1} [(\lambda - a_0(\xi))_{j,i}^+]_{k,\sigma(k)} \right|.$$

Since $[(\lambda - a_0(\xi))_{j,i}^+]_{k,\sigma(k)}$ is just a component of the matrix $\lambda - a_0(\xi)$ and by $|\xi^\beta D^\beta \xi^\alpha| \leq |\xi^\alpha|$ and $|\xi^\beta D^\beta \lambda| \leq |\lambda|$, we obtain

$$|\xi^\beta D^\beta [(\lambda - a_0(\xi))_{j,i}^+]_{k,\sigma(k)}| \leq C(N, M) (|\xi|^m + |\lambda|) \quad (\beta \leq \gamma).$$

Altogether we therefore have

$$\|\xi^\gamma D^\gamma (\lambda - a_0(\xi))^\#\|_{\mathbb{C}^{N \times N}} \leq C(N, M, n) \cdot (|\xi|^m + |\lambda|)^{N-1} \quad (11)$$

for $\gamma \in \{0,1\}^n$. In order to estimate $\xi^{\alpha-\gamma} D^{\alpha-\gamma} \frac{\xi^\beta}{\det(\lambda - a_0(\xi))}$, we apply Lemma 5.10, the parabolicity condition, and again the Leibniz rule to obtain

$$\begin{aligned} \left| \xi^{\alpha-\gamma} D^{\alpha-\gamma} \frac{\xi^\beta}{\det(\lambda - a_0(\xi))} \right| &\leq \sum_{\delta \leq \alpha-\gamma} \left| \xi^{(\alpha-\gamma)-\delta} D^{(\alpha-\gamma)-\delta} \xi^\beta \right| \left| \xi^\delta D^\delta \frac{1}{\det(\lambda - a_0(\xi))} \right| \\ &\leq C(m, n, N, M, C_P) \frac{|\xi|^{|\beta|}}{(|\xi|^m + |\lambda|)^N}. \end{aligned} \quad (12)$$

Now (10), (11), and (12) imply

$$\begin{aligned} \|\xi^\alpha D^\alpha m_{\beta,\lambda}(\xi)\|_{\mathbb{C}^{N \times N}} &\leq C(n, m, N, M, C_P) \frac{(|\lambda|^{1/m})^{m-|\beta|} |\xi|^{|\beta|}}{(|\xi|^m + |\lambda|)} \\ &\leq C(n, m, N, M, C_P) \frac{(|\xi| + |\lambda|^{1/m})^m}{(|\xi|^m + |\lambda|)} \leq C(n, m, N, M, C_P). \end{aligned}$$

Note that the operator $T_{m_\beta(\cdot, \lambda)}$ commutes with Λ_k and therefore

$$[T_{m_\beta(\cdot, \lambda)}]_{|W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} = \Lambda_{-k} T_{m_\beta(\cdot, \lambda)} \Lambda_k.$$

Assertion (iii) now follows immediately from (ii). \square

Proposition 5.12 (Model-problem for $W^{k,p}$). *Let $A(D)$ be given as in (9) and assume that it is parameter-elliptic in $\bar{\Sigma}_\theta$ for a $\theta \in (0, \pi)$ and a constant $C_P > 0$. Furthermore, assume that we have $\sum_{|\alpha|=m} \|a_\alpha\|_{\mathbb{C}^{N \times N}} \leq M$ for an $M > 0$. Then, there hold the following assertions for the $W^{k,p}$ -realization $A_{k,p}$ ($k \in \mathbb{N}_0$) of $A(D)$:*

(i) $\overline{\Sigma}_\theta \setminus \{0\} \subset \rho(A_{k,p})$.

(ii) For every $\beta \in \mathbb{N}_0^n$ with $0 \leq |\beta| \leq m$ we have, that $\{\lambda^{\frac{m-|\beta|}{m}} D^\beta (\lambda - A_{k,p})^{-1} : \lambda \in \overline{\Sigma}_\theta \setminus \{0\}\} \subset L(W^{k,p}(\mathbb{R}^n, \mathbb{C}^N))$ is \mathcal{R} -bounded. In addition, we have $\lambda^{\frac{m-|\beta|}{m}} D^\beta (\lambda - A_{k,p})^{-1} = [T_{m,\beta,\lambda}]_{|W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)}$ for $m_{\beta,\lambda}(\xi) := \lambda^{\frac{m-|\beta|}{m}} \xi^\beta (\lambda - a_0(\xi))^{-1}$ ($\lambda \in \overline{\Sigma}_\theta \setminus \{0\}, \xi \in \mathbb{R}^n$). Furthermore, the \mathcal{R}_p -bound is bounded from above by a constant only depending on k, p, n, m, N, C_P and M . In particular, this yields the \mathcal{R} -sectoriality of $A_{k,p}$ with $\varphi_{\mathcal{R}}(A_{k,p}) \geq \theta$.

(iii) For all $\lambda \in \overline{\Sigma}_\theta \setminus \{0\}$ there exists a constant $C_\lambda(k, p, n, m, N, C_P, M) > 0$ such that

$$\|(\lambda - A_{k,p})^{-1}\|_{L(W^{k,p}(\mathbb{R}^n, \mathbb{C}^N), W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N))} \leq C_\lambda.$$

Proof. For $\lambda \in \overline{\Sigma}_\theta \setminus \{0\}$ and $0 \leq |\beta| \leq m$ we define

$$T_{\beta,\lambda} := [T_{m,\beta,\lambda}]_{|W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} \in L(W^{k,p}(\mathbb{R}^n, \mathbb{C}^N), W^{k+m-|\beta|,p}(\mathbb{R}^n, \mathbb{C}^N)).$$

Then it follows easily $\lambda(\lambda - A_{k,p})^{-1} = T_{0,\lambda}$ or rather $\lambda^{\frac{m-|\beta|}{m}} D^\beta (\lambda - A_{k,p})^{-1} = T_{\beta,\lambda}$. Therefore we proved (i) and

$$\{\lambda^{\frac{m-|\beta|}{m}} D^\beta (\lambda - A_{k,p})^{-1} : \lambda \in \overline{\Sigma}_\theta \setminus \{0\}\} = \mathcal{T}_\beta^k.$$

By Lemma 5.11 (iii) assertion (ii) follows.

To prove (iii), we use $\lambda(\lambda - A_{k,p})^{-1} = T_{0,\lambda}$ and (ii) to obtain the estimate

$$\begin{aligned} \|(\lambda - A_{k,p})^{-1}\|_{L(W^{k,p}(\mathbb{R}^n, \mathbb{C}^N), W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N))} &\leq \frac{C}{|\lambda|} \max_{|\beta| \leq m} \|T_{\xi^\beta m_0, \lambda}\|_{L(L^p(\mathbb{R}^n, \mathbb{C}^N))} \\ &\leq C(\lambda) C(p, n, m, N, C_P, M) =: C_\lambda. \end{aligned}$$

Hence the assertion is proved. \square

By means of interpolation this results extends to the model-problem of the (\mathcal{F}, k, p) -realization of $A(D)$.

Corollary 5.13 (Model-problem for (\mathcal{F}, k, p) -realization). *Let $p, q \in (1, \infty)$ and let \mathcal{F} be an L^q -compatible interpolation functor of type h . If $A(D)$ is parameter-elliptic in $\overline{\Sigma}_\theta$ ($\theta \in (0, \pi)$) with constant $C_P > 0$, then the (\mathcal{F}, k, p) -realization of $A(D)$ is \mathcal{R} -sectorial with $\varphi_{\mathcal{R}}(A_{\mathcal{F},k,p}) \geq \theta$, provided that*

$$W^{k,p}(\mathbb{R}^n, \mathbb{C}^N) \hookrightarrow_d \mathcal{F}(\{L^p(\mathbb{R}^n, \mathbb{C}^N), W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)\}).$$

Proof. For simplicity we set

$$\begin{aligned} X_0 &:= L^p(\mathbb{R}^n, \mathbb{C}^N), & D(A_0) &:= W^{m,p}(\mathbb{R}^n, \mathbb{C}^N), \\ X_1 &:= W^{k,p}(\mathbb{R}^n, \mathbb{C}^N), & D(A_1) &:= W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N). \end{aligned}$$

According to Remark 2.2 the spaces X_0 and X_1 are of class \mathcal{HT} . Setting $A_0 := A_p$ and $A_1 := A_{k,p}$, Proposition 5.12 yields the \mathcal{R} -sectoriality of A_0 and A_1 with $\min_{j=0,1} \varphi_{\mathcal{R}}(A_j) \geq \theta$. Hence we obtain the \mathcal{R} -sectoriality of the (\mathcal{F}, k, p) -realization by Theorem 3.22. Note that we use graph norms on $D(A_j)$ in Theorem 3.22. On the other hand, we have the equivalence of the graph norms and the Sobolev norms by the fact that $A_0 \in L(W^{m,p}(\mathbb{R}^n, \mathbb{C}^N), L^p(\mathbb{R}^n, \mathbb{C}^N))$ and $A_1 \in L(W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N), W^{k,p}(\mathbb{R}^n, \mathbb{C}^N))$. Theorem 3.22 also yields $\varphi_{\mathcal{R}}(A_{\mathcal{F},k,p}) \geq \theta$. \square

Remark 5.14. *In particular, Corollary 5.13 holds for the real and the complex interpolation method. This follows directly from Remark 3.24.*

Corollary 5.15. *The result of Proposition 5.12 and Corollary 5.13 is also true for a $\theta > \frac{\pi}{2}$, if we regard parabolic model problems. This follows immediately from Remark 5.9.*

Example 5.16. *Let $X_s \in \{B_{p,q}^s(\mathbb{R}^n), H^{s,p}(\mathbb{R}^n)\}$. For all $s \in (0, \infty)$, $1 < p, q < \infty$ the Laplace operator $\Delta : D(\Delta) \subset X_s \rightarrow X_s$ with $D(\Delta) := X_{s+2}$ is \mathcal{R} -sectorial on X_s , and we have $\varphi_{\mathcal{R}}(\Delta) = \pi$.*

5.2.2 Perturbation results

To handle the case of slightly varying coefficients we provide suitable perturbation results for \mathcal{R} -sectorial operators. The following notation as well as Theorem 5.20 are taken from [14].

Definition 5.17. Let $A : D(A) \subset X \rightarrow X$ be an operator on a Banach space X . For $1 < p < \infty$ we define:

$$\Theta_{\mathcal{R}}(A) := \{\theta \in (0, \pi) : \Sigma_{\theta} \subset \rho(A) \wedge \{\lambda(\lambda - A)^{-1} \in L(X) : \lambda \in \Sigma_{\theta}\} \text{ } \mathcal{R}\text{-bounded}\}.$$

It is obvious that A is \mathcal{R} -sectorial with $\varphi_{\mathcal{R}}(A) = \sup \Theta_{\mathcal{R}}(A)$ provided that $\Theta_{\mathcal{R}}(A) \neq \emptyset$. For $\theta \in \Theta_{\mathcal{R}}(A)$ we define :

$$\begin{aligned} N_{\theta}(A) &:= \sup\{\|\lambda(\lambda - A)^{-1}\|_{L(X)} : \lambda \in \Sigma_{\theta}\}, & R_{\theta,p}(A) &:= \mathcal{R}_p(\{\lambda(\lambda - A)^{-1} : \lambda \in \Sigma_{\theta}\}), \\ \tilde{N}_{\theta}(A) &:= \sup\{\|A(\lambda - A)^{-1}\|_{L(X)} : \lambda \in \Sigma_{\theta}\}, & \tilde{R}_{\theta,p}(A) &:= \mathcal{R}_p(\{A(\lambda - A)^{-1} : \lambda \in \Sigma_{\theta}\}). \end{aligned}$$

Of course $(0, \varphi_{\mathcal{R}}(A)) \subset \Theta_{\mathcal{R}}(A)$, but $\varphi_{\mathcal{R}}(A) \notin \Theta_{\mathcal{R}}(A)$ in general.

Remark 5.18. It can be easily seen that $\tilde{R}_{\theta,p}(A) \leq 1 + R_{\theta,p}(A)$, $R_{\theta,p}(A) \leq 1 + \tilde{R}_{\theta,p}(A)$, $N_{\theta}(A) \leq R_{\theta,p}(A)$ and $\tilde{N}_{\theta}(A) \leq \tilde{R}_{\theta,p}(A)$.

The next lemma is an obvious consequence of the definition of \mathcal{R} -sectoriality.

Lemma 5.19 (Shift of \mathcal{R} -sectorial operators). Let $A : D(A) \subset X \rightarrow X$ be an \mathcal{R} -sectorial operator and $\theta \in \Theta_{\mathcal{R}}(A)$, $\mu > 0$ be arbitrary. Then we have $\tilde{R}_{\theta,p}(A - \mu) \leq C'_{\theta} + C_{\theta} \tilde{R}_{\theta,p}(A)$ and therefore the \mathcal{R} -sectoriality of $A - \mu : D(A) \subset X \rightarrow X$ with constants $C_{\theta}, C'_{\theta} > 0$ only depending on θ . Furthermore, $\theta \in \Theta_{\mathcal{R}}(A - \mu)$ and $\varphi_{\mathcal{R}}(A - \mu) \geq \theta$.

Theorem 5.20. Let X be a Banach space, $A : D(A) \subset X \rightarrow X$ be an \mathcal{R} -sectorial operator, and $\theta \in \Theta_{\mathcal{R}}(A)$. Assume that $B : D(B) \subset X \rightarrow X$ is an operator satisfying $D(A) \subset D(B)$ and

$$\|Bx\|_X \leq a\|Ax\|_X + b\|x\|_X \quad (x \in D(A))$$

for some $a, b \geq 0$ such that $a < \left(\tilde{N}_{\theta}(A)(C'_{\theta} + C_{\theta} \tilde{R}_{\theta,p}(A))\right)^{-1}$ (C'_{θ}, C_{θ} from Lemma 5.19). Then there is a constant

$$C(a, b, \theta, A) := \frac{bN_{\theta}(A_{k,p})(C'_{\theta} + C_{\theta} \tilde{R}_{\theta,p}(A_{k,p}))}{1 - a\tilde{N}_{\theta}(A_{k,p})(C'_{\theta} + C_{\theta} \tilde{R}_{\theta,p}(A_{k,p}))} > 0$$

such that for all $\mu > C(a, b, \theta, A)$ the operator $A + B - \mu : D(A) \subset X \rightarrow X$ is \mathcal{R} -sectorial with $\theta \in \Theta_{\mathcal{R}}(A + B - \mu)$ and $\varphi_{\mathcal{R}}(A + B - \mu) \geq \theta$. Moreover, for all $\lambda \in \Sigma_{\theta}$ the resolvent is represented through

$$(\lambda - (A - \mu + B))^{-1} = (\lambda + \mu - A)^{-1}(1 - B(\lambda + \mu - A)^{-1})^{-1} \quad (13)$$

and we have $\|(1 - B(\lambda + \mu - A)^{-1})^{-1}\|_{L(X)} \leq \left(1 - \left[a\tilde{N}_{\theta}(A) + b\frac{1}{\mu}N_{\theta}(A)\right] \tilde{R}_{\theta,p}(A - \mu)\right)^{-1}$.

Lemma 5.21. Let $k \in \mathbb{N}$ ($k \neq 0$) and $A(D) = \sum_{|\alpha|=m} a_{\alpha} D^{\alpha}$ be parameter-elliptic in $\overline{\Sigma_{\theta}}$ with constant $C_P > 0$ and constant coefficients $a_{\alpha} \in \mathbb{C}^{N \times N}$ such that $\sum_{|\alpha|=m} \|a_{\alpha}\|_{\mathbb{C}^{N \times N}} \leq M$. Then there exists a constant $C = C(k, m, n, p, N, M, C_P) > 0$ such that

$$\|D^{\alpha} f\|_{W^{k-1,p}(\mathbb{R}^n, \mathbb{C}^N)} \leq C|\lambda|^{-\frac{1}{m}} \|A_{k,p} f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} + C|\lambda|^{1-\frac{1}{m}} \|f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)}$$

for all $f \in W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N)$, $|\alpha| = m$, and $\lambda \in \overline{\Sigma_{\theta}} \setminus \{0\}$.

Proof. Let $f \in W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N)$, $|\alpha| = m$, and $\lambda \in \overline{\Sigma_{\theta}} \setminus \{0\}$ be arbitrary. Choose any $\beta \in \mathbb{N}_0^n$ with $|\beta| = m - 1$ and $\beta_i \leq \alpha_i$ for all $i = 1, \dots, n$. Proposition 5.12 yields $\lambda \in \rho(A_{k,p})$ and therefore

$$D^{\alpha} f = \lambda^{\frac{|\beta|-m}{m}} D^{\alpha-\beta} \left[\lambda^{\frac{m-|\beta|}{m}} D^{\beta} (\lambda - A_{k,p})^{-1} \right] (\lambda - A_{k,p}) f.$$

According to Proposition 5.12 (ii) there exists a constant $C > 0$ only depending on p, n, m, N, M, k and C_P such that $\max_{|\beta|=m-1} \mathcal{R}_p(\mathcal{T}_\beta^k) \leq C$. In view of $|\beta| = m-1$ and the triangle inequality we then deduce

$$\|D^\alpha f\|_{W^{k-1,p}(\mathbb{R}^n, \mathbb{C}^N)} \leq C|\lambda|^{\frac{|\beta|-m}{m}} (|\lambda| \|f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} + \|A_{k,p} f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)}).$$

□

We are now in position to handle perturbations of the principal part of a parameter-elliptic differential operator with constant coefficients.

Notation: In the following context the constants k, m, n, p, N are always fixed. So, we do not mention the dependence of them explicitly.

Proposition 5.22 (Perturbation of the model-problem of a $W^{k,p}$ -realization). *Let $M, C_P, \tau > 0$, $k \in \mathbb{N}_0$, $1 < p < \infty$, and $\theta \in (0, \pi)$. There exist constants $\varepsilon(M, C_P, \theta) > 0$, $K(M, C_P, \theta) > 0$, and $\mu(M, C_P, \tau, \theta) > 0$ such that for all $A(D) := \sum_{|\alpha|=m} a_\alpha D^\alpha$ and $S(x, D) := \sum_{|\alpha|=m} s_\alpha(x) D^\alpha$ with*

(i) $A(D)$ parameter-elliptic in $\bar{\Sigma}_\theta$ with constant C_P and $a_\alpha \in \mathbb{C}^{N \times N}$ such that $\sum_{|\alpha|=m} \|a_\alpha\|_{\mathbb{C}^{N \times N}} \leq M$,

(ii) $s_\alpha \in W^{k,\infty}(\mathbb{R}^n, \mathbb{C}^{N \times N})$ satisfying $\sum_{|\alpha|=m} \|s_\alpha\|_\infty < \varepsilon$

and in the case of $k \neq 0$ additionally that $\max_{0 < |\gamma| \leq k, |\alpha|=m} \|D^\gamma s_\alpha\|_\infty \leq \tau$

we have that

$$R_{\theta,p}(A_{k,p} + S_{k,p} - \mu) \leq K.$$

Thus the $W^{k,p}$ -realization $A_{k,p} + S_{k,p} - \mu$ is \mathcal{R} -sectorial and we have $\theta \in \Theta_{\mathcal{R}}(A_{k,p} + S_{k,p} - \mu)$, i.e., in particular that $\varphi_{\mathcal{R}}(A_{k,p} + S_{k,p} - \mu) \geq \theta$. Furthermore, for all $\lambda \in \Sigma_\theta$ there exists a constant $C_\lambda(M, C_P, \mu) > 0$ such that

$$\|(\lambda - (A_{k,p} - \mu + S_{k,p}))^{-1}\|_{L(W^{k,p}(\mathbb{R}^n, \mathbb{C}^N), W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N))} \leq C_\lambda. \quad (14)$$

Proof. We define $m_\alpha(\xi) := \xi^\alpha a(\xi)^{-1}$ with $|\alpha| = m$. Due to the homogeneity of m_α this symbol is a Fourier multiplier. Note that

$$T_{m_\alpha} A(D)g = T_{m_\alpha} \mathcal{F}^{-1} a \mathcal{F} g = \mathcal{F}^{-1} \xi^\alpha \mathcal{F} g = D^\alpha g. \quad (15)$$

Thanks to $T_{m_\alpha} \in \mathcal{T}_{\alpha,0}$ Lemma 5.11 (ii) yields

$$\max_{|\alpha|=m} \|T_{m_\alpha}\|_{L(L^p(\mathbb{R}^n, \mathbb{C}^N))} \leq \eta \quad (16)$$

for a constant $\eta(M, C_p) > 0$ that does not depend explicitly on the coefficients a_α .

According to Proposition 5.12 we have the \mathcal{R} -sectoriality of $A_{k,p}$ with $\theta \in \Theta_{\mathcal{R}}(A_{k,p})$ and a constant $\mathcal{K}(M, C_P) > 0$ with $\tilde{N}_\theta(A_{k,p}), N_\theta(A_{k,p}), \tilde{R}_{\theta,p}(A_{k,p}), R_{\theta,p}(A_{k,p}) < \mathcal{K}/2$. In the following we aim for an application of Theorem 5.20. Thus we have to show that

$$\|S_{k,p} f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} \leq a \|A_{k,p} f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} + b \|f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} \quad (17)$$

for $a := a(M, C_P, \theta) := (\mathcal{K}(C'_\theta + C_\theta \mathcal{K}))^{-1} < (\tilde{N}_\theta(A_{k,p})(C'_\theta + C_\theta \tilde{R}_{\theta,p}(A_{k,p})))^{-1}$ and a $b(M, C_P, \tau) \geq 0$.

Step 1: Proof of estimate (17).

Here we will only give the proof for the case $k \geq 1$, since the case $k = 0$ is given in [14]:

Let $g \in W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)$ with $|\alpha| = m$. Then we have thanks to assumption (ii) that

$$\begin{aligned} \|s_\alpha g\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} &\leq C \left[\sum_{|\beta| \leq k} \|s_\alpha\|_\infty \|D^\beta g\|_{L^p(\mathbb{R}^n, \mathbb{C}^N)} + \sum_{|\beta| \leq k} \sum_{\gamma < \beta} \|D^{\beta-\gamma} s_\alpha\|_\infty \|D^\gamma g\|_{L^p(\mathbb{R}^n, \mathbb{C}^N)} \right] \\ &\leq C \|s_\alpha\|_\infty \|g\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} + C(\tau) \|g\|_{W^{k-1,p}(\mathbb{R}^n, \mathbb{C}^N)}. \end{aligned} \quad (18)$$

Summing up, we obtain for $f \in W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N)$ that

$$\|S_{k,p}f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} \leq C \max_{|\alpha|=m} \|s_\alpha\|_\infty \sum_{|\alpha|=m} \|D^\alpha f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} + C(\tau) \sum_{|\alpha|=m} \|D^\alpha f\|_{W^{k-1,p}(\mathbb{R}^n, \mathbb{C}^N)}.$$

Now we have to find estimates for $\|D^\alpha f\|_{W^{k-1,p}(\mathbb{R}^n, \mathbb{C}^N)}$ and $\|D^\alpha f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)}$. By applying (15) and (16) we can conclude

$$\|D^\alpha f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} \leq \eta \|A_{k,p}f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)}. \quad (19)$$

For $\|D^\alpha f\|_{W^{k-1,p}(\mathbb{R}^n, \mathbb{C}^N)}$ we have by Lemma 5.21 for $\lambda_0 > 0$ and a constant $C(M, C_P) > 0$ that

$$\|D^\alpha f\|_{W^{k-1,p}(\mathbb{R}^n, \mathbb{C}^N)} \leq C(M, C_P) \lambda_0^{-\frac{1}{m}} \|A_{k,p}f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} + C(M, C_P, \lambda_0) \|f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)}. \quad (20)$$

This results in

$$\begin{aligned} \|S_{k,p}f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} &\leq \left[C(\eta) \max_{|\alpha|=m} \|s_\alpha\|_\infty + C(M, C_P, \tau) \lambda_0^{-\frac{1}{m}} \right] \|A_{k,p}f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} \\ &\quad + C(M, C_P, \tau, \lambda_0) \cdot \|f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)}. \end{aligned} \quad (21)$$

Step 2: Application of Theorem 5.20

We set $\varepsilon(M, C_P, \theta) := \frac{1}{2} \frac{a}{C(\eta)} > 0$ and fix $\lambda_0 > 0$ such that

$$C(M, C_P, \tau) \lambda_0^{-\frac{1}{m}} < \frac{1}{2} a$$

with a as be given before. Since the choice of λ_0 only depends on the variables M, C_P, τ and θ we obtain that $b := C(M, C_P, \tau, \lambda_0)$ does not depend explicitly on the coefficients a_α and s_α . Since $\sum_{|\alpha|=m} \|s_\alpha\|_\infty < \varepsilon$ we obtain

$$\|S_{k,p}f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} \leq a \|A_{k,p}f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} + b \|f\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)}.$$

By the fact that $\mathcal{K}/2 > \tilde{N}_\theta(A_{k,p})$, $\mathcal{K} > N_\theta(A_{k,p})$, $\tilde{R}_\theta(A_{k,p})$ and since

$$\mu(M, C_P, \tau, \theta) := \frac{b\mathcal{K}(C'_\theta + C_\theta\mathcal{K})}{1 - a\frac{\mathcal{K}}{2}(C'_\theta + C_\theta\mathcal{K})} > \frac{bN_\theta(A_{k,p})(C'_\theta + C_\theta\tilde{R}_{\theta,p}(A_{k,p}))}{1 - a\tilde{N}_\theta(A_{k,p})(C'_\theta + C_\theta\tilde{R}_{\theta,p}(A_{k,p}))} > 0,$$

we obtain the \mathcal{R} -sectoriality of $A_{k,p} + S_{k,p} - \mu$ with $\theta \in \Theta_{\mathcal{R}}(A_{k,p} + S_{k,p} - \mu)$. Observe that we have $\mu = 2b\mathcal{K}(C'_\theta + C_\theta\mathcal{K})$. Moreover, Theorem 5.20 yields the estimate

$$\begin{aligned} R_{\theta,p}(A_{k,p} + S_{k,p} - \mu) &\leq \frac{R_{\theta,p}(A_{k,p} - \mu)}{1 - [a\tilde{N}_\theta(A_{k,p}) + b\frac{1}{\mu}N_\theta(A_{k,p})]\tilde{R}_{\theta,p}(A_{k,p} - \mu)} \\ &\leq \frac{R_{\theta,p}(A_{k,p} - \mu)}{1 - \frac{\mathcal{K}}{2}[a + b\frac{1}{\mu}](C'_\theta + C_\theta\mathcal{K})} \\ &\leq 4(1 + C'_\theta + C_\theta\mathcal{K}) =: K(M, C_P, \theta), \end{aligned}$$

where we used that $\tilde{R}_{\theta,p}(A_{k,p} - \mu) \leq C'_\theta + C_\theta\tilde{R}_{\theta,p}(A_{k,p}) \leq C'_\theta + C_\theta\mathcal{K}$ and $R_{\theta,p}(A_{k,p} - \mu) \leq 1 + \tilde{R}_{\theta,p}(A_{k,p} - \mu)$.

The proof of (14) with $\lambda \in \Sigma_\theta$ follows directly from

$$\begin{aligned} \|(1 - S_{k,p}(\lambda + \mu - A_{k,p})^{-1})^{-1}\|_{L(W^{k,p}(\mathbb{R}^n, \mathbb{C}^N))} &\leq \left(1 - \left[a\tilde{N}_\theta(A_{k,p}) + b\frac{1}{\mu}N_\theta(A_{k,p}) \right] \tilde{R}_{\theta,p}(A_{k,p} - \mu) \right)^{-1} \\ &\leq 4 \end{aligned}$$

and the representation of the resolvent in (13) and Proposition 5.12 (iii). \square

5.2.3 Some helpful facts on diagonal-operators

To prove our main theorem for parabolic systems we next establish some facts on diagonal operators. Diagonal operators appear in a natural way during the process of localization. The proof of the following results is rather elementary and therefore omitted.

Lemma 5.23 (Diagonal-operator). *Let $(T_l)_{l \in \mathbb{N}}$ be a sequence of operators on a Banach space X with $D(T_l) := Y$ for another Banach space $Y \hookrightarrow X$. If there additionally hold the conditions*

- (i) $T_l \in L(Y, X)$ for all $l \in \mathbb{N}$,
- (ii) $\sup_{l \in \mathbb{N}} \|T_l\|_{L(Y, X)} < \infty$,

then the diagonal-operator

$$\mathbb{A} : D(\mathbb{A}) \subset \mathbb{X} \rightarrow \mathbb{X}, \quad (u_l)_{l \in \mathbb{N}} \mapsto (T_l u_l)_{l \in \mathbb{N}}$$

with $\mathbb{X} := \ell^p(\mathbb{N}, X)$ and $D(\mathbb{A}) := \mathbb{Y} := \ell^p(\mathbb{N}, Y)$ is well-defined and we have $\mathbb{A} \in L(\mathbb{Y}, \mathbb{X})$.

If the T_l 's are densely defined, then \mathbb{A} is densely defined as well and we have

$$\rho(\mathbb{A}) = \left\{ \lambda \in \bigcap_{l=1}^{\infty} \rho(T_l) : \exists C_\lambda > 0 : \sup_{l \in \mathbb{N}} \|(\lambda - T_l)^{-1}\|_{L(X, Y)} \leq C_\lambda \right\}.$$

Furthermore, we obtain for all $\lambda \in \rho(\mathbb{A})$ and $(u_l)_{l \in \mathbb{N}} \in \mathbb{X}$ the representation

$$(\lambda - \mathbb{A})^{-1}(u_l)_{l \in \mathbb{N}} = ((\lambda - T_l)^{-1}u_l)_{l \in \mathbb{N}}.$$

Lemma 5.24 (\mathcal{R} -sectoriality of diagonal-operators). *Assume that $(T_l)_l$ is a sequence satisfying the conditions of Lemma 5.23. Then the diagonal-operator \mathbb{A} is \mathcal{R} -sectorial with $\theta \in \Theta_{\mathcal{R}}(\mathbb{A})$, if there exists a $\theta \in (0, \pi)$ such that we have:*

- (i) T_l is \mathcal{R} -sectorial with $\theta \in \Theta_{\mathcal{R}}(T_l)$ for all $l \in \mathbb{N}$.
- (ii) There is a $1 < p < \infty$ such that there exists a $K_p > 0$ with $\sup_{l \in \mathbb{N}} R_{\theta, p}(T_l) \leq K_p$.
- (iii) For all $\lambda \in \Sigma_\theta$ there exists a $C_\lambda > 0$ with $\sup_{l \in \mathbb{N}} \|(\lambda - T_l)^{-1}\|_{L(X, Y)} \leq C_\lambda$.

In particular, we have $\varphi_{\mathcal{R}}(\mathbb{A}) \geq \theta$ and $R_{\theta, p}(\mathbb{A}) \leq K_p$ in this case.

Proof. Thanks to Lemma 5.23 and condition (iii), we have $\Sigma_\theta \subset \rho(\mathbb{A})$ and

$$(\lambda - \mathbb{A})^{-1}(u_l)_{l \in \Gamma} = ((\lambda - T_l)^{-1}u_l)_{l \in \Gamma}$$

for all $(u_l)_{l \in \Gamma} \in \mathbb{X}$ and $\lambda \in \Sigma_\theta$.

It remains to prove the \mathcal{R} -boundedness of $\{\lambda(\lambda - \mathbb{A})^{-1} : \lambda \in \Sigma_\theta\}$. Let $M \in \mathbb{N}$, $(\lambda_k)_{k=1, \dots, M} \subset \Sigma_\theta$, and let $(x_k)_{k=1, \dots, M} \subset \mathbb{X}$ with $x_k =: (u_l^{(k)})_{l \in \mathbb{N}}$. Then we obtain

$$\begin{aligned} \left\| \sum_{k=1}^M r_k \lambda_k (\lambda_k - \mathbb{A})^{-1} x_k \right\|_{L^p([0, 1], \mathbb{X})} &= \left(\sum_{l \in \mathbb{N}} \int_0^1 \left\| \sum_{k=1}^M r_k(t) \lambda_k (\lambda_k - T_l)^{-1} u_l^{(k)} \right\|_X^p dt \right)^{1/p} \\ &\leq \left(\sum_{l \in \mathbb{N}} K_p^p \int_0^1 \left\| \sum_{k=1}^M r_k(t) u_l^{(k)} \right\|_X^p dt \right)^{1/p} = K_p \left\| \sum_{k=1}^M r_k x_k \right\|_{L^p([0, 1], \mathbb{X})}. \end{aligned}$$

This implies $\mathcal{R}_p(\{\lambda(\lambda - \mathbb{A})^{-1} : \lambda \in \Sigma_\theta\}) \leq K_p$ and $\varphi_{\mathcal{R}}(\mathbb{A}) \geq \theta$. \square

5.2.4 Main result on parabolic systems of differential equations

First we make a preliminary remark on the approach we use in this section: our first aim is to prove \mathcal{R} -sectoriality of the $W^{k,p}$ -realization of a parabolic system. To this end, we essentially follow the approach given in [14, Chapter 6]. However, since we deal with Sobolev spaces of arbitrary order we need differentiability of the localized coefficients, which is not required for the localization in L^p as performed in [14, Chapter 6]. Therefore, we have to slightly modify the method used in [14, Chapter 6] by introducing a smoother localization that assures the well-definedness of the localized operators on the Sobolev space $W^{k,p}$.

Our second aim is the transference of \mathcal{R} -sectoriality to the realization of a parabolic system on certain interpolation spaces. This, in turn, is then obtained as an easy consequence of Theorem 3.18. As a well-known fact we first have

Lemma 5.25 (Partition of unity). *For all $r > 0$ there exists a $\varphi \in C_0^\infty(\mathbb{R}^n)$ with $0 \leq \varphi \leq 1$, $\text{supp}\varphi \subset (-r, r)^n$ and*

$$\sum_{l \in \Gamma} \varphi_l^2(x) = 1 \text{ for all } x \in \mathbb{R}^n.$$

Here we set $\Gamma := r\mathbb{Z}^n$ and $\varphi_l := \varphi(\cdot - l)$ for all $l \in \Gamma$. Additionally, we have $\text{supp}\varphi_l \subset l + [-\frac{3}{4}r, \frac{3}{4}r]^n$.

Definition and Lemma 5.26 (Localization- and patching-together-operator). *Let*

$$\mathbb{X}_k := \ell^p(\Gamma, W^{k,p}(\mathbb{R}^n, \mathbb{C}^N))$$

for $1 < p < \infty$, $N \in \mathbb{N}$, and $k \in \mathbb{N}_0$ with $\Gamma = r\mathbb{Z}$, $r > 0$. Let the sequence $(\varphi_l)_{l \in \Gamma}$ be as given in Lemma 5.25 and define the 'localization-operator'

$$J : L^p(\mathbb{R}^n, \mathbb{C}^N) \rightarrow \mathbb{X}_0, \quad f \mapsto (\varphi_l f)_{l \in \Gamma}$$

and the 'patching-together-operator'

$$P : \mathbb{X}_0 \rightarrow L^p(\mathbb{R}^n, \mathbb{C}^N), \quad (f_l)_{l \in \Gamma} \mapsto \left(x \mapsto \sum_{l \in \Gamma} \varphi_l(x) f_l(x) \right).$$

For $k \in \mathbb{N}_0$ we have

$$J|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} \in L(W^{k,p}(\mathbb{R}^n, \mathbb{C}^N), \mathbb{X}_k), \quad P|_{\mathbb{X}_k} \in L(\mathbb{X}_k, W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)), \quad (22)$$

and $PJ = \text{id}_{L^p(\mathbb{R}^n, \mathbb{C}^N)}$.

Proof. Follows easily by the smoothness of φ . □

Next we recall an interpolation inequality for \mathbb{C}^N -valued functions. The scalar case is proved in [1, 5.2 Theorem], for instance. The \mathbb{C}^N -valued case follows directly from the scalar case. We will apply this inequality later to verify the conditions of the perturbation result.

Lemma 5.27 (Interpolation inequality). *For every $m \in \mathbb{N}_0$ and each $\varepsilon_0 > 0$ there exists a constant $\mathcal{K}(p, n, k, N, \varepsilon_0) > 0$ such that for all $\varepsilon \in (0, \varepsilon_0]$ and $j \in \mathbb{N}_0$ with $0 \leq j < k$ and $u \in W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)$ we have*

$$\|u\|_{W^{j,p}(\mathbb{R}^n, \mathbb{C}^N)} \leq \mathcal{K}(\varepsilon \|u\|_{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} + \varepsilon^{-j/(k-j)} \|u\|_{L^p(\mathbb{R}^n, \mathbb{C}^N)}).$$

Theorem 5.28 ($W^{k,p}$ -realizations of parabolic systems). *Let $1 < p < \infty$, $k \in \mathbb{N}_0$, and the differential operator $A(x, D) := \sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha$ be given. Furthermore, assume that the coefficients satisfy the following regularities:*

$$a_\alpha \in BUC(\mathbb{R}^n, \mathbb{C}^{N \times N}) \cap C_b^k(\mathbb{R}^n, \mathbb{C}^{N \times N}) \text{ for } |\alpha| = m, \quad a_\alpha \in W^{k,p_\infty}(\mathbb{R}^n, \mathbb{C}^{N \times N}) \text{ for } |\alpha| < m$$

(see (8) for the definition of p_∞). If $A(x, D)$ is parabolic with constant $C_P > 0$, then there exists a $\nu > 0$ such that the $W^{k,p}$ -realization $A_{k,p} - \nu$ is \mathcal{R} -sectorial with $\varphi_{\mathcal{R}}(A_{k,p} - \nu) > \frac{\pi}{2}$.

Proof. 1. Localization (“Freezing the coefficients”):

First Remark 5.9 yields a $\theta \in (\frac{\pi}{2}, \pi)$ such that we even have the parameter-ellipticity of $A(x, D)$ in $\bar{\Sigma}_\theta$. For $M := \sum_{|\alpha|=m} \|a_\alpha\|_\infty$ let $\varepsilon = \varepsilon(M, C_P, \theta)$, $K = K(M, C_P, \theta) > 0$ be the constants as given in the statement of Proposition 5.22. By the uniform continuity of all a_α with $|\alpha| = m$ there exists a $\delta > 0$ such that

$$\sum_{|\alpha|=m} \|a_\alpha(x) - a_\alpha(y)\|_{\mathbb{C}^{N \times N}} < \varepsilon \quad (x, y \in \mathbb{R}^n, |x - y| < \delta).$$

Next, we choose $r > 0$ such that $\text{diam}(-r, r)^n < \delta$. For this r we choose a $\varphi \in C_0^\infty(\mathbb{R}^n)$ as in Lemma 5.25. Furthermore, we choose $\chi \in C_0^\infty(\mathbb{R}^n)$ satisfying $\chi(x) = 1$ for $\|x\|_1 := \sum_{k=1}^n |x_k| \leq \frac{7}{8}r$, $\chi(x) = 0$ for $\|x\|_1 \geq 1$, and $0 \leq \chi(x) \leq 1$ for all $x \in \mathbb{R}^n$.

Then we define for $l \in \Gamma := r\mathbb{Z}$ the localized differential operator

$$A^l(x, D) := \sum_{|\alpha|=m} a_\alpha^l(x) D^\alpha$$

with coefficients

$$a_\alpha^l(x) := a_\alpha \left(l + \chi \left(\frac{x-l}{r} \right) (x-l) \right) = \begin{cases} a_\alpha(x) & , \|x-l\|_1 \leq \frac{7}{8}r \\ a_\alpha(l) & , \|x-l\|_1 \geq r \\ a_\alpha(\tilde{x}) \text{ for some } \tilde{x} \in Q_l & , \text{ otherwise} \end{cases}$$

for $|\alpha| = m, l \in \Gamma$, and $Q_l := l + (-r, r)^n$. Next, we analyze the structure of these coefficients. For $\Phi_l : x \mapsto l + \chi \left(\frac{x-l}{r} \right) (x-l) \in Q_l$ we have $\Phi_l \in C_b^\infty(\mathbb{R}^n, \mathbb{R})$ and there exists a constant $C_\chi(r) > 0$ such that for all $l \in \Gamma$, $\gamma \in \mathbb{N}_0^n$ with $1 \leq |\gamma| \leq k$ we have $\|D^\gamma \Phi_l\|_\infty < C_\chi$. Thanks to $a_\alpha \in C_b^k(\mathbb{R}^n, \mathbb{C}^{n \times n})$ ($|\alpha| = m$) the classical chain rule applies and we obtain that $a_\alpha^l \in C_b^k(\mathbb{R}^n, \mathbb{C}^{N \times N})$. Moreover, there exists a constant $\tau > 0$ only depending on χ, r , and $\max_{\substack{0 \leq |\gamma| \leq k \\ |\alpha|=m}} \|D^\gamma a_\alpha\|_\infty$ such that

$$\max_{0 \leq |\gamma| \leq k} \|D^\gamma a_\alpha^l\|_\infty \leq \tau \text{ for all } l \in \Gamma, |\alpha| = m.$$

As before, let $A_{k,p}^l$ be the $W^{k,p}$ -realization of A^l . Due to Lemma 5.23 we can form the diagonal-operator

$$\mathbb{A}_{k,p} : D(\mathbb{A}_{k,p}) \subset \mathbb{X}_k \rightarrow \mathbb{X}_k, \quad (u_l)_{l \in \Gamma} \mapsto (A_{k,p}^l u_l)_{l \in \Gamma}$$

with $D(\mathbb{A}_{k,p}) := \mathbb{X}_{k+m}$.

2. \mathcal{R} -sectoriality of $\mathbb{A}_{k,p} - \mu$:

For the formal differential operator $A^l(x, D)$ we have the decomposition

$$A^l(x, D) = \underbrace{\sum_{|\alpha|=m} [a_\alpha^l(x) - a_\alpha(l)] D^\alpha}_{=: A'(x, D)} + \underbrace{\sum_{|\alpha|=m} a_\alpha(l) D^\alpha}_{=: A''(D)}.$$

Note that the second part $A''(D)$ is parameter-elliptic in $\bar{\Sigma}_\theta$ with the same constant C_P . For all $l \in \Gamma$, we have

$$\begin{aligned} \sum_{|\alpha|=m} \|a_\alpha(l)\|_{\mathbb{C}^{N \times N}} &\leq M & \sum_{|\alpha|=m} \|a_\alpha^l(\cdot) - a_\alpha(l)\|_\infty &< \varepsilon \\ \max_{\substack{0 < |\gamma| \leq k \\ |\alpha|=m}} \|D^\gamma (a_\alpha^l(\cdot) - a_\alpha(l))\|_\infty &\leq \tau & \text{for } k \neq 0. \end{aligned}$$

Applying Proposition 5.22 yields a constant $\mu(M, C_P, \tau, \theta) > 0$ independent of $l \in \Gamma$ such that $A_{k,p}'' + A_{k,p}^l - \mu = A_{k,p}^l - \mu$ is \mathcal{R} -sectorial with $\theta \in \Theta_{\mathcal{R}}(A_{k,p}^l - \mu)$ and $R_{\theta,p}(A_{k,p}^l - \mu) \leq K$. Since K according to Proposition 5.22 only depends on C_P, M , and θ and since these constants are uniformly in l , we obtain

$$R_{\theta,p}(A_{k,p}^l - \mu) \leq K \quad (l \in \Gamma).$$

Additionally, Proposition 5.22 yields

$$\sup_{l \in \Gamma} \|(\lambda + \mu - A_{k,p}^l)^{-1}\|_{L(W^{k,p}(\mathbb{R}^n, \mathbb{C}^N), W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N))} \leq C_\lambda,$$

where the constant C_λ only depends on λ, M, C_P und μ . Thus, Lemma 5.24 implies the \mathcal{R} -sectoriality of $\mathbb{A}_{k,p} - \mu$ with

$$\theta \in \Theta_{\mathcal{R}}(\mathbb{A}_{k,p} - \mu) \quad \text{and} \quad R_{\theta,p}(\mathbb{A}_{k,p} - \mu) \leq K.$$

3. Determination of $JA_{k,p} - \mathbb{A}_{k,p}J$ and $A_{k,p}P - P\mathbb{A}_{k,p}$:

In essentially the same way as in [14], we obtain

$$(JA_{k,p} - \mathbb{A}_{k,p}J)u = \mathbb{B}_{k,p}Ju, \quad (23)$$

$$(A_{k,p}P - P\mathbb{A}_{k,p})(u_l)_{l \in \Gamma} = P\mathbb{D}_{k,p}(u_l)_{l \in \Gamma}, \quad (24)$$

where we set $D(\mathbb{B}_{k,p}) := \mathbb{X}_{k+m-1}$, $D(\mathbb{D}_{k,p}) := \mathbb{X}_{k+m-1}$, $A_{\text{low}}(x, D) := A(x, D) - A_0(x, D)$, and where the operators $\mathbb{B}_{k,p}$ and $\mathbb{D}_{k,p}$ are defined as

$$\begin{aligned} \mathbb{B}_{k,p} : D(\mathbb{B}_{k,p}) \subset \mathbb{X}_k &\rightarrow \mathbb{X}_k \\ (u_l)_{l \in \Gamma} &\mapsto \left(A_{\text{low}}(x, D)u_l + \sum_{\substack{j \in \Gamma: \\ Q_j \cap Q_l \neq \emptyset}} (\varphi_l A(x, D) - A(x, D)\varphi_l)(\varphi_j u_j) \right)_{l \in \Gamma}, \end{aligned}$$

$$\begin{aligned} \mathbb{D}_{k,p} : D(\mathbb{D}_{k,p}) \subset \mathbb{X}_k &\rightarrow \mathbb{X}_k \\ (u_l)_{l \in \Gamma} &\mapsto \left(A_{\text{low}}(x, D)u_l + \sum_{\substack{j \in \Gamma: \\ Q_j \cap Q_l \neq \emptyset}} \varphi_l (A(x, D)\varphi_j - \varphi_j A(x, D))u_j \right)_{l \in \Gamma}. \end{aligned}$$

Observe that in $\mathbb{B}_{k,p}$ and $\mathbb{D}_{k,p}$ there appear only derivatives of order less or equal to $m - 1$.

5. Perturbation of $\mathbb{A}_{k,p} - \mu$:

Our aim is to apply Theorem 5.20 to the operator $\mathbb{A}_{k,p}^{(\mu)} := \mathbb{A}_{k,p} - \mu$ combined with the perturbation $\mathbb{B}_{k,p}$ and to $\mathbb{A}_{k,p}^{(\mu)}$ combined with $\mathbb{D}_{k,p}$. At first we have $1 \in \rho(\mathbb{A}_{k,p}^{(\mu)})$ and $(1 - \mathbb{A}_{k,p}^{(\mu)})^{-1} \in L(\mathbb{X}_k, \mathbb{X}_{k+m})$ due to Lemma 5.23. Next we set $\varepsilon_0 := 1$ and choose $0 < \varepsilon' < 1$ such that

$$a := \varepsilon' \mathcal{K}M_{\mathbb{B}, \mathbb{D}} \cdot \left\| (1 - \mathbb{A}_{k,p}^{(\mu)})^{-1} \right\|_{L(\mathbb{X}_k, \mathbb{X}_{k+m})} < \left(\tilde{N}_\theta \left(\mathbb{A}_{k,p}^{(\mu)} \right) \left(C'_\theta + C_\theta \tilde{R}_{\theta,p} \left(\mathbb{A}_{k,p}^{(\mu)} \right) \right) \right)^{-1},$$

with $M_{\mathbb{B}, \mathbb{D}} := \max\{\|\mathbb{B}_{k,p}\|_{L(\mathbb{X}_{k+m-1}, \mathbb{X}_k)}, \|\mathbb{D}_{k,p}\|_{L(\mathbb{X}_{k+m-1}, \mathbb{X}_k)}\}$ and with $\mathcal{K} > 0$ from Lemma 5.27. For $(u_l)_{l \in \Gamma} \in \mathbb{X}_{k+m-1}$, Lemma 5.27 and the boundedness of $\mathbb{B}_{k,p}$ and $\mathbb{D}_{k,p}$ yield

$$\left. \begin{aligned} \|\mathbb{B}_{k,p}(u_l)_{l \in \Gamma}\|_{\mathbb{X}_k} \\ \|\mathbb{D}_{k,p}(u_l)_{l \in \Gamma}\|_{\mathbb{X}_k} \end{aligned} \right\} \leq a' \|(u_l)_{l \in \Gamma}\|_{\mathbb{X}_{k+m}} + b' \|(u_l)_{l \in \Gamma}\|_{\mathbb{X}_0} \leq a' \|(u_l)_{l \in \Gamma}\|_{\mathbb{X}_{k+m}} + b' \|(u_l)_{l \in \Gamma}\|_{\mathbb{X}_k}$$

for $a' := \varepsilon' \mathcal{K}M_{\mathbb{B}, \mathbb{D}}$ and $b' := \mathcal{K}M_{\mathbb{B}, \mathbb{D}} \cdot (\varepsilon')^{-(k+m-1)}$. By virtue of

$$\|(u_l)_{l \in \Gamma}\|_{\mathbb{X}_{k+m}} \leq \left\| (1 - \mathbb{A}_{k,p}^{(\mu)})^{-1} \right\|_{L(\mathbb{X}_k, \mathbb{X}_{k+m})} \cdot \left(\left\| \mathbb{A}_{k,p}^{(\mu)}(u_l)_{l \in \Gamma} \right\|_{\mathbb{X}_k} + \|(u_l)_{l \in \Gamma}\|_{\mathbb{X}_k} \right)$$

we conclude

$$\left. \begin{aligned} \|\mathbb{B}_{k,p}(u_l)_{l \in \Gamma}\|_{\mathbb{X}_k} \\ \|\mathbb{D}_{k,p}(u_l)_{l \in \Gamma}\|_{\mathbb{X}_k} \end{aligned} \right\} \leq a \left\| \left(\mathbb{A}_{k,p}^{(\mu)} u_l \right)_{l \in \Gamma} \right\|_{\mathbb{X}_k} + \left(b' + a' \left\| (1 - \mathbb{A}_{k,p}^{(\mu)})^{-1} \right\|_{L(\mathbb{X}_k, \mathbb{X}_{k+m})} \right) \cdot \|(u_l)_{l \in \Gamma}\|_{\mathbb{X}_k}.$$

Hence, from Theorem 5.20 we infer that there exists an $\eta > 0$ such that $\mathbb{A}_{k,p}^{(\mu)} + \mathbb{B}_{k,p} - \eta$ and $\mathbb{A}_{k,p}^{(\mu)} + \mathbb{D}_{k,p} - \eta$ are \mathcal{R} -sectorial with $\theta \in \Theta_{\mathcal{R}} \left(\mathbb{A}_{k,p}^{(\mu)} + \mathbb{B}_{k,p} - \eta \right) \cap \Theta_{\mathcal{R}} \left(\mathbb{A}_{k,p}^{(\mu)} + \mathbb{D}_{k,p} - \eta \right)$.

6. Determination of the resolvent $(A_{k,p} - (\mu + \eta))^{-1}$:

Let $\lambda \in \Sigma_{\theta}$ and $\nu := \mu + \eta > 0$; Due to (23) and (24) we obtain a left inverse of $\lambda - (A_{k,p} - \nu)$ in form of

$$P \left(\lambda + \eta - \left(\mathbb{A}_{k,p}^{(\mu)} + \mathbb{B}_{k,p} \right) \right)^{-1} J_{|W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} \quad (25)$$

and a right inverse given as

$$P \left(\lambda + \eta - \left(\mathbb{A}_{k,p}^{(\mu)} + \mathbb{D}_{k,p} \right) \right)^{-1} J_{|W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)}. \quad (26)$$

Hence (25) and (26) coincide and we have $\lambda \in \rho(A_{k,p} - \nu)$. The \mathcal{R} -boundedness follows directly from

$$\lambda (\lambda - (A_{k,p} - \nu))^{-1} = P \left[\lambda \left(\lambda - \left(\mathbb{A}_{k,p}^{(\mu)} + \mathbb{D}_{k,p} - \eta \right) \right)^{-1} \right] J \quad (\lambda \in \Sigma_{\theta})$$

and the fact that $P|_{\mathbb{X}_k} \in L(\mathbb{X}_k, W^{k,p}(\mathbb{R}^n, \mathbb{C}^N))$ and $J_{|W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)} \in L(W^{k,p}(\mathbb{R}^n, \mathbb{C}^N), \mathbb{X}_k)$. Consequently $R_{\theta,p}(A_{k,p} - \nu) \leq C(P, J)R_{\theta,p} \left(\mathbb{A}_{k,p}^{(\mu)} + \mathbb{D}_{k,p} - \eta \right)$, which implies $A_{k,p} - \nu$ to be \mathcal{R} -sectorial with \mathcal{R} -angle $\varphi_{\mathcal{R}}(A_{k,p} - \nu) \geq \theta > \frac{\pi}{2}$. \square

With the help of Theorem 3.22 the above result generalizes to parabolic systems realized on interpolation spaces.

Theorem 5.29 (\mathcal{R} -sectoriality of (\mathcal{F}, k, p) -realizations of parabolic systems). *Let $k \in \mathbb{N}_0$ and $A(x, D) := \sum_{|\alpha| \leq m} a_{\alpha}(x)D^{\alpha}$ be a formal differential operator, where we assume the coefficients to have the following regularities:*

$$a_{\alpha} \in BUC(\mathbb{R}^n, \mathbb{C}^{N \times N}) \cap C_b^k(\mathbb{R}^n, \mathbb{C}^{N \times N}) \text{ for } |\alpha| = m, \quad a_{\alpha} \in W^{k,p\infty}(\mathbb{R}^n, \mathbb{C}^{N \times N}) \text{ for } |\alpha| < m.$$

Let $1 < p, q < \infty$ and \mathcal{F} be an L^q -compatible interpolation functor of type h such that

$$W^{k,p}(\mathbb{R}^n, \mathbb{C}^N) \hookrightarrow_d \mathcal{F}(\{L^p(\mathbb{R}^n, \mathbb{C}^N), W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)\}),$$

and let $A_{\mathcal{F},k,p}$ be the (\mathcal{F}, k, p) -realization of $A(x, D)$ as defined in (7). If $A(x, D)$ is parabolic, then there exists a $\nu > 0$ such that $A_{\mathcal{F},k,p} - \nu$ is \mathcal{R} -sectorial with $\varphi_{\mathcal{R}}(A_{\mathcal{F},k,p} - \nu) > \frac{\pi}{2}$.

Proof. By the assumed regularity of the coefficients the L^p - and $W^{k,p}$ -realizations A_p and $A_{k,p}$ are well-defined. Theorem 5.28 yields the \mathcal{R} -sectoriality of $A_0 := A_p - \nu$ and $A_1 := A_{k,p} - \nu$ for a $\nu > 0$ with $\varphi_{\mathcal{R}}(A_p - \nu) > \frac{\pi}{2}$ and $\varphi_{\mathcal{R}}(A_{k,p} - \nu) > \frac{\pi}{2}$. We set

$$\begin{aligned} X_0 &:= L^p(\mathbb{R}^n, \mathbb{C}^N), & D(A_0) &:= W^{m,p}(\mathbb{R}^n, \mathbb{C}^N), \\ X_1 &:= W^{k,p}(\mathbb{R}^n, \mathbb{C}^N), & D(A_1) &:= W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N). \end{aligned}$$

By the same arguments as in Corollary 5.13 we obtain in combination with Theorem 3.22 the \mathcal{R} -sectoriality of the operator

$$A_{\mathcal{F},k,p} - \nu : D(A_{\mathcal{F},k,p}) \subset \mathcal{F}(\{L^p(\mathbb{R}^n, \mathbb{C}^N), W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)\}) \rightarrow \mathcal{F}(\{L^p(\mathbb{R}^n, \mathbb{C}^N), W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)\})$$

with $D(A_{\mathcal{F},k,p}) := \mathcal{F}(\{W^{k,p}(\mathbb{R}^n, \mathbb{C}^N), W^{k+m,p}(\mathbb{R}^n, \mathbb{C}^N)\})$ and $(A_{\mathcal{F},k,p} - \nu)f := (A(x, D) - \nu)f$ for $f \in D(A_{\mathcal{F},k,p})$. Moreover, Theorem 3.22 yields $\varphi_{\mathcal{R}}(A_{\mathcal{F},k,p} - \nu) > \frac{\pi}{2}$, hence the assertion is proved. \square

Corollary 5.30. *The parabolic system described in Theorem 5.29 has maximal L^p -regularity on the space $\mathcal{F}(\{L^p(\mathbb{R}^n, \mathbb{C}^N), W^{k,p}(\mathbb{R}^n, \mathbb{C}^N)\})$.*

Proof. The characterization of maximal L^p -regularity by \mathcal{R} -sectoriality with \mathcal{R} -angle bigger than $\pi/2$ yields the assertion. This characterization can be found in [19], [14], or [5]. \square

Corollary 5.31 (\mathcal{R} -sectoriality of parabolic systems on Besov- and Bessel-potential-spaces).

Assume the situation of Theorem 5.29 to be given. Then the Besov- and Bessel-potential-space realizations $A_{s,p,q}^{\mathcal{B}}$ and $A_{s,p}^{\mathcal{H}}$ as defined in Remark 5.5 are \mathcal{R} -sectorial on $B_{p,q}^s(\mathbb{R}^n, \mathbb{C}^N)$ and $H^{s,p}(\mathbb{R}^n, \mathbb{C}^N)$ with $\varphi_{\mathcal{R}}(A_{s,p,q}^{\mathcal{B}} - \nu) > \frac{\pi}{2}$ and $\varphi_{\mathcal{R}}(A_{s,p}^{\mathcal{H}} - \nu) > \frac{\pi}{2}$, respectively.

Proof. This follows directly from Theorem 5.29 and Remark 3.24. \square

Acknowledgement: We want to thank Lutz Weis for helpful hints concerning the first part of this paper. Furthermore, we also want to express our gratitude to Robert Denk for his cooperativeness and a plenty of fruitful discussions.

References

- [1] R.A. Adams and J.J.F. Fournier. *Sobolev Spaces*. Pure and Applied Mathematics. Academic Press, 2003. second edition.
- [2] M.S. Agranovich and M.I. Vishik. Elliptic problems with a parameter and parabolic problems of general form. *Russ. Math. Surv.*, 19(3):53–157, 1964. Engl. transl. of *Usp. Mat. Nauk.* 19(3):53-161, 1964.
- [3] H. Amann. *Linear and Quasilinear Parabolic Problems. Vol. I*. Birkhäuser, 1995.
- [4] H. Amann. Anisotropic function spaces and maximal regularity for parabolic problems. part 1: Function spaces. *Jindrich Necas Center for Mathematical Modeling Lecture Notes*, 6, 2009.
- [5] R. Denk, M. Hieber, and J. Prüß. \mathcal{R} -boundedness, Fourier multipliers and problems of elliptic and parabolic type. *Mem. Amer. Math. Soc.*, 166:viii+114, 2003.
- [6] R. Denk, J. Saal, and J. Seiler. Inhomogeneous symbols, the Newton polygon, and maximal L^p -regularity. *Russian J. Math. Phys.*, 15(2):171–192, 2008.
- [7] R. Denk, J. Saal, and J. Seiler. Bounded H_{∞} -calculus for pseudo-differential Douglis-Nirenberg systems of mild regularity. *Math. Nachr.*, 282(3):386–407, 2009.
- [8] J. Diestel, H. Jarchow, and A. Tonge. *Absolutely Summing Operators*, volume 43 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 1995.
- [9] G. Dore and A. Venni. H^{∞} -calculus for an elliptic operator on a halfspace with general boundary conditions. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (5)*, 1(3):487–543, 2002.
- [10] G. Dore and A. Venni. H^{∞} functional calculus for sectorial and bisectorial operators. *Studia Math.*, 166(3):221–241, 2005.
- [11] Nigel Kalton and Stephen Montgomery-Smith. Interpolation of Banach spaces. In *Handbook of the geometry of Banach spaces, Vol. 2*, pages 1131–1175. North-Holland, Amsterdam, 2003.
- [12] N.J. Kalton, P.C. Kunstmann, and L. Weis. Perturbation and interpolation theorems for the H^{∞} -calculus with applications to differential operators. *Math. Ann.*, 336(4):747–801, 2006.
- [13] N.J. Kalton and L. Weis. The H^{∞} -calculus and sums of closed operators. *Math. Ann.*, 321:319–345, 2001.
- [14] P.C. Kunstmann and L. Weis. *Maximal L_p -regularity for Parabolic Equations, Fourier Multiplier Theorems, and H^{∞} -functional Calculus*, volume 1855 of *Functional Analytic Methods for Evolution Equations, Lecture notes in Math*. Springer, 2004.

-
- [15] J. Prüß, J. Saal, and G. Simonett. Existence of analytic solutions for the classical Stefan problem. *Math. Ann.*, 338:703–755, 2007.
- [16] P. Rabier. An isomorphism theorem for linear evolution problems on the line. *Journal of Dynamics and Differential Equations*, 15(4):779–806, 2003.
- [17] V.A. Solonnikov. Estimates in L_p of solutions of elliptic and parabolic systems. *Proc. Steklov Inst. Math.*, 102:157–185, 1970. Engl. transl. of Tr. Mat. Inst. Steklova 102:137-160, 1967.
- [18] H. Triebel. *Interpolation Theory, Function Spaces, Differential Operators*. North Holland, 1978.
- [19] L. Weis. Operator-valued multiplier theorems and maximal L^p -regularity. *Math. Ann.*, 319(4):735–758, 2001.
-