

Parasite identification, succession and infection pathways in perch fry (*Perca fluviatilis*): new insights through a combined morphological and genetic approach

JASMINCA BEHRMANN-GODEL*

Limnological Institute, University of Konstanz, 78464 Konstanz, Germany

(Received 5 March 2012; revised 20 September and 18 October 2012; accepted 18 October 2012; first published online 2 January 2013)

SUMMARY

Identification of parasite species is particularly challenging in larval and juvenile hosts, and this hampers the understanding of parasite acquisition in early life. The work described here employs a new combination of methods to identify parasite species and study parasite succession in fry of perch (*Perca fluviatilis*) from Lake Constance, Germany. Classical morphological diagnostics are combined with sequence comparisons between parasite life-stages collected from various hosts within the same ecosystem. In perch fry at different stages of development, 13 different parasite species were found. Incomplete morphological identifications of cestodes of the order Proteocephalidea, and trematodes of the family Diplostomatidae were complemented with sequences of mitochondrial DNA (cytochrome oxidase 1) and/or nuclear (28S rDNA) genes. Sequences were compared to published data and used to link the parasites in perch to stages from molluscs, arthropods and more easily identifiable developmental stages from other fishes collected in Lake Constance, which both aided parasite identification and clarified transmission pathways. There were distinct changes in parasite community composition and abundance associated with perch fry age and habitat shifts. Some parasites became more abundant in older fish, whereas the composition of parasite communities was more strongly affected by the ontogenetic shifts from the pelagic to the littoral zone.

Key words: parasite community, transmission, Diplostomatidae, Proteocephalidea, barcoding genes, 28S rDNA, CO1, ITS1-5.8S-ITS2 rDNA.

INTRODUCTION

Many marine and freshwater fish species produce vast numbers of eggs. Only a minority will be fertilized, hatch, develop into juveniles and reach maturity. High levels of larval and juvenile mortality are due mainly to predation but starvation, disease and parasitism also take a significant toll (Wootton, 1998). An increasing number of studies show that the early developmental stages of fish are targeted by several species of endo- and ectoparasites (Balbuena *et al.* 2000; King and Cone, 2009; Pracheil and Muzzall, 2009; Skovgaard *et al.* 2009a). Parasitic infections may be detrimental to fish fry (Grutter *et al.* 2009; Skovgaard *et al.* 2009b; Kelly *et al.* 2010; Nendick *et al.* 2011), but there still is a tremendous lack of knowledge about the relationship, especially in freshwater systems. The timing of first parasite encounter and subsequent parasite succession are almost unstudied. The study of parasite infections in fish fry presents challenges because the young age of the hosts means that any infections observed must be very recent. Thus the parasites are themselves in early stages of development (Kuchta *et al.* 2009), and may lack important morphological characteristics

required for precise species determination. Classically, this problem has been addressed with life-cycle studies including also experimental infections. However, such studies are laborious and usually limited to small numbers of congeneric parasites. In contrast, difficulty in the identification of parasite species often scales up to higher levels, such as that among cestodes of the order Proteocephalidea, and trematodes of the family Diplostomatidae (Wootton 1974; Kennedy, 1978).

Molecular identification techniques based on sequences of nuclear genes such as ribosomal rRNA and the mitochondrial cytochrome oxidase gene can be helpful in otherwise tricky species diagnoses (Zehnder and Mariaux, 1999; Scholz *et al.* 2007; Sonnenberg *et al.* 2007; Locke *et al.* 2010a,b). An increasing number of sequences are becoming available for comparison in various databases. Unfortunately, not all are backed up by solid taxonomic work, without which reference sequences cannot be fully relied on for species identification purposes. Furthermore, accordance between a reliable reference and the investigated sequence must be very good to allow confident species determination. This study develops a method that combines classical morphological analysis with a genetic survey of various parasite life-cycle stages to solve this problem. Preliminary identification of

* Corresponding author: Tel: ++49 7531 884536. Fax: ++49 7531 883533. E-mail: Jasminca.Behrmann@uni-konstanz.de

species of parasites infecting perch fry was carried out using morphological characteristics, and was supported by sequencing of DNA of other, more easily identifiable developmental stages from the same ecosystem. This method was also applied to investigate transmission between hosts. The data are used in an ecological investigation of macroparasite infection of perch fry in Lake Constance.

The Eurasian perch (*Perca fluviatilis* L.) is a widely distributed fish species of the Palaearctic region. Perch harbour a variety of ecto- and endoparasites belonging to various orders (Balling and Pfeiffer, 1997a–d; Carney and Dick, 1999; Morozinska-Gogol, 2008), but very little is known about parasites of the larval and early juvenile life stages. The author knows of only 1 study, by Kuchta *et al.* (2009), which compared infection rates of perch fry from 3 spatially distinct subpopulations, epipelagic, bathypelagic and littoral, in a deep, canyon-shaped reservoir in the Czech Republic. The authors found 6 species of endoparasites. Infection rates increased with fish age and the highest infection rates were found in the littoral population. In deep, natural lakes in Europe and North America, the life histories of perch and its sister species yellow perch (*Perca flavescens*) typically include early habitat shifts. Perch spawn in May/June in the littoral zones. Soon after hatching, larvae are found in the pelagic zone where they remain up until metamorphosis. Soon afterwards they return to the littoral and inhabit shallow water until autumn when they move into deeper water to overwinter (Coles, 1981; Wang and Eckmann, 1994; Imbrock *et al.* 1996; Miehl and Dettmers, 2011). Thus perch occupy different habitats and exploit different food sources early in their ontogeny, switching from planktonic to benthic prey (Wang and Appenzeller, 1998; Weber *et al.* 2011). These niche shifts alter the exposure and vulnerability of young perch to a number of parasite species (Johnson *et al.* 2004; McCairns and Fox, 2004; Zelmer and Arai, 2004; Bertrand *et al.* 2008; Pracheil and Muzzall, 2009). For fish larvae and early juveniles, the consumption of zooplankton is a major route of infection with helminths. Their occurrence in the zooplankton is often spatially and temporally synchronized with the presence of intermediate and definitive hosts in spring (Rosenthal, 1967; Marcogliese, 1995; Johnson *et al.* 2006; Lahnsteiner *et al.* 2009). Young perch respond to seasonal changes in abiotic parameters such as water temperature and day length, which can also strongly influence the occurrence and density of infective parasite stages, including those whose complex life cycles involve many intermediate hosts, for example most cestodes and the digenean trematodes (Marcogliese, 1995; Faltynkova *et al.* 2009). Another challenge for larval and early juvenile fish is the slow maturation of their immune system during ontogeny. However, both the innate and the adaptive immune functions of perch can play a major

role in the defence against helminth infections (Secombes and Chappell, 1996).

In this contribution, macroparasites were surveyed in order to characterize the chronology of infection in perch during the first 3 months of life in Lake Constance. Specifically, the aims were to (1) identify parasites of perch fry, (2) study parasite succession in perch fry, (3) investigate the effects of increasing age and the ontogenetic habitat shift of host fish on the development of the parasite community and (4) clarify transmission pathways between hosts.

MATERIALS AND METHODS

Sampling and age determination of perch fry

Perch were sampled from the large pre-alpine Upper Lake Constance in south Germany. Many previous studies on the perch life cycle have been conducted in this system and have shown that perch here undergo typical ontogenetic habitat shifts (Wang and Eckmann, 1994; Eckmann and Imbrock, 1996; Imbrock *et al.* 1996; Wang and Appenzeller, 1998). Larval perch were sampled from the pelagic zone of the lake at night, using an 800 μm mesh ichthyoplankton net towed close to the surface by boat approximately 500 m offshore. These pelagic perch samples were from 2 different years. The 2 youngest samples were from 8 and 24 June 2010. Routine weekly samplings taken during and after the spawning season of this year, recorded the first yolk sac larvae in the pelagic zone on 1 June. Since perch use up their yolk sac within 1 week, the hatching date of the 2010 cohort was estimated to be 23 May. Hence, the ages of the first two pelagic samples were estimated at 2 and 4 weeks post-hatch (p.h.). The third pelagic and all littoral perch samples were from 2009 (pelagic sample from 1 and 7 July, littoral samples from 8 and 16 July and 14 and 16 August). Using 3 fish from each 2009 sample ($n=18$), age was determined by counting daily increments on the lapilli (Campana, 1992). From these counts, a hatching date of 20 May was estimated for the 2009 cohort. Thus, the ages of the third pelagic and both littoral samples were estimated to be 7, 8 and 12 weeks p.h. respectively. The two littoral perch samples were netted with beach seines of 2 and 4 mm mesh sizes hauled from 1 m depth to the shore at night. Live perch were transported in bottles of aerated lake water and killed with an overdose of MS 222. Fish that died during sampling were stored in 0.64% NaCl solution in small bags kept on ice, and examined within 24 h.

Parasite assessment and morphological identification

Perch fry were squeezed between glass plates for microscopical analysis and every organ and tissue was examined for the presence of macroparasites. Additionally, the NaCl solutions in which dead fish had been stored were screened for escaped parasites.

Parasites were identified based on typical morphological features identified in previous studies (Bykhovskaya-Pavlovskaya *et al.* 1964; Wootten, 1974; Schäperclaus, 1979; Scholz *et al.* 1998). All cestodes from the suborder Proteocephalidae and some of the Triaenophoridae were observed to be in very early developmental stages, and could thus be identified only to family level. For species determination, cestode subsets were removed from perch (2–4 weeks p.h.) of the 2010 cohort and from additional perch fry of 1 sampling event in 2011 and stored in 95% ethanol for genetic analysis. To obtain comparative sequences, cestodes from both suborders were also sampled from various adult fish hosts in Lake Constance. Fish species sampled were adult perch ($n=4$), whitefish *Coregonus lavaretus* ($n=2$), arctic char *Salvelinus alpinus* ($n=1$), northern pike *Esox lucius* ($n=1$) and burbot *Lota lota* ($n=1$). Fish were dissected and late developmental stage and adult cestodes were identified to the species level based on morphological characteristics as described above. Samples were stored in 95% ethanol for genetic analysis.

Pelagic copepods (*Cyclops* spp.) were collected in June 2011 and from April to September 2012. Sampling for copepods took place in approximately the same location as perch fry had been caught, using an ichthyoplankton net 1·10 m in diameter with 200 μm mesh. The net was towed approximately 0·5 m below the water surface behind a boat. Plankton samples were transported to the laboratory, sieved through 500 μm mesh, collected in beakers with filtered lake water and stored at 4 °C until analysis. A small portion of the collected sample was placed in a glass Petri dish, anaesthetized with carbonated water and examined under a dissection binocular microscope. For further investigation cyclopoid copepods (copepodids C5 and adults) were dissected using Dumont tweezers and preparation needles and analysed for proceroid infection. All proceroids found were stored in 95% ethanol. Then 19 proceroids (9 from 2011 and 10 from 2012) were selected at random for genetic analysis.

Eye flukes (metacercariae of diplostomatid trematodes) from perch fry were determined morphologically to family level only. For species determination via genetic comparison, metacercariae were sampled from the eye lenses and the vitreous body of several adult fish hosts in Lake Constance: perch ($n=9$), ruffe *Gymnocephalus cernuus* ($n=1$), whitefish ($n=2$) and roach *Rutilus rutilus* ($n=1$). To investigate the emergence of eye fluke cercariae in Lake Constance, lymnaeid snails *Radix auricularia* and *R. labiata* were sampled at 3 littoral locations between May and October 2010. Snails were taken to the laboratory and placed individually in 150 ml plastic bowls for a maximum of 7 weeks. The temperature was 20 °C and bowls were illuminated artificially with light: dark cycles corresponding to natural conditions. The

snails were fed with commercial fish food (Tetramin[®]) and frozen chironomids and the water was exchanged every 2–3 days. Before water exchange, bowls were screened for shed cercariae. Cercariae of infected snails were observed under a microscope after vital staining with Neutral Red. Diplostomatid cercariae were morphologically determined to the family and where possible to the species level with the help of available literature (Niewiadomska and Kieseliene, 1994; Niewiadomska and Laskowski, 2002; Faltynkova, 2005). Cercariae were collected and stored in 95% ethanol for later genetic analysis. From every infected snail, 2 cercariae were genetically analysed.

Direct sequencing

Parasite DNA was extracted from whole individuals, or from small pieces in the case of adult cestodes, using the Chelex[™] extraction technique (Criscione and Blouin, 2004). The primer combinations LSU D1-D2 fw1 and LSU D1-D2 rev2 were used to amplify the D1-D2 fragment of the LSU rDNA gene region according to the protocol of Sonnenberg *et al.* (2007). For trematodes, part of the mitochondrial cytochrome oxidase subunit 1 (CO1) and the complete internal transcribed spacer 1 and 2 (ITS1-5·8S-ITS2) of the rDNA gene were amplified with the use of established primers and protocols: Plat-diploCOX1dF/R (Moszczyńska *et al.* 2009) and D1/2 (Galazzo *et al.* 2002). PCR products were purified using Exo/FastAP (Fermentas), prepared for sequencing using the BigDye termination reaction chemistry (Applied Biosystems) and sequenced in both directions in an ABI 3130 capillary DNA sequencer (Applied Biosystems). All unique traces were compared to previously published sequences from GenBank with a BLAST search in order to find identical sequences. The LSU D1-D2, CO1 and ITS1-5·8S-ITS2 sequences from parasites found in this study were aligned with the help of BioEdit (Hall, 1999). Neighbour-joining (NJ) trees (Saitou and Nei, 1987) were constructed from a Kimura-2-parameter distance matrix (Kimura, 1980) using MEGA software (Tamura *et al.* 2011).

Representative unique sequences of genes newly sequenced here were submitted to the GenBank database. These include 5 LSU rDNA gene region sequences from cestodes (GenBank ID: JQ639165–JQ639169), and 35 CO1 sequences (GenBank ID: JQ639170–JQ639204) and 5 ITS1-5·8S-ITS2 sequences from trematodes (GenBank ID: JQ665456–JQ665460).

Statistical analysis of the parasite community development

To investigate the development of the parasite community of perch fry without interference or variability due to inter-annual variation, analyses

Table 1. Parasite species, prevalence and mean intensity of infection found in perch of Lake Constance sampled at different age

(First 3 columns (2, 4 and 7 weeks p.h. (wph), grey box) represent the pelagic, the other 2 (8 and 12 weeks p.h.) the littoral samples. *Proteocephalus* spp. includes *P. percae* and *P. longicollis*). Added is information on the invertebrate hosts (1.IH = first, 2.IH = second intermediate host), the species names are given in parentheses if known for Lake Constance from own observations.)

Age wph (n fish)	2 (51)	4 (85)	7 (52)	8 (60)	12 (60)	
Year of sampling	2010	2010	2009	2009	2009	
Parasite species	Prevalence (mean intensities)					Invertebrate host
Cestoda						
<i>Triaenophorus nodulosus</i> *	0(0)	0(0)	52(1.5)	40(1.2)	50(1.5)	
<i>T. nodulosus</i> **	0(0)	0(0)	19(1.4)	2(1.0)	3(1.0)	1.IH: Copepods (<i>Cyclops</i> spp.)
<i>Proteocephalus</i> spp.	0(0)	12(1.1)	71(4.8)	42(3.3)	13(1.3)	1.IH: Copepods (<i>Cyclops</i> spp.)
<i>Eubothrium crassum</i>	0(0)	70(3.0)	63(3.5)	88(6.8)	40(3.8)	1.IH: Copepods
Trematoda						
<i>Bunodera luciopercae</i>	0(0)	6(1.2)	63(2.7)	87(4.9)	93(13.2)	1.IH: Bivalves; 2.IH: Copepods, cladocerans, ostracods, amphipods, ephemeroptera
<i>Diplostomum baeri</i>	0(0)	0(0)	0(0)	13(1.3)	67(7.2)	1.IH: Snails
<i>Thyodelphys</i> sp.2	0(0)	0(0)	0(0)	52(2.1)	100(64)	1.IH: Snails (<i>Radix auricularia</i> , <i>R. labiata</i>)
<i>Ichthyocotylurus variegatus</i>	0(0)	0(0)	0(0)	0(0)	15(4.7)	1.IH: Snails (<i>Valvata piscinalis</i>)
<i>Cotylurus pileatus</i>	0(0)	0(0)	0(0)	8(1.4)	35(7.2)	1.IH: Snails
<i>Hysteromorpha triloba</i>	0(0)	0(0)	0(0)	25(4.0)	22(1.8)	1.IH: Snails
<i>Bucephalus polymorphus</i>	0(0)	0(0)	0(0)	5(1.6)	0(0)	1.IH: Bivalves
Nematoda						
<i>Raphidascaris acus</i>	0(0)	0(0)	0(0)	2(1.0)	0(0)	—
Maxillopoda						
<i>Ergasilus sieboldi</i>	0(0)	0(0)	0(0)	0(0)	1(1.0)	—
<i>Argulus foliaceus</i>	0(0)	0(0)	0(0)	5(2.0)	5(1.6)	—

* Plerocercoids; ** Proceroids.

were restricted to data for 1 year class, including 7, 8 and 12-week-old fish of the 2009 cohort. In this analysis, the abundance of each parasite species was determined for every fish. A Bray-Curtis dissimilarity matrix was then calculated and the relationship between fish age and parasite community structure was visualized with 2-dimensional nonmetric multi-dimensional scaling (nMDS; Kruskal, 1964) using the software PRIMER-E v6 (Clarke, 1993). The nMDS yields a plot in which the parasite community of each individual fish is represented as a dot. The closer together the dots appear on the plot, the more similar the parasite communities of individual fish are to one another. In labelling fish by age (weeks 7, 8 and 12), age-related clustering can be visualized. The difference in parasite communities between the 3 age groups was tested for significance with an analysis of similarity (ANOSIM) implemented in the software PRIMER-E.

RESULTS

Parasite species identification based on morphological characteristics

The perch fry sampled in this study were infected with 13 different parasite taxa (Table 1). Nine species

could be identified unambiguously to species level based on morphological characters: *Bunodera luciopercae*, *Triaenophorus nodulosus* (hook carrying plerocercoids in perch liver), *Cotylurus pileatus*, *Ichthyocotylurus variegatus*, *Hysteromorpha triloba*, *Bucephalus polymorphus*, *Raphidascaris acus*, *Argulus foliaceus* and *Ergasilus sieboldi*. The cestode *Eubothrium*, could be identified to family level since the typical scolex grooves had already developed.

From adult fish hosts the following species were identified: *Proteocephalus percae* from perch, *Proteocephalus longicollis* from white fish, *Eubothrium salvelini* from arctic char, *Eubothrium crassum* from burbot and *Triaenophorus nodulosus* from pike.

Parasite species identification via direct sequencing

Based on sequence analysis 5 different cestode and 5 different trematode (eye fluke) species were identified from various hosts (Table 2, Fig. 1).

Sequence analysis of D1-D2 fragments from the LSU rDNA gene region of cestodes from adult fish hosts revealed several mutational steps between the different parasite sequences (Supplementary Fig. S1, online version only), which resulted in

Table 2. Fish parasites analysed via direct sequencing from various host species of Lake Constance (LC)

(Added also is information on the typical invertebrate host species from the literature.)

Parasite species	Host species LC	No. parasites found	Location in host LC	Gene(s) sequenced	First intermediate host described	References
Cestodes						
<i>Proteocephalus percae</i>	Perch adult	10	Gut	LSU rDNA	<i>Cyclops</i> spp.	Scholz, 1999
	Perch fry	1	Gut		<i>Macrocyclus</i> spp.	
	Burbot	1	Gut		<i>Megacyclus</i> spp. <i>Eudiptomus</i> spp.	
<i>Proteocephalus longicollis</i>	Perch fry	13	Gut	LSU rDNA	<i>Cyclops</i> spp.	Scholz, 1999
	Whitefish	8	Gut		<i>Megacyclus</i> spp.	
	<i>Cyclops</i> spp.	15	Haemocoel		<i>Mesocyclops</i> spp. <i>Eudiptomus</i> spp.	
<i>Eubothrium crassum</i>	Perch fry	21	Gut	LSU rDNA	<i>Cyclops</i> spp.	Hanzelová <i>et al.</i> 2002; Kennedy, 1978
	Burbot	1	Gut			
	<i>Cyclops</i> spp.	1	Haemocoel			
<i>Eubothrium salvelini</i>	Arctic charr	5	Gut	LSU rDNA	<i>Cyclops</i> spp.	Hanzelová <i>et al.</i> 2002; Kennedy, 1978; Poulin <i>et al.</i> 1992
<i>Triaenophorus nodulosus</i>	Perch fry	2	Liver/gut	LSU rDNA	<i>Cyclops</i> spp.	Kuperman, 1981; Pronin, 1990
	Pike	1	Gut		<i>Eudiptomus</i> spp.	
	<i>Cyclops</i> spp.	3	Haemocoel			
Trematodes						
<i>Diplostomum baeri</i>	Perch adult	39	Eye vitreous	CO1/ITS1 + 2	<i>Radix auricularia</i>	Niewiadomska and Kiseliene, 1994
	Ruffe [#]	2	Eye vitreous			
<i>Diplostomum pseudospathaceum</i>	Ruffe [#]	1	Eye lens	CO1/ITS1 + 2	<i>Radix auricularia</i> <i>Radix labiata</i> <i>Lymnaea stagnalis</i> <i>Galba palustris</i>	Niewiadomska and Kiseliene, 1994; Niewiadomska and Laskowski 2002; Faltýnková, 2005
	<i>Radix labiata</i>	9	Cercaria release			
<i>Diplostomum spathaceum</i>	Roach*	2	Eye lens	CO1/ITS1 + 2	<i>Radix auricularia</i> <i>Radix labiata</i> <i>Lymnaea stagnalis</i>	Niewiadomska and Kiseliene, 1994; Vojtec, 1989
	<i>Radix auricularia</i>	1	Cercaria release			
<i>Diplostomum paracaudum</i>	Ruffe [#]	1	Eye lens	CO1/ITS1 + 2	<i>Radix auricularia</i> <i>Galba palustris</i> <i>Lymnaea stagnalis</i>	Niewiadomska and Szymanski, 1991; Faltýnková, 2005
	Whitefish [#]	1	Eye lens			
	<i>Radix auricularia</i>	1	Cercaria release			
<i>Tylodelphys</i> sp 2.	Perch adult	7	Eye vitreous	CO1/ITS1 + 2	<i>Radix auricularia</i> <i>Radix labiata</i>	<i>Tylodelphys</i> spp. : Vojtec, 1989; Faltýnková, 2005
	Whitefish [#]	5	Eye vitreous			
	<i>Radix auricularia</i>	5	Cercaria release			
	<i>Radix labiata</i>	3	Cercaria release			

* Only CO1 gene fragment sequenced.

Only ITS1 + 2 region sequenced.




Fig. 1. NJ trees from Kimura-2 parameter distances of trematodes and cestodes (1000 bootstraps, only bootstrap support >70% is given on the nodes). (A) CO1 sequences (388 bp fragment) of trematodes. (B) ITS1 sequences (577 bp fragment) of trematodes, grey box highlights species found in the eye lenses of fish hosts. Sequences from GenBank are marked in bold. (C) D1-D2 fragment of the LSU rDNA gene region (870 bp fragment) of cestodes.

clear clustering in the cestode NJ phenogram (Fig. 1C). However, no intraspecific variation was detected.

The comparison with published sequences from GenBank revealed only 1 accordance, a 100% match with a sequence published for *Eubothrium crassum*, GenBank Accession number AF286947, described by Olson *et al.* (2001) (Fig. 1). For all other cestode species, no good accordance could be found with sequences from specimens sampled in this study. Thus the sequences of both *Proteocephalus* species sampled in this study differed from those published for both *P. longicollis* (syn. *P. exiguus*) GenBank Accession number AJ388626.1 and *P. percae* GenBank Accession number AJ288594.1 described by Zehnder and Mariaux (1999). In both instances there were numerous substitutions (3 and 21 respectively) and sequence gaps (8 and 3 respectively). No sequences were available in GenBank for *Eubothrium salvelini* or *Triaenophorus nodulosus*.

Proteocephalus percae as identified from adult perch was also present in 1 juvenile perch from the pelagic zone and in burbot. *Proteocephalus longicollis*, identified from whitefish, also occurred in perch fry and in 15 copepods from the pelagic zone. *Eubothrium crassum*, identified in burbot, was also found in perch fry and 1 copepod. *Eubothrium salvelini* was identified from Arctic char and found only in this species. *Triaenophorus nodulosus* was identified from pike and also occurred in juvenile perch and 3 copepods from the pelagic zone (Table 2).

Sequencing a fragment of the CO1 gene and the rDNA ITS 1+2 region from trematodes obtained from various host species (Table 2) allowed further species identification based on numerous mutational differences between parasites (Supplementary Figs S2 and S3 respectively, online version only). While no intraspecific variation was found based on the alignment of the rDNA ITS 1+2 region, high intraspecific variation occurred when considering the alignment of the mitochondrial CO1 subunit (Fig. 1A, and Supplementary Fig. S2, online version only). A BLAST search based solely on the ITS 1 gene region of the *Diplostomum* specimen found in this study allowed specimens to be assigned to 4 species (*Diplostomum baeri* GenBank Accession number AF419274.1, *Diplostomum pseudospathaceum* GenBank Accession number AF419273.1, *Diplostomum spathaceum* GenBank Accession number AF419276.1 and *Diplostomum paracaudum* GenBank Accession number AF419272.1) described by Niewiadomska and Laskowski (2002). Accordance between the sequences found here and the sequences of Niewiadomska and Laskowski (2002) was between 100% (*Diplostomum pseudospathaceum*) and 99% (remaining 4 species) (Fig. 1B). It must be noted, that *Diplostomum spathaceum* and *Diplostomum parviventosum* (GenBank Accession number AF419277.1) cannot be separated according to their

ITS1 sequences (Niewiadomska and Laskowski, 2002). However, based on typical morphological characteristics of the cercariae (Niewiadomska and Kiseliene, 1994), *D. spathaceum* and *D. parviventosum* can be distinguished. *D. spathaceum* is in general larger than *D. parviventosum* (total length 717–887 µm for *D. spathaceum* and 529–645 µm for *D. parviventosum* respectively) and its acetabulum exceeds the width of the anterior organ whereas it is much smaller in *D. parviventosum*. The cercariae, shed by *R. auricularia* sampled in this study were 750 µm long and their acetabulum exceeded the anterior organ, thus they could be identified as *D. spathaceum* (Supplementary Fig. S4, online version only). All *Tylodelphys* specimens found could be assigned to 1 species. In contrast to the various ITS1 *Diplostomum* sequences available in GenBank, only a single *Tylodelphys* sequence was found *Tylodelphys* sp. GenBank Accession number AY386164. Our sequence differed from that published in GenBank by 3 base-pair substitutions, and thus we referred to it hereafter as *Tylodelphys* sp. 2 (Fig. 1B). In perch, only *Diplostomum baeri* and *Tylodelphys* sp. 2 were found (Tables 1 and 2). No mitochondrial CO1 sequences were available in GenBank for European trematodes and our sequences differed substantially from the North American *Diplostomum* species described by Locke *et al.* (2010a,b).

Chronology of infection and parasite community development

The Lake Constance perch fry sampled in this study exhibited a very specific chronology of parasite infection (Table 1). The first fish larvae caught in the pelagic zone in 2010 were 2 weeks old and appeared to be free of parasites. At 4 weeks of age, pelagic juvenile fish from the same cohort were infected with 3 parasite species: *Proteocephalus* (*P. percae* and *P. longicollis*), *Eubothrium crassum* and *Bunodera luciopercae*. The 7-week-old juveniles caught in 2009 in the pelagic zone were infected with the same parasite species as the 4-week-old fish in 2010 and, additionally, with *Triaenophorus nodulosus*. Perch caught in the littoral zone at 8 and 12 weeks of age (2009 cohort) were further infected with several species of digenean trematodes (Table 1). The eye flukes *Tylodelphys* sp. 2 and *Diplostomum baeri*, both found in the vitreous body (Table 2), reached high levels of prevalence, while 3 other species (*Cotylurus pileatus*, *Ichthyocotylurus variegatus* and *Hysteromorpha triloba*) occurred either at lower densities or were found only sporadically (*Bucephalus polymorphus*). In addition, 1 nematode, *Raphidascaris acus*, and 2 maxillopods, *Argulus foliaceus* and *Ergasilus sieboldi*, were found at low densities.

Parasite communities from fish of similar age clustered together in the nMDS plot, with younger fish occupying the left side of the graph and older

Table 3. Sampling details of invertebrate intermediate hosts from Lake Constance (The number of hosts investigated and the parasite prevalence in% (in parentheses) is given for cestodes (in *Cyclops* spp.) and trematodes of the family Diplostomatidae (in both *Radix* species).)

Intermediate host	Year	Total (n)	March	April	May	June	July	August	Sept.	Oct.
<i>Cyclops</i> spp.	2011	224	ns	ns	ns	224 (6.3)	ns	ns	ns	ns
<i>Cyclops</i> spp.	2012	3983	200 (1.5)	700 (0.1)	544 (1.5)	469 (4.3)	470 (5.3)	300 (13.3)	300 (14.6)	—
<i>Radix auricularia</i>	2010	1043	ns	ns	641 (0)	14 (0)	151 (5.3)	49 (2.0)	72 (4.1)	116 (1.7)
<i>Radix labiata</i>	2010	960	ns	ns	0 (-)	0 (-)	344 (2.6)	328 (1.8)	10 (0)	278 (0.4)

ns, no sampling.




Fig. 2. Two-dimensional nonmetric multidimensional scaling (nMDS) plot calculated using Bray-Curtis ordination of parasite abundance data of perch fry sampled in Lake Constance in 2009. Triangles denote perch, sampled at week 7, circles at week 8 and squares at week 12 post-hatch. Perch sampled at week 7 were caught in the pelagic, perch sampled at weeks 8 and 12 in the littoral zone of the lake.

perch appearing towards the right (Fig. 2). The stress value for the 2-dimensional nMDS analysis was 0.15 which indicated a reasonably good resolution for reliable interpretation of the data. The differences observed between the parasite communities of fish of the 3 age groups was significant based on ANOSIM (Global R = 0.69, P = 0.001). Parasite communities between fish caught at week 7 in the pelagic and week 8 in the littoral had more overlap (Fig. 2) and a quite low R value in the ANOSIM (R = 0.23, P = 0.001), whereas the parasite communities of the 7- and 8-week-old fish groups were very different from those from fish caught at week 12 in the littoral (ANOSIM: week 7 vs 12: R = 0.90, P = 0.001; week 8 vs 12: R = 0.86, P = 0.001).

Emergence of infective stages from intermediate hosts and transmission to perch fry

In total, 4207 cyclopoid copepods (*Cyclops* spp.) and 2003 snails (*Radix auricularia* and *R. labiata*) were analysed for infection with fish pathogenic species. The parasite prevalence varied substantially between sampling months (Table 3). In the pelagic *Cyclops* spp., cestode prevalence was very low from March to May (around 1%) but increased thereafter to 15% in September. In the littoral zone, no cercariae were observed hatching from snails before July, but from July onwards, prevalence varied between 1 and 5% until sampling ended in October. Genetic survey of parasite life-cycle stages indicated transmission of *Proteocephalus longicollis*, *Eubothrium crassum* and *Triaenophorus nodulosus* to perch fry via consumption of infected pelagic copepods, as all 3 species have been detected in both perch fry and *Cyclops* spp. (Table 2). *Proteocephalus percae* could not be found in

pelagic *Cyclops* spp. but was detected in littoral *Cyclops* spp. in a more recent survey in Lake Constance (sampled 4 July 2012, data not shown).

Of all the different trematode cercariae found in the littoral snails, transmission to perch was indicated only for *Tyloodelphys* sp.2 from *Radix auricularia* and *R. labiata* (Table 2). *Tyloodelphys* sp.2 was also found infecting whitefish. Unfortunately, the snail host for *Diplostomum baeri* has not been identified thus far but other transmission pathways between snail and fish intermediate hosts were detected for the first time in Lake Constance (Table 2). For example, *Diplostomum pseudospathaceum* cercariae from *Radix labiata* were found infecting ruffe, *Diplostomum spathaceum* cercariae from *Radix auricularia* were found infecting roach and *Diplostomum paracaudum* cercariae from *R. auricularia* were found infecting ruffe and whitefish.

DISCUSSION

Parasite succession and development of the parasite community

Perch fry in Lake Constance first encounter macro-parasites in the pelagic zone between 2 and 7 weeks p.h., when ingestion of zooplankton intermediate hosts leads to successive infection with helminths (cestodes and one trematode). Further evidence for this is the detection of infective parasite stages in pelagic copepods. After perch juveniles begin to return to the littoral zone at approximately 8 weeks p. h., they are exposed to direct infection by trematodes hatching from various littoral snail intermediate hosts, and to sporadic infection with nematodes and maxillopods.

The observed changes in parasite infracommunities seemed to be caused by different mechanisms. The first shift, between week 7 and week 8 is mainly influenced by the addition of new species. This nicely reflects the move from a pelagic to a littoral habitat. Week 7 fish are infected exclusively with parasites from pelagic intermediate hosts, while by week 8, host fish are additionally infected with trematodes stemming from littoral intermediate hosts. In contrast, the shift in parasite infracommunities between week 8 and week 12 is mainly characterized by intensification of existing infections, mainly of the littoral trematodes. Thus the parasite community of perch fry is influenced by both increasing age of perch and by ontogenetic habitat shift.

During the first days of life, perch larvae migrate from the littoral to the pelagic zone of the lake. In this study, the first larvae caught in the pelagic zone at the beginning of June were still in the yolk-sac stage. No larvae from this first sample were infected with macroparasites, indicating that either parasites were not encountered prior to the first habitat shift, or if that rapid mortality following direct and indirect

infections meant that parasitized larvae died before reaching the pelagial. The data also showed that the first encounter with macroparasites in the pelagial happens via ingestion of infected prey. Interestingly, prevalence of procercoid infection in pelagic copepods began to increase in June, which matches well with the arrival of perch fry in the pelagial.

First infections with trematodes from snail intermediate hosts occurred after the juvenile perch had migrated back to the littoral zone. This is not surprising, because the snail intermediate hosts do not occur in the pelagial, and cercariae (infective free-living stages of trematodes that hatch from snails) are relatively short-lived (max. 52 and 73 h for *Diplostomum* spp. and *Tyloodelphys* sp.2 respectively, own unpublished observations). Additionally, due to their small size, cercariae are not very mobile. Despite the possibility of passive transport with offshore water currents we do not expect many cercariae in the pelagial and thus infection of perch before they migrate back to the littoral zone is highly unlikely. Interestingly, the timing of first cercariae release from littoral snails matches very precisely with that of the return migration of perch to the littoral zone in mid-July.

Almost all parasite species that were found in larval and juvenile perch are typical of adult perch from various regions (Andersen, 1978; Andrews, 1979; Valtonen *et al.* 2003; Morozinska-Gogol, 2008) and some have also been previously observed in adult perch from Lake Constance (Balling and Pfeiffer, 1997a-d). However, the genetic analysis revealed 2 different *Proteocephalus* species in perch fry. *Proteocephalus percae* is a typical cestode parasite of perch, while *P. longicollis* (synonyms: *P. exiguuus*, *P. neglectus*) is more usually associated with salmoniform fish (reviewed by Scholz *et al.* 2007). Amazingly, however, our genetic analyses of a subset of cestodes from perch of the pelagic samples yielded only 1 specimen of *P. percae*. The remainder were all *P. longicollis*. We hypothesize that this prevalence of *P. longicollis* is caused by the consumption of infected plankton in the pelagial, which is the major habitat of the parasite's final host, *Coregonus wartmanni* (Reckel *et al.* 1999). Older perch, which are the definitive hosts of *P. percae*, are littorally oriented (Wang and Eckmann, 1994) and thus a majority of *P. percae*-infected copepods may be found in littoral zooplankton. However, prevalence and also intensity of *Proteocephalus* in perch fry decreased after 7 weeks p.h. Possible explanations for this finding include: prevalence and intensity of both *Proteocephalus* species rise and peak until infected zooplankton are no longer available and/or perch switch to other prey items, followed by a decrease due to (i) successive elimination of *P. longicollis* by the host's maturing immune system, (ii) deficient establishment of most parasite specimens as has been shown for a number of *Proteocephalus* species (reviewed in Scholz, 1999), or

(iii) intra- and interspecific competition between young worms in the host intestine whereby *P. percae* outcompetes *P. longicollis*. These 3 hypotheses are not mutually exclusive; however, the result is the complete loss of *P. longicollis* during the later stages of perch development. The last 2 explanations could similarly lead to the observed decrease in *Eubothrium crassum* in perch fry. The decrease in *Triaenophorus nodulosus* procercooids, however, is most likely the result of a decline in infected zooplankton in spring (Lahnsteiner *et al.* 2009) and the additional migration of procercooids to the host's liver and subsequent development into plerocercoids.

Identification of parasite species

In total, 17 parasite species were identified in this study, including 13 found in perch fry and four additional ones found in other fish or intermediate hosts sampled in Lake Constance. Referring to perch fry, only 9 of the 13 species could be identified unambiguously by morphological characteristics alone. But an approach using a combination of informative gene sequences, gained either from unambiguously identified adult parasites sampled from older fish (cestodes) or using previously published sequences in GenBank (trematodes), allowed the successful species determination of all remaining parasites except *Tylodelphys*. Without additional genetic information, we certainly would have failed in distinguishing both *Proteocephalus* species because no morphological difference was apparent in their procercooids. Similarly, the metacercariae of trematode eye flukes of several fish hosts could not have been identified based on morphology alone (Moszczyńska *et al.* 2009; Locke *et al.* 2010a, b). Here, it was the comparison of our sequences to those published in GenBank that allowed identification to the species level. However, reference sequences from macroparasites of aquatic hosts are still scarce in the databases. For trematodes, the mitochondrial CO1 sequences seem to be more useful for species determination than the rDNA sequences, because of their high inter- and lower intraspecific variability, and they have proved useful in the detection of cryptic species in North American Diplostomoidea (Moszczyńska *et al.* 2009; Locke *et al.* 2010a). The high variability between CO1 sequences found in this study, and the unambiguous clustering supported by high bootstrap values has helped delineate different species. Unfortunately few CO1 sequences from European species are yet available for comparative analysis, and there is a need for more elaborate sampling to be carried out in the future.

In conclusion, this study showed that the genetic connection of different parasite life-cycle stages within one ecosystem provides a powerful tool for the correct identification of parasite species. It may

also open the door to a new means of studying parasite transmission within ecosystems. Genetic surveys could provide independent confirmation of 'classical' life-cycle studies, such as experimental infection of various hosts. These and many further ecological, evolutionary and conservation scenarios can be imagined, in which accompanying genetic surveys provide new insights into parasite transmission pathways, parasite-host co-evolution or the spread of invasive parasite species.

ACKNOWLEDGEMENTS

The author wants to thank the following people for their help in data collection: Michael Donner, Dominik Geray and Daniela Harrer caught and analysed perch fry. Simon Weigl helped with parasite sampling from adult fish hosts and copepods and Martina Knauer sampled cercariae from snails during her diploma thesis. Julia Behr, Simon Weigl and Martina Knauer assisted with the molecular analysis. I am grateful to Reiner Eckmann, Martin Kalbe and Irene Samonte-Padilla for critical comments on the manuscript and to Amy-Jane Beer for proof reading. Additionally, the author wants to thank an anonymous referee whose comments greatly improved the manuscript.

FINANCIAL SUPPORT

This research was financially supported by the 'Stiftung für Umwelt und Wohnen' of the University of Konstanz.

REFERENCES

- Andersen, K. (1978). The helminths in the gut of perch (*Perca fluviatilis* L.) in a small oligotrophic lake in southern Norway. *Zeitschrift für Parasitenkunde* **56**, 17–27.
- Andrews, C. (1979). Host specificity of the parasite fauna of perch *Perca fluviatilis* L. from the British Isles, with special reference to a study at Llyn Tegid (Wales). *Journal of Fish Biology* **15**, 195–209.
- Balbuena, J. A., Karlsbakk, E., Kvenseth, A. M., Saksvik, M. and Nylund, A. (2000). Growth and emigration of third-stage larvae of *Hysterothylacium aduncum* (nematoda: anisakidae) in larval herring *Culpea havengus*. *Journal of Parasitology* **86**, 1271–1275.
- Balling, T. E. and Pfeiffer, W. (1997a). Frequency distribution of fish parasites in the perch *Perca fluviatilis* L. from Lake Constance. *Parasitology Research* **83**, 370–373.
- Balling, T. E. and Pfeiffer, W. (1997b). The parasitism of fish from Lake Constance: A comparison of present and earlier data. *Parasitology Research* **83**, 793–796.
- Balling, T. E. and Pfeiffer, W. (1997c). Seasonal differences in infestation of the perch *Perca fluviatilis* L. from Lake Constance with digenean trematodes. *Parasitology Research* **83**, 789–792.
- Balling, T. E. and Pfeiffer, W. (1997d). Location-dependent infection of fish parasites in Lake Constance. *Journal of Fish Biology* **51**, 1025–1032.
- Bertrand, M., Marcogliese, D. J. and Magnan, P. (2008). Trophic polymorphism in brook charr revealed by diet, parasites and morphometrics. *Journal of Fish Biology* **72**, 555–572.
- Bykhovskaya-Pavlovskaya, I., Gusev, A., Dubinina, M., Izyumova, N., Smirnova, T., Sokolovskaya, I., Shtein, G., Shul'man, S. and Epshtein, V. (1964). *Key to Parasites of Freshwater Fish of the U.S.S.R.* Akademiya Nauk SSSR.
- Campana, S. E. (1992). Measurement and interpretation of the microstructure of fish otoliths. In *Otolith Microstructure, Examination and Analysis* (ed. Stevenson, K. D. and Campana, S. E.). Canadian Special Publication of Fisheries and Aquatic Sciences. **117**, pp. 59–71.
- Carney, J. P., and Dick, T. A. (1999). Enteric helminths of perch (*Perca fluviatilis* L.) and yellow perch (*Perca flavescens* Mitchell): stochastic or predictable assemblages? *Journal of Parasitology* **85**, 785–795.
- Clarke, K. R. (1993). Non-parametric multivariate analyses of changes in community structure. *Australian Journal of Ecology* **18**, 117–143.

- Coles, T. (1981). The distribution of perch, *Perca fluviatilis* L. throughout their first year of life in Llyn Tegid, North Wales. *Archiv für Fischereiwissenschaft* **15**, 193–204.
- Criscione, C. D. and Blouin, M. S. (2004). Life cycles shape parasite evolution: comparative population genetics of salmon trematodes. *Evolution* **58**, 198–202.
- Eckmann, R. and Imbrock, F. (1996). Distribution and diel vertical migration of Eurasian perch (*Perca fluviatilis* L.) during winter. *Annales Zoologici Fennici* **33**, 679–686.
- Faltýnková, A. (2005). Laval trematodes (Digenea) in molluscs from small water bodies near Ceske Budejovice, Czech Republic. *Acta Parasitologica* **50**, 49–55.
- Faltýnková, A., Karvonen, A., Jyrkkä, M. and Valtonen, E. T. (2009). Being successful in the world of narrow opportunities: transmission patterns of the trematode *Ichthyocotylurus pileatus*. *Parasitology* **136**, 1375–1382.
- Galazzo, D. E., Dayanandan, S., Marcogliese, D. J. and McLaughlin, J. D. (2002). Molecular systematics of some North American species of *Diplostomum* (Digenea) based on rDNA-sequence data and comparisons with European congeners. *Canadian Journal of Zoology* **80**, 2207–2217.
- Gutter, A. S., Cribb, T. H., McCallum, H., Pickering, J. L. and McCormick, M. I. (2009). Effects of parasites on larval and juvenile stages of the coral reef fish *Pomacentrus moluccensis*. *Coral Reefs* **29**, 31–40.
- Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium* **41**, 95–98.
- Hanzelová, V., Scholz, T., Gerdeaux, D. and Kuchta, R. (2002). A comparative study of *Eubothrium salvelini* and *E. crassum* (Cestoda: Pseudophyllidea) parasites of arctic charr and brown trout in Alpine lakes. *Environmental Biology of Fishes* **64**, 245–256.
- Imbrock, F., Appenzeller, A. and Eckmann, R. (1996). Diel and seasonal distribution of perch in Lake Constance: a hydroacoustic study and *in situ* observations. *Journal of Fish Biology* **49**, 1–13.
- Johnson, M. W., Nelson, P. A. and Dick, T. A. (2004). Structuring mechanisms of yellow perch (*Perca flavescens*) parasite communities: host age, diet, and local factors. *Canadian Journal of Zoology* **82**, 1291–1301.
- Johnson, P. T. J., Stanton, D. E., Preu, E. R., Forshay, K. J. and Carpenter, S. R. (2006). Dining on disease: How interactions between infection and Environment affect predation risk. *Ecology* **87**, 1973–1980.
- Kelly, D. W., Thomas, H., Thielges, D. W., Poulin, R. and Tompkins, D. M. (2010). Trematode infection causes malformations and population effects in a declining New Zealand fish. *Journal of Animal Ecology* **79**, 445–452.
- Kennedy, C. R. (1978). The biology, specificity and habitat of the species of *Eubothrium* (Cestoda: Pseudophyllidea), with reference to their use as biological tags: a review. *Journal of Fish Biology* **12**, 393–410.
- Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* **16**, 111–120.
- King, S. D. and Cone, D. K. (2009). Infections of *Dactylogyrus pectenatus* (Monogenea: Dactylogyridae) on Larvae of *Pimephales promelas* (Teleostei: Cyprinidae) in Scott Lake, Ontario, Canada. *Comparative Parasitology* **76**, 110–112.
- Kruskal, J. B. (1964). Nonmetric multidimensional scaling: A numerical method. *Psychometrika* **29**, 115–129.
- Kuchta, R., Cech, M., Scholz, T., Soldanova, M., Levron, C. and Skorikova, B. (2009). Endoparasites of European perch *Perca fluviatilis* fry: role of spatial segregation. *Diseases of Aquatic Organisms* **86**, 87–91.
- Kuperman, B. I. (1981). *Tapeworms of the genus Triaenophorus*, Parasites of Fishes. Amerind Publishing, New York, USA.
- Lahnsteiner, F., Kletzl, M. and Weismann, T. (2009). The risk of parasite transfer to juvenile fishes by live copepod food with the example *Triaenophorus crassus* and *Triaenophorus nodulosus*. *Aquaculture* **295**, 120–125.
- Locke, S. A., Daniel McLaughlin, J. and Marcogliese, D. J. (2010a). DNA barcodes show cryptic diversity and a potential physiological basis for host specificity among Diplostomoidea (Platyhelminthes: Digenea) parasitizing freshwater fishes in the St. Lawrence River, Canada. *Molecular Ecology* **19**, 2813–2827.
- Locke, S. A., McLaughlin, J. D., Dayanandan, S. and Marcogliese, D. J. (2010b). Diversity and specificity in *Diplostomum* spp. metacercariae in freshwater fishes revealed by cytochrome c oxidase I and internal transcribed spacer sequences. *International Journal for Parasitology* **40**, 333–343.
- Marcogliese, D. J. (1995). The role of zooplankton in the transmission of helminth parasites to fish. *Reviews in Fish Biology and Fisheries* **5**, 336–371.
- McCairns, R. J. S. and Fox, M. G. (2004). Habitat and home range fidelity in a trophically dimorphic pumpkinseed sunfish (*Lepomis gibbosus*) population. *Oecologia* **140**, 271–279.
- Miehl, S. M. and Dettmers, J. M. (2011). Factors Influencing Habitat Shifts of Age-0 Yellow Perch in Southwestern Lake Michigan. *Transactions of the American Fisheries Society* **140**, 1317–1329.
- Morozynska-Gogol, J. (2008). A check-list of parasites of percid fishes (Actinopterygii: Percidae) from the estuaries of the Polish coastal zone. *Helminthologia* **45**, 196–203.
- Moszczyńska, A., Locke, S. A., McLaughlin, J. D., Marcogliese, D. J. and Crease, T. J. (2009). Development of primers for the mitochondrial cytochrome c oxidase I gene in digenetic trematodes (Platyhelminthes) illustrates the challenge of barcoding parasitic helminths. *Molecular Ecology Resources* **9**, 75.
- Nendick, L., Sackville, M., Tang, S., Brauner, C. J. and Farrell, A. P. (2011). Sea lice infection of juvenile pink salmon (*Oncorhynchus gorbuscha*): effects on swimming performance and postexercise ion balance. *Canadian Journal of Fisheries and Aquatic Sciences* **68**, 241–249.
- Niewiadomska, K. and Szymanski, S. (1991). Host-induced variability of *Diplostomum paracaudum* (Iles, 1959) metacercariae (Digenea). *Acta Parasitologica Warszawa* **36**, 11–17.
- Niewiadomska, K. and Kiseliene, V. (1994). *Diplostomum* cercariae (Digenea) in snails from Lithuania. II. Survey of species. *Acta Parasitologica Warszawa* **39**, 179–186.
- Niewiadomska, K. and Laskowski, Z. (2002). Systematic relationships among six species of *Diplostomum* Nordmann, 1832 (Digenea) based on morphological and molecular data. *Acta Parasitologica Warszawa* **47**, 20–28.
- Olson, P. D., Littlewood, D. T., Bray, R. A. and Mariaux, J. (2001). Interrelationships and evolution of the tapeworms (Platyhelminthes: Cestoda). *Molecular Phylogenetics and Evolution* **19**, 443–467.
- Pracheil, B. M. and Muzzall, P. M. (2009). Chronology and development of juvenile bluegill parasite communities. *Journal of Parasitology* **95**, 838–845.
- Pronin, N. M. (1990). Structure of the cestode population of *Triaenophorus nodulosus* (Pseudophyllidea, Triaenophoridae) in the ecosystem of Lake Shchuchic and the mortality of the helminth at various stages of development. *Ekologija. Lietuvos Mokslu Akademija Vilnius* **3**, 48–55.
- Reckel, F., Melzer, R. R. and Smola, U. (1999). Ultrastructure of the retina of two subspecies of *Coregonus lavaretus* (Teleostei) from Lake Constance (Germany). *Acta Zoologica* **80**, 153–162.
- Rosenthal, H. (1967). Parasites in larvae of the herring (*Clupea harengus* L.) fed with wild plankton. *Marine Biology* **1**, 10–15.
- Saitou, N. and Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* **4**, 406–425.
- Schäperclaus, W. (1979). *Fischkrankheiten*. Vol 4. Akademie-Verlag, Berlin, Germany.
- Scholz, T., Drabek, R. and Hanzelova, V. (1998). Scolex morphology of *Proteocephalus* tapeworms (Cestoda: Proteocephalidae), parasites of freshwater fish in the Palearctic Region. *Folia Parasitologica Praha* **45**, 27–43.
- Scholz, T. (1999). Life cycle of species of *Proteocephalus*, parasites of fishes in the Palearctic Region: a review. *Journal of Helminthology* **73**, 1–19.
- Scholz, T., Hanzelova, V., Skerikova, A., Shimazu, T. and Rolbiecki, L. (2007). An annotated list of species of the *Proteocephalus* Weinland, 1858 aggregate sensu de Chambrier et al. (2004) (Cestoda: Proteocephalidae), parasites of fishes in the Palearctic Region, their phylogenetic relationships and a key to their identification. *Systematic Parasitology* **67**, 139–156.
- Secombes, C. J. and Chappell, L. H. (1996). Fish immune responses to experimental and natural infection with helminth parasites. *Annual Review of Fish Diseases* **6**, 167–177.
- Skovgaard, A., Bahlool, Q. Z. M., Munk, P., Berge, T. and Buchmann, K. (2009a). Infection of North Sea cod, *Gadus morhua* L., larvae with the parasitic nematode *Hysterothylacium aduncum* Rudolphi. *Journal of Plankton Research* **33**, 1311–1316.
- Skovgaard, A., Meneses, I. and Angélico, M. M. (2009b). Identifying the lethal fish egg parasite *Ichthyodinium chabelardi* as a member of Marine Alveolate Group I. *Environmental Microbiology* **11**, 2030–2041.
- Sonnenberg, R., Nolte, A. W. and Tautz, D. (2007). An evaluation of LSU rDNA D1-D2 sequences for their use in species identification. *Frontiers in Zoology* **4**, 6.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. and Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* **28**, 2731–2739.
- Valtonen, E. T., Holmes, J. C., Aronen, J. and Rautalahti, I. (2003). Parasite communities as indicators of recovery from pollution: parasites of

- roach (*Rutilus rutilus*) and perch (*Perca fluviatilis*) in central Finland. *Parasitology* **126** (Suppl), S43–S52.
- Vojtek, J.** (1989). The present situation of the research into the stages of development of trematodes in Czechoslovakia. *Scripta Facultatis Scientiarum Naturalium Universitatis Prukynianae Brunensis. Brno* **19**, 339–352.
- Wang, N. and Eckmann, R.** (1994). Distribution of perch (*Perca fluviatilis* L.) during their first year of life in Lake Constance. *Hydrobiologia* **277**, 135–143.
- Wang, N. and Appenzeller, A.** (1998). Abundance, depth distribution, diet composition and growth of perch (*Perca fluviatilis*) and burbot (*Lota lota*) larvae and juveniles in the pelagic zone of Lake Constance. *Ecology of Freshwater Fish* **7**, 176–183.
- Weber, M. J., Dettmers, J. M. and Wahl, D. H.** (2011). Growth and survival of age-0 yellow perch across habitats in southwestern Lake Michigan: early life history in a large freshwater environment. *Transactions of the American Fisheries Society* **140**, 1172.
- Wootton, R.** (1974). Studies on the life history and development of *Proteocephalus percae* (Müller) (Cestoda: Proteocephalidea). *Journal of Helminthology* **48**, 269–281.
- Wootton, J. R.** (1998). *Ecology of Teleost Fishes*, 2nd Edn. Chapman and Hall, New York, USA.
- Zehnder, M. P. and Mariaux, J.** (1999). Molecular systematic analysis of the order Proteocephalidea (Eucestoda) based on mitochondrial and nuclear rDNA sequences. *International Journal for Parasitology* **29**, 1841–1852.
- Zelmer, D. A. and Arai, H. P.** (2004). Development of nestedness: host biology as a community process in parasite infracommunities of yellow perch (*Perca flavescens* (Mitchill)) from Garner Lake, Alberta. *Journal of Parasitology* **90**, 435–436.