

**EQUILIBRIA AND KINETICS IN THE CO-INDUCED, REVERSIBLE DISPROPORTIONATION OF  $((C_5H_5)_2TiCl)_2$  TO  $(C_5H_5)_2TiCl_2$  AND  $(C_5H_5)_2Ti(CO)_2$ . AN IR STUDY AT ELEVATED CO GAS PRESSURES**

EDITH U. VAN RAAIJ, C. DAVID SCHMULBACH \* and HANS H. BRINTZINGER \*\*

*Fakultät für Chemie, Universität Konstanz, 7750 Konstanz (F.R.G.)*

**Summary**

IR studies of toluene solutions of  $((C_5H_5)_2TiCl)_2$  under elevated CO pressures indicate the formation of a carbonyl complex, probably  $(C_5H_5)_2Ti(CO)Cl$ , with  $\nu(CO)$  2068  $cm^{-1}$ . This species predominates at CO pressures above 100 bar, and is reasonably stable at temperatures around  $-30^\circ C$ . At higher temperatures, disproportionation to  $(C_5H_5)_2Ti(CO)_2$  and  $(C_5H_5)_2TiCl_2$  occurs. This disproportionation is reversible; at  $+30^\circ C$  and 1 bar of CO, roughly equal concentrations of  $((C_5H_5)_2TiCl)_2$  and of dicarbonyl and dichloride products are in equilibrium with each other. Practically complete formation of these products occurs at CO pressures above 10 bar. Rate laws and reaction mechanisms have been studied for the disproportionation reaction and for its reverse, the comproportionation of  $(C_5H_5)_2Ti(CO)_2$  and  $(C_5H_5)_2TiCl_2$  to  $((C_5H_5)_2TiCl)_2$ , which predominates at lower CO pressures.

**Introduction**

The comproportionation of the  $Ti^{IV}$  and  $Ti^{II}$  derivatives  $Cp_2TiCl_2$  and  $Cp_2Ti(CO)_2$  ( $Cp = \eta^5-C_5H_5$ ) has been shown to be a useful method for the preparation of the  $Ti^{III}$  derivative  $(Cp_2TiCl)_2$  [1]; the CO-induced disproportionation of  $(Cp_2TiCl)_2$ , on the other hand, has been reported to occur in the reduction of  $Cp_2TiCl_2$  to  $Cp_2Ti(CO)_2$  by metals such as Zn or Al [2,3]. We describe here the results of an IR study, conducted at elevated CO pressures by use of an internal-circulation IR cell, of the equilibria and kinetics in the reaction of  $(Cp_2TiCl)_2$  with CO, and of intermediates participating in this reaction system.

\* Department of Chemistry and Biochemistry, S. Illinois University, Carbondale, Illinois 62901 (U.S.A.).

\*\* To whom correspondence should be addressed.

## Experimental

Dimeric titanocene monochloride,  $(Cp_2TiCl)_2$ , free of zinc halide contamination, was prepared by a published procedure [4]. All materials were handled with strict exclusion of air. Solids were transferred and weighed in an inert atmosphere glovebox, and solutions were prepared and transferred by standard vacuum-line techniques.

For IR measurements under elevated CO pressures, a pressure-resistant IR cell was constructed in which isobaric and isothermal circulation of gas-saturated solutions between a gas-liquid mixing chamber and the IR window gap is maintained by means of an internal, magnetically driven centrifugal rotor (Fig. 1). The cell block, manufactured from stainless steel (DIN 14571), was thermally insulated by use of foam-rubber linings inside a polyvinyl casing; its temperature was controlled within  $\pm 1^\circ C$  by a circulation thermostat with isopropanol as heat-transfer liquid. A solution volume of 30–50 ml is required for operation of the cell.

IR absorbance data were collected by means of a double-beam IR spectrophotometer (Perkin-Elmer 398) and transferred to a microcomputer for further processing. Raw absorbance values thus obtained were scaled to constant path length, using the toluene absorption band at  $1802\text{ cm}^{-1}$  as a standard, to compensate for increases in optical path lengths at elevated CO pressures. The relationship  $l = A_{1802} \cdot 0.0329\text{ cm}$ , derived from absorbance measurements in calibrated IR cells [6], was used to calculate the cell path length  $l$ . The solvent spectrum was then subtracted to give the spectra of the species in solution.




Fig. 1. Pressure-resistant IR cell with internal, isobaric and isothermal circulation. Essential parts: 1, inner, pressure-resistant  $CaF_2$  windows (13 mm thick, 18 mm diam.); 2, channels for thermostatted heat-transfer liquid; 3, internal centrifugal rotor, driven by rotating external magnet bar; 4, inlet/outlet for sample solutions and gases; 5, inlet/outlet for heat transfer liquid; 6, evacuation of space between inner and outer  $CaF_2$  windows; 7, removable foam-rubber insulation; 8, polyvinyl casing; 9, outer  $CaF_2$  windows. Direction of circulation of the sample solution between rotor-agitated gas-liquid mixing chamber, solution reservoir and IR window gap as indicated by small arrows.

TABLE 1  
SOLUBILITY DATA FOR CO IN TOLUENE AT ELEVATED CO PRESSURES

Temp. (°C)	Mol fractions of CO ( $\times 10^2$ ) at $P_{\text{CO}} =$				$d_{\text{tol}}^a$	
	20	60	100	138 bar		
20	1.49(7)	4.38(7)	7.14(11)	9.38(3)	0.8614	
0	1.43(4)	4.24(4)	6.79(14)	8.64(3)	0.8845	
-40	1.37(5)	4.02(3)	5.99(5)	7.73(6)	0.9212	
Temp. (°C)	Concentration of CO (mol/l) at $P_{\text{CO}} =$				$\alpha (\times 10^3)$	$\beta (\times 10^8)$
	20	60	100	138 bar		
20	0.141	0.428	0.719	0.967	7.1	0.3
0	0.139	0.425	0.699	0.908	7.1	2.3
-40	0.139	0.419	0.637	0.837	7.1	5.7

<sup>a</sup> Derived from data given in Internat. Critical Tables, Vol. III, E.W. Washburn (Ed.), New York 1928, p. 29 and in K. Schäfer and G. Beggerow, (Eds.), Landolt-Börnstein, Zahlenwerte und Funktionen, Vol. II., Berlin 1971, part 1, p. 641.

The concentration-proportionality of the absorbance values obtained in this way was checked for solutions of acetonitrile in toluene; absorbances at  $2254 \text{ cm}^{-1}$  of 1.0 to 6.0 M  $\text{CH}_3\text{CN}$  solutions obeyed the Lambert-Beer law up to absorbances above 1, at both 20 and  $-40^\circ\text{C}$ .

The solubility of CO in toluene at elevated CO pressures, for which we could not find useful data in the literature, was determined by gas-volumetric measurements [7]. In a small (40 ml) autoclave, thermally controlled within  $\pm 1^\circ\text{C}$  by means of a circulation thermostat, toluene was saturated with CO by several hours' stirring with a magnetic stirring bar, the CO pressure  $P_{\text{CO}}$  being read from a pressure transducer with an accuracy of ca.  $\pm 1$  bar. The CO-saturated toluene solutions were drained, by way of a dip-tube, into a vacuum manifold, where the toluene fractions (about 2 to 10 g) were condensed, at  $-80^\circ\text{C}$ , into a cold trap and weighed, while the CO released was transferred, by means of a Töpler pump, into a calibrated reservoir connected to a mercury manometer, and its volume determined.

After discarding of a sufficient forerun, four to six successive samples were measured at each temperature and CO pressure. The CO solubilities determined in this manner are presented in Tab. 1 as mol fractions  $X_{\text{CO}}$  with estimated mean deviations for the last digit indicated in parentheses. These solubility data were analysed, as recommended [7], with respect to the Krichevsky-Kasarnosky equation [8],  $\ln(f_{\text{CO}}/X_{\text{CO}}) = \ln(K_{\text{CO, tol}}) + V_{\text{CO}} \cdot P_{\text{CO}}/(RT)$  using, as a substitute for the unknown fugacities  $f_{\text{CO}}$  of CO in the equilibrium gas phase, fugacity values of pure, gaseous CO [9]. Plots of  $\ln(f_{\text{CO}}/X_{\text{CO}})$  vs.  $P_{\text{CO}}$  at constant temperature reveal substantial curvature. Whether these variations in  $d(\ln(f_{\text{CO}}/X_{\text{CO}}))/d(P_{\text{CO}})$  do, indeed, indicate changes in the partial volume  $V_{\text{CO}}$  of CO in these solutions, or are rather due to some systematic error either in our solubility data or in the fugacity data used, is at present an open question.

The utilisation of the CO solubility data represented in Tab. 1 for the calculations of equilibrium and rate constants, comparable with those for other, non-volatile ligand species, requires a conversion from mol fractions  $X_{\text{CO}}$  to concentrations  $C_{\text{CO}}$  in mol/l. For this purpose, it is necessary to know the volumes of the CO solutions formed from 1 mol of toluene. As an approximation, we have used the

TABLE 2

MOLAR EXTINCTION COEFFICIENTS  $\epsilon(\text{CO})$  AND HALF-WIDTHS  $\Delta\nu$  OF THE IR BAND AT  $2134 \text{ cm}^{-1}$  OF CO IN TOLUENE SOLUTION

Temp. ( $^{\circ}\text{C}$ )	20	0	-20	-40
$\epsilon(\text{CO})$ ( $\text{l mol}^{-1} \text{cm}^{-1}$ )	37.1(8)	40.0(9)	41.6(6)	45.8(9)
$\Delta\nu$ ( $\text{cm}^{-1}$ )	55	40	32	26

molar volume of the toluene solvent at the relevant temperature and at ambient pressure: At  $20^{\circ}\text{C}$  and  $P_{\text{CO}}$  100 bar, for instance, the compressibility of toluene would give rise to a volume decrease by ca. 1%, while the volume of the dissolved CO (as a liquid) would amount to ca. 2% of that of the solvent. Since the partial volume of CO in these solutions is undoubtedly smaller than the liquid volume of CO, these two effects would each be smaller than the estimated data error and would, furthermore, tend to cancel each other out. The CO concentrations  $C_{\text{CO}}$  in Tab. 1 were, therefore, derived from the corresponding mol fractions  $X_{\text{CO}}$  by the relationship  $C_{\text{CO}} = X_{\text{CO}} \cdot d_{\text{tol}} \cdot 1000 / ((1 - X_{\text{CO}}) \cdot 92.15)$ , where 92.15 is the molar mass of toluene and  $d_{\text{tol}}$  is its density at the temperature indicated, at ambient pressure.

The CO solubility data thus obtained show an evident saturation at higher CO pressures, which is particularly pronounced at lower temperatures. At each temperature, the data are reproduced, within experimental error, by the two-term equation  $C_{\text{CO}} = P_{\text{CO}} \cdot \alpha - (P_{\text{CO}})^3 \cdot \beta$ , with values of  $\alpha$  and  $\beta$  as indicated in Tab. 1. Since  $\alpha$  is essentially independent of temperature and since  $\beta$  appears to decrease practically linearly with increasing temperature, the empirical relation  $C_{\text{CO}} = P_{\text{CO}} \cdot 7.1 \cdot 10^{-3} - (P_{\text{CO}})^3 \cdot (2.2 - \delta \cdot 9.0 \cdot 10^{-2}) \cdot 10^{-8}$  with  $\delta$  being the temperature in  $^{\circ}\text{C}$ , was used to obtain the CO concentrations in these solutions from known CO pressures  $P_{\text{CO}}$ .

As a check on these data, IR absorbances of dissolved CO in CO-saturated toluene solutions were measured, at temperatures between  $-40$  and  $20^{\circ}\text{C}$ , over a range of CO saturation pressures between 20 and 130 bar. The CO absorbances at  $2134 \text{ cm}^{-1}$  thus obtained show, at any given temperature, the same saturation effects as the concentration data in Tab. 1. The rather satisfactory proportionality between these CO absorbances and the CO concentration values calculated for each of these solutions from the dependence of  $C_{\text{CO}}$  on  $P_{\text{CO}}$  derived above provides further confirmation of the reliability of this relation, which was used to convert  $P_{\text{CO}}$  to  $C_{\text{CO}}$  values for the subsequent calculations of equilibrium and rate constants.

The molar extinction coefficient  $\epsilon_{\text{CO}}$  of these CO solutions decreases noticeably with increasing temperature, together with a significant broadening of its IR band (Tab. 2). The relation  $\epsilon_{\text{CO}} = 40.0 - 2.9 \cdot \delta$  was used to obtain, at a given temperature  $\delta$  in  $^{\circ}\text{C}$ , the extinction coefficient  $\epsilon_{\text{CO}}$  necessary to check the equilibrium concentration of CO in a reaction mixture by means of its CO absorbance at  $2134 \text{ cm}^{-1}$ .

## Results and discussion

Upon admission of higher CO pressures to  $(\text{Cp}_2\text{TiCl})_2$  in toluene solution, particularly at temperatures below  $0^{\circ}\text{C}$ , a new IR absorption band appears at 2068




Fig. 2. IR spectra of a 0.025 *M* solution of  $(\text{Cp}_2\text{TiCl})_2$  in toluene, at  $-5^\circ\text{C}$ , 15 to 285 min after admission of 130 bar of CO.

$\text{cm}^{-1}$ . This IR band is fully formed within the 4–5 min required for the first measurement; in the course of time it decays again, as a simultaneous increase of IR absorption bands at 1966 and 1883  $\text{cm}^{-1}$  indicates the formation of the  $\text{Ti}^{\text{II}}$  complex  $(\text{Cp}_2\text{Ti}(\text{CO})_2$  (Fig. 2).

The reaction of  $(\text{Cp}_2\text{TiCl})_2$  with CO was first studied at  $-30^\circ\text{C}$ , where the secondary reaction is sufficiently slow to allow the new IR band at 2068  $\text{cm}^{-1}$  to remain unchanged for several hours. The changes of its intensity with CO pressure – about 25–35 bar of CO are required for half-formation of this species – and with initial concentrations of the dimer  $(\text{Cp}_2\text{TiCl})_2$  indicate that the carbonyl complex responsible for this IR band is the species  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$ , formed according to equilibrium (1).



Least-squares fitting of the IR data \* yields an equilibrium constant  $K_1 = 0.65 \pm 0.1$  l/mol for reaction (1) at  $-30^\circ\text{C}$  and a molar extinction coefficient  $\epsilon(\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}) = 360 \pm 20$  l  $\text{mol}^{-1}$   $\text{cm}^{-1}$ .

At  $-5^\circ\text{C}$ , where the ensuing disproportionation is still slow enough to allow the determination of initial IR absorbances at 2068  $\text{cm}^{-1}$  by extrapolation back to the actual time of CO admission, a value of  $K_1 = 0.25 \pm 0.05$  l/mol is found for

\* The IR absorbance at 2068  $\text{cm}^{-1}$  had to be corrected for a fringe contribution from the IR band of dissolved CO (cf. Fig. 2).

TABLE 3

FIRST-ORDER RATE CONSTANTS  $k_{\text{app}}$  AND  $k_{\text{dis}}$  FOR DISPROPORTIONATION OF  $(\text{Cp}_2\text{TiCl})_2$  IN DEPENDENCE OF CO PRESSURES, AT  $+30^\circ\text{C}$

$P_{\text{CO}}$ (bar)	5	7	9	10	14
$C_{\text{CO}}$ (mol/l)	0.035	0.050	0.064	0.071	0.099
$k_{\text{app}}$ ( $\text{min}^{-1}$ )	2.1	2.9	4.3	4.0	$7.7 \times 10^{-2}$
$k_{\text{dis}}$ ( $1 \text{ mol}^{-1} \text{ min}^{-1}$ )	5.7	5.6	6.7	5.6	$7.7 \times 10^{-1}$

reaction (1) if the value of  $\epsilon(\text{Cp}_2\text{Ti}(\text{CO})\text{Cl})$  previously determined at  $-30^\circ\text{C}$  is used to analyse the IR absorption data. This temperature dependence of  $K_1$  yields estimates of  $\Delta H^0 = -20 \pm 5 \text{ kJ}$  and  $\Delta S^0 = -85 \pm 20 \text{ J/K}$  for reaction (1).

The species  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$ , responsible for the IR band at  $2068 \text{ cm}^{-1}$ , is closely related to the more stable  $\text{Ti}^{\text{III}}$  carbonyl complexes  $\text{Cp}_2\text{Ti}(\text{CO})\text{C}_6\text{F}_5$  and  $(\text{C}_5(\text{CH}_3)_5)_2\text{Ti}(\text{CO})\text{Cl}$ , for which Teuben and coworkers have reported IR bands at  $2060$  and  $2000 \text{ cm}^{-1}$ , respectively [10].

The CO-induced disproportionation of  $(\text{Cp}_2\text{TiCl})_2$  to  $\text{Cp}_2\text{Ti}(\text{CO})_2$  and  $\text{Cp}_2\text{TiCl}_2$  was studied at  $+30^\circ\text{C}$ , where both this reaction and its reverse occur at conveniently measurable rates. After equilibration, the IR absorptions at  $1966$  and  $1883 \text{ cm}^{-1}$  depend on CO pressure in a manner which is in accord with the existence of equilibrium (2).



At a CO pressure of 1 bar, about one half of the  $\text{Ti}^{\text{III}}$  dimer is converted to the dicarbonyl and dichloride products at a total Ti concentration of  $[\text{Ti}]_{\text{tot}} = 0.05 \text{ M}$ ; above 10–15 bar of CO this conversion is essentially complete\*. For reaction (2) at  $30^\circ\text{C}$ , an equilibrium constant  $K_2 = 207 \pm 30 \text{ l/mol}$  is obtained if the rather low solubility of  $\text{Cp}_2\text{TiCl}_2$  in toluene, which we found to be  $9.8 \times 10^{-3} \text{ mol/l}$  at  $+30^\circ\text{C}$ , is assumed to keep this species at constant concentration.

The kinetics of the CO-induced disproportionation reaction was studied, at  $+30^\circ\text{C}$ , by following the increase in IR absorbances at  $1966$  and  $1883 \text{ cm}^{-1}$  caused by an increase in CO pressure from  $P_{\text{CO}} = 1$  bar to the value indicated in Tab. 3. The pattern of the CO-induced growth of these absorptions of  $\text{Cp}_2\text{Ti}(\text{CO})_2$  with time corresponds with an apparent first-order dependence on  $[(\text{Cp}_2\text{TiCl})_2]$ . The apparent first-order rate constants for the CO-induced disproportionation increase approximately linearly with CO concentration (Tab. 3), so that the empirical rate law would be given by eq. 3.

$$V_{\text{dis}} = k_{\text{dis}} \cdot [(\text{Cp}_2\text{TiCl})_2] \cdot [\text{CO}] \quad (3)$$

In order to satisfy the equilibrium condition (2), the rate of the reverse, disproportionation reaction must be in accord with eq. 4.


$$V_{\text{com}} = k_{\text{com}} \cdot [\text{Cp}_2\text{Ti}(\text{CO})_2] \cdot [\text{Cp}_2\text{TiCl}_2] / [\text{CO}] \quad (4)$$

\* The complete conversion of  $(\text{Cp}_2\text{TiCl})_2$  in  $\text{Cp}_2\text{Ti}(\text{CO})_2$  and  $\text{Cp}_2\text{TiCl}_2$  at  $P_{\text{CO}} > 15$  bar was confirmed by comparison with calibrated solutions of  $\text{Cp}_2\text{Ti}(\text{CO})_2$ . Battaglia et al. (ref. 3) had previously observed an incomplete conversion in THF solution at  $P_{\text{CO}} = 1$  bar, and ascribed this to formation of a complex between  $(\text{Cp}_2\text{TiCl})_2$  and  $\text{Cp}_2\text{TiCl}_2$ .

If the concentration of  $[\text{Cp}_2\text{TiCl}_2]$  is again assumed to be fixed by its solubility limit at  $9.8 \times 10^{-3}$  mol/l, eq. 4 would predict a first-order dependence of the disproportionation rate on  $[\text{Cp}_2\text{Ti}(\text{CO})_2]$ . A first-order decrease of  $\text{Cp}_2\text{Ti}(\text{CO})_2$  absorbances at 1966 and 1883  $\text{cm}^{-1}$  was indeed observed when the CO pressure was lowered, from ca. 70 bar, where equilibrium (2) lies far to the right, to 1 bar, where  $(\text{Cp}_2\text{TiCl})_2$ ,  $\text{Cp}_2\text{Ti}(\text{CO})_2$  and  $\text{Cp}_2\text{TiCl}_2$  are present in comparable concentrations.

In determining  $k_{\text{dis}}$  and  $k_{\text{com}}$  from these data, it is necessary to take into account the fact that both disproportionation and comproportionation were studied under conditions where these reactions approach equilibrium, so that the apparent first-order rate constants  $k_{\text{app}}$  derived, e.g., from a plot of  $\ln([A]_t - [A]_e)/([A]_0 - [A]_e)$  vs.  $t$  contain additive contributions from forward and backward reaction rates,  $k_{\text{app}} = k_{\text{frw}} + k_{\text{bkw}}$ , with  $k_{\text{frw}}$  and  $k_{\text{bkw}}$  being the overall first-order rate constants for each of these two reactions, for which eqs. 3 and 4 yield  $k_{\text{frw}} = k_{\text{dis}} \cdot [\text{CO}]$  and  $k_{\text{bkw}} = k_{\text{com}} \cdot [\text{Cp}_2\text{TiCl}_2]/[\text{CO}]$ . With  $K_2 = k_{\text{dis}}/k_{\text{com}}$ , we can determine  $k_{\text{dis}}$  and  $k_{\text{com}}$  from  $k_{\text{app}}$  as  $k_{\text{dis}} = k_{\text{app}} \cdot K_2 \cdot [\text{CO}]/(K_2 \cdot [\text{CO}]^2 + [\text{Cp}_2\text{TiCl}_2])$  and  $k_{\text{com}} = k_{\text{app}} \cdot [\text{CO}]/(K_2 \cdot [\text{CO}]^2 + [\text{Cp}_2\text{TiCl}_2])$ . Reasonably constant values of  $k_{\text{dis}} = 6.3 \pm 0.8 \cdot 10^{-1}$  (l/mol)  $\cdot$  min $^{-1}$  are thus obtained from the data in Tab. 3, i.e. in the range of 5 to 14 bar of CO. From  $k_{\text{app}} = 9.8 \pm 3.4 \cdot 10^{-3}$  min $^{-1}$ , measured for the reverse reaction at  $P_{\text{CO}} = 1$  bar, one obtains  $k_{\text{com}} = 3.4 \pm 1.2 \cdot 10^{-3}$  min $^{-1}$ .

A central question concerning the course of the CO-induced disproportionation of  $(\text{Cp}_2\text{TiCl})_2$  involves the role of the monocarbonyl complex  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$ . The first-order increase of the disproportionation rate with  $C_{\text{CO}}$  suggests a rate-determining step for this reaction in which the monocarbonyl complex  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$  reacts with monomeric  $\text{Cp}_2\text{TiCl}$  (formed by dissociation of  $(\text{Cp}_2\text{TiCl})_2$ ) according to eq. 5,



with  $\text{Cp}_2\text{Ti}(\text{CO})$  then taking up a second CO in a fast subsequent step. Such a reaction sequence would obey the overall rate law (eq. 6).

$$V_{\text{dis}} = k_6 \cdot [\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}] \cdot [\text{Cp}_2\text{TiCl}] \quad (6)$$

The concentrations of the two  $\text{Ti}^{\text{III}}$  species in eq. (6) are determined by equilibrium reaction (7); the product of their concentrations would, therefore, be given by  $[\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}] \cdot [\text{Cp}_2\text{TiCl}] = K_7 \cdot [(\text{Cp}_2\text{TiCl})_2] \cdot [\text{CO}]$ . Equation 6 is thus equivalent to the observed rate law (3) as long as the CO pressures are low enough to keep equilibrium (7) essentially on the left hand side, i.e. as long as  $(\text{Cp}_2\text{TiCl})_2$  is the predominant  $\text{Ti}^{\text{III}}$  species present.



At higher CO pressures, where the formation of  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$  approaches completion, the involvement of this species in the rate-determining step should cause the disproportionation rate  $V_{\text{dis}}$  to level off as  $C_{\text{CO}}$  is increased further. With  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$  becoming the predominant  $\text{Ti}^{\text{III}}$  species, the disproportionation reaction should, furthermore, approach a second-order rate law with respect to the total, residual  $\text{Ti}^{\text{III}}$  concentration  $[\text{Ti}]_{\text{tot}}$ . In order to clarify these aspects, the dispro-

\* The expressions  $[A]_t$ ,  $[A]_0$  and  $[A]_e$  refer to the concentrations of the decaying species ( $(\text{Cp}_2\text{TiCl})_2$  for disproportionation, and  $\text{Cp}_2\text{Ti}(\text{CO})_2$  for comproportionation reactions) at time  $t$ , for zero reaction, and after establishment of equilibrium, respectively.




Fig. 3. Disproportionation of  $(Cp_2TiCl)_2$  ( $[Ti]_{tot} = 0.05 M$  in toluene) at  $-5^\circ C$  under CO pressures of 20 and 140 bar as measured by absorbance  $A$  at  $1883\text{ cm}^{-1}$ ;  $A_{max}$ :  $A$  at time  $t > 1000$  min. A: First-order plot ( $\ln(1-A/A_{max})$  vs.  $t$ ); solid line: first-order reaction with  $k = 7.1 \times 10^{-3}\text{ min}^{-1}$ . B: Second-order plot ( $1/(1-A/A_{max})$  vs.  $t$ ); solid line: second-order reaction with  $k = 1.8 \times 10^{-2}\text{ min}^{-1}$ .

TABLE 4

INITIAL RATES  $V_{\text{dis}}$  FOR THE DISPROPORTIONATION OF  $(\text{Cp}_2\text{TiCl})_2$ , IN DEPENDENCE OF THE CONCENTRATIONS OF CO, AT  $-5^\circ\text{C}$

$P_{\text{CO}}$ (bar)	20	40	80	130	140
$C_{\text{CO}}$ (mol/l)	0.142	0.283	0.554	0.865	0.918
$[\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}]$	0.0100	0.0179	0.0286	0.0361	0.0370 <sup>a</sup>
$[(\text{Cp}_2\text{TiCl})_2]$	0.0200	0.0160	0.0107	0.0070	0.0065 <sup>b</sup>
$[\text{Cp}_2\text{TiCl}]/K_9^{1/2}$	0.141	0.126	0.103	0.083	0.081 <sup>c</sup>
$V_{\text{dis}}$ (calcd.) <sup>d</sup>	1.4	2.3	2.9	3.0	$3.0 (\times 10^{-4})$
$V_{\text{dis}}$ (exptl.) <sup>e</sup>	1.8	2.1	2.7	3.0	$3.0 (\times 10^{-4})$

<sup>a</sup>  $[\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}] = -A/2 + 1/2 \cdot (A^2 + 4 \cdot A \cdot [\text{Ti}]_{\text{tot}})^{1/2}$  mol/l, with  $A = K_1 \cdot [\text{CO}]^2/2$ ,  $K_1 = 0.25$  l/mol and  $[\text{Ti}]_{\text{tot}} = 0.05$  mol/l. <sup>b</sup>  $[(\text{Cp}_2\text{TiCl})_2] = ([\text{Ti}]_{\text{tot}} - [\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}])/2$ . <sup>c</sup>  $[\text{Cp}_2\text{TiCl}]/K_9^{1/2} = ([(\text{Cp}_2\text{TiCl})_2])^{1/2}$  (mol/l)<sup>1/2</sup>. <sup>d</sup> According to eq. 8 with  $k_6 \cdot K_9^{1/2} = 0.10$ ; in (mol/l) min<sup>-1</sup>. <sup>e</sup> Derived from initial slope,  $k_{\text{ini}}$ , of  $\ln(1 - A_t/A_{\text{max}})$  vs.  $t$  by  $V_{\text{dis}} = k_{\text{ini}} \cdot [(\text{Cp}_2\text{TiCl})_2]_0 = k_{\text{ini}} \cdot [\text{Ti}]_{\text{tot}}/2$  (mol/l) min<sup>-1</sup>.

portionation kinetics were studied in the  $P_{\text{CO}}$  range of 20 to 140 bar, at a temperature of  $-5^\circ\text{C}$ , where reaction rates were conveniently measurable even at higher CO pressures, and where the rate of the comproportionation reaction had been found to be negligible. It is apparent from Fig. 3 that the disproportionation still closely adheres to first-order kinetics at  $P_{\text{CO}} = 20$  bar, whereas at 140 bar the change to second-order kinetics appears to be practically complete. The initial disproportionation rates,  $V_{\text{dis}}$ , determined under these conditions (Tab. 4) increase indeed much less than proportionally to  $C_{\text{CO}}$ , and become practically independent of  $C_{\text{CO}}$  at higher CO concentrations.

The rate data in Tab. 4 were analysed in terms of their agreement with rate equation (6) by calculating concentrations of  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$  from equilibrium (1) using a value of  $K_1 = 0.25$  l/mol, as found at  $-5^\circ\text{C}$ , and concentrations of  $\text{Cp}_2\text{TiCl}$  from the residual concentrations of  $(\text{Cp}_2\text{TiCl})_2$  as  $[\text{Cp}_2\text{TiCl}] = (K_9 \cdot [(\text{Cp}_2\text{TiCl})_2])^{1/2}$  (cf. eq. 8) \*;  $K_9$  is the dissociation constant of  $(\text{Cp}_2\text{TiCl})_2$  according to eq. 9.


$$V_{\text{dis}} = k_6 \cdot [\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}] \cdot (K_9 \cdot [(\text{Cp}_2\text{TiCl})_2])^{1/2} \quad (8)$$



Since neither  $K_7$  nor  $K_9$  are known under the reaction conditions used \*\*, we can only determine the product  $k_6 \cdot K_9^{1/2}$  from the rate data given in Tab. 4. The acceptable agreement of the experimental data for  $V_{\text{dis}}$  with the values for  $V_{\text{dis}}$  calculated from eq. 8 with a mean value of  $k_6 \cdot K_9^{1/2} = 0.10 \pm 0.02$  (l/mol)<sup>1/2</sup> · min<sup>-1</sup>, in particular their levelling-off at higher CO concentrations, together with the concomitant change from first-order to second-order kinetics, represent reasonable evidence that a bimolecular reaction between  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$  and monomeric  $\text{Cp}_2\text{TiCl}$  according to eq. 6 is, indeed, rate-determining for the CO-induced disproportionation of  $(\text{Cp}_2\text{TiCl})_2$ .

\* For this we assume that only a small fraction of  $(\text{Cp}_2\text{TiCl})_2$  is dissociated into  $\text{Cp}_2\text{TiCl}$  in these solutions, as indicated by a value of  $2.6 \cdot 10^{-6}$  mol/l reported for  $K_9$  in toluene at  $80^\circ\text{C}$  (ref. 11).

\*\*  $K_1$ ,  $K_7$  and  $K_9$  are interconnected by the relation  $K_7 = (K_1 \cdot K_9)^{1/2}$ .



SCHEME 1. Intermediates in the CO-induced interconversion between  $(\text{Cp}_2\text{TiCl})_2$ ,  $\text{Cp}_2\text{Ti}(\text{CO})_2$  and  $\text{Cp}_2\text{TiCl}_2$ . Rate-determining step indicated by broken arrows ( $\rightleftharpoons$ ).

Such a reaction could occur either via an outer-sphere electron transfer to yield  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}^-$  and  $\text{Cp}_2\text{TiCl}^+$ , from which the final products could then be formed by rapid ligand exchange, or via an inner-sphere halogen transfer from  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$  to  $\text{Cp}_2\text{TiCl}$ . Electrochemical studies on the oxidation of  $\text{Cp}_2\text{TiCl}$  and the reduction of  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$  [12,13] indicate that an electron transfer between these species would be endothermic by ca. 100–150 kJ/mol in THF solution (and hence even more so in toluene), which would let an outer-sphere path appear rather unlikely.

An inner-sphere halogen transfer, on the other hand, would require the intermediate formation of a Cl-bridged binuclear species of the type  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}-(\text{Cl})\text{TiCp}_2$ . Such a species might conceivably be accessible directly from  $(\text{Cp}_2\text{TiCl})_2$  and CO without prior formation of  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$  and  $\text{Cp}_2\text{TiCl}$ . Our spectral data indicate, however, that the species absorbing at  $2068 \text{ cm}^{-1}$  has the composition  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$  rather than  $(\text{Cp}_2\text{TiCl})_2(\text{CO})$ ; a Cl-bridged, binuclear carbonyl species would thus probably be a collision complex arising from  $\text{Cp}_2\text{Ti}(\text{CO})\text{Cl}$  and  $\text{Cp}_2\text{TiCl}$ .

In any case, microreversibility requires that in the comproportionation of  $\text{Cp}_2\text{TiCl}_2$  and  $\text{Cp}_2\text{Ti}(\text{CO})_2$ , the reverse of reaction 5, i.e. a bimolecular reaction of  $\text{Cp}_2\text{Ti}(\text{CO})$  with  $\text{Cp}_2\text{TiCl}_2$  must be rate-determining, with  $\text{Cp}_2\text{Ti}(\text{CO})$  arising from  $\text{Cp}_2\text{Ti}(\text{CO})_2$  by reversible loss of CO. In a recent study of ligand substitution reactions of  $\text{Cp}_2\text{Ti}(\text{CO})_2$ , Basolo and coworkers [14] have shown that many of these substitution reactions do, indeed, proceed via a reversible predissociation of CO from  $\text{Cp}_2\text{Ti}(\text{CO})_2$ . It is entirely reasonable, therefore, that a coordinatively unsaturated complex  $\text{Cp}_2\text{Ti}(\text{CO})$  is an essential intermediate also in redox reactions of  $\text{Cp}_2\text{Ti}(\text{CO})_2$  such as that with  $\text{Cp}_2\text{TiCl}_2$  considered here, especially if it proceeds via an inner-sphere halogen-ligand transfer route.

Essential steps of the interconversion between  $(\text{Cp}_2\text{TiCl})_2$ ,  $\text{Cp}_2\text{Ti}(\text{CO})_2$  and  $\text{Cp}_2\text{TiCl}_2$ , as derived from our study, are summarised in Scheme 1.

### Acknowledgements

Financial support of this work by Deutsche Forschungsgemeinschaft (grant Br 510 and a guest professorship for C.D.S.), by Fonds der Chemischen Industrie and

by funds of the University of Konstanz is gratefully acknowledged. We thank Dr. H. Schwemlein for help with the preparation of  $(Cp_2TiCl)_2$ . Thanks are due to Wissenschaftliche Werkstätten of the University of Konstanz for expert collaboration in the construction of the pressure-resistant IR cell.

## References

- 1 C. Floriani and G. Fachinetti, *J. Chem. Soc., Dalton Trans.*, (1973) 1954.
- 2 B. Demerseman, G. Bouquet and M. Bigorgne, *J. Organomet. Chem.*, 101 (1975) C24.
- 3 L.P. Battaglia, M. Nardelli, C. Pelizzi, G. Predieri and G.P. Chiusoli, *J. Organomet. Chem.*, 259 (1983) 301.
- 4 M.L.H. Green and C.R. Lucas, *J. Chem. Soc., Dalton Trans.*, (1972) 1000.
- 5 E.U. van Raaij, Dissertation, University of Konstanz, 1987.
- 6 H. Weitkamp and R. Barth, *Einführung in die quantitative Infrarot-Spektrophotometrie*, Stuttgart 1976, p. 6 ff.
- 7 H.L. Clever and R. Battino, in M.R.J. Dack (Ed.), *Techniques in Chemistry*, Vol 8; New York, 1975, Part 1, Chapter VII.
- 8 I.R. Krichevsky and J.S. Kasarnovsky, *J. Amer. Chem. Soc.*, 57 (1935) 2168.
- 9 W.E. Deming and L.E. Shupe, *Phys. Rev.*, 38 (1931) 2245.
- 10 E.J.M. De Boer, L.C. Ten Cate, A.G.J. Staring and J.H. Teuben, *J. Organomet. Chem.*, 181 (1979) 61.
- 11 P.E. Matkovskii, L.I. Chernaya, F.S. D'yachkovskii and N.M. Chirkov, *Zh. Obsh. Khim.*, 44 (1974) 542.
- 12 N. El Murr and A. Chaloyard, *J. Organomet. Chem.*, 231 (1982) 1.
- 13 H. Schwemlein, W. Tritschler, H. Kiesele and H.H. Brintzinger, *J. Organomet. Chem.*, 293 (1985) 353.
- 14 G.T. Palmer, F. Basolo, L.B. Kool and M.D. Rausch, *J. Amer. Chem. Soc.*, 108 (1986) 4417.