

# Exponential stability for wave equations with non-dissipative damping<sup>☆</sup>

Jaime E. Muñoz Rivera<sup>a,b</sup>, Reinhard Racke<sup>c,\*</sup>

<sup>a</sup> *Department of Research and Development, National Laboratory for Scientific Computation, Rua Getúlio Vargas 333, Quitandinha, CEP 25651-070 Petrópolis, RJ, Brazil*

<sup>b</sup> *UF RJ, Rio de Janeiro, Brazil*

<sup>c</sup> *Department of Mathematics and Statistics, University of Konstanz, 78457 Konstanz, Germany*

## Abstract

We consider the nonlinear wave equation  $u_{tt} - \sigma(u_x)_x + a(x)u_t = 0$  in a bounded interval  $(0, L) \subset \mathbb{R}^1$ . The function  $a$  is allowed to change sign, but has to satisfy  $\bar{a} = \frac{1}{L} \int_0^L a(x) dx > 0$ . For this non-dissipative situation we prove the exponential stability of the corresponding linearized system for: (I) possibly large  $\|a\|_{L^\infty}$  with small  $\|a(\cdot) - \bar{a}\|_{L^2}$ , and (II) a class of pairs  $(a, L)$  with possibly negative moment  $\int_0^L a(x) \sin^2(\pi x/L) dx$ . Estimates for the decay rate are also given in terms of  $\bar{a}$ . Moreover, we show the global existence of smooth, small solutions to the corresponding nonlinear system if, additionally, the negative part of  $a$  is small enough.

MSC: 35L70; 35B40

*Keywords:* Indefinite damping; Wave equation; Exponential stability

## 1. Introduction

We consider the following nonlinear wave equation:

$$u_{tt} - \sigma(u_x)_x + a(x)u_t = 0 \tag{1.1}$$

for a function  $u = u(t, x)$ ,  $t \geq 0$ ,  $x \in (0, L) \subset \mathbb{R}^1$ ,  $L > 0$  fixed, with initial conditions

$$u(t = 0) = u_0, u_t(t = 0) = u_1 \tag{1.2}$$

and Dirichlet type boundary conditions

$$u(\cdot, 0) = u(\cdot, L) = 0. \tag{1.3}$$

<sup>☆</sup> Supported by a CNPq-DLR grant.

\* Corresponding author.

*E-mail addresses:* rivera@lncc.br (J.E. Muñoz Rivera), reinhard.racke@uni-konstanz.de (R. Racke).

We assume that  $a \in L^\infty((0, L))$  for the part on the exponential stability of the associated semigroup, and  $a \in C^3([0, L])$  for the discussion of the nonlinear system, as well as

$$\bar{a} := \frac{1}{L} \int_0^L a(x)dx > 0, \tag{1.4}$$

and, in particular,  $a$  may change sign in  $[0, L]$  or be zero in open subsets. The nonlinear function  $\sigma$  is assumed to satisfy

$$\sigma \in C^3(\mathbb{R}), \quad d_0 := \sigma'(0) > 0, \quad \text{and} \quad \sigma''(0) = 0. \tag{1.5}$$

**Remark.** This is, for instance, satisfied for  $\sigma$  corresponding to a vibrating string,

$$\sigma(y) = \frac{y}{\sqrt{1 + y^2}}.$$

Rewriting (1.1) as

$$u_{tt} - d_0 u_{xx} + au_t = b(u_x)u_{xx} \tag{1.6}$$

with

$$b(u_x) := \sigma'(u_x) - d_0 = \sigma'(u_x) - \sigma'(0) \tag{1.7}$$

the associated linearized system is

$$u_{tt} - d_0 u_{xx} + au_t = 0 \tag{1.8}$$

together with the initial conditions (1.2) and the boundary conditions (1.3). Since  $a$  may change sign we have a non-dissipative system still regarding  $au_t$  as a non-local but indefinite damping. There are many papers on solutions to (1.1) and on decay rates for (1.1) or (1.8) if  $a \geq 0$ , i.e. if  $a$  does not change sign; see for example the papers of Cox and Overton [3], Cox and Zuazua [5], Kawashima et al. [9], Nakao [13,14], da Silva Ferreira [19] or Zuazua [20] and the references therein. If  $a(x) \geq a_0 > 0$  is strictly positive, the exponential decay of solutions to (1.8) and also to (1.1), for small data, easily follows.

The non-dissipative case with indefinite  $a$  seems to have been posed first by Chen et al. [2] where it was conjectured that the energy

$$E_0(t) = \int_0^L (u_t^2 + u_x^2)(t, x)dx \tag{1.9}$$

decays exponentially if

$$\exists \gamma > 0 \quad \forall n = 1, 2, \dots : \int_0^L a(x) \sin^2(n\pi x/L)dx \geq \gamma \tag{1.10}$$

holds. Later Freitas [6] found that (1.10) is not sufficient to guarantee exponential stability when  $\|a\|_{L^\infty}$  is large. Replacing  $a$  by  $\varepsilon a$ , Freitas and Zuazua [7] proved that when  $a$  is of bounded variation and (1.10) holds, then there is  $\varepsilon^* = \varepsilon^*(a)$  such that for all  $\varepsilon \in (0, \varepsilon^*)$  the energy does indeed decay exponentially. This result was extended to a differential equation of the type

$$u_{tt} - u_{xx} + \varepsilon a(x)u_t + b(x)u = 0$$

by Benaddi and Rao [1]. Liu et al. [10] gave an abstract treatment of these results under certain conditions on the abstract damping operator. An extension to higher space dimensions was presented by Liu et al. [11].

Here we show that solutions to the linearized system (1.8), (1.2) and (1.3) decay exponentially for

(I) possibly large  $\|a\|_{L^\infty}$  with small  $\|a(\cdot) - \bar{a}\|_{L^2}$ , and

(II) a class of pairs  $(a, L)$  with possibly negative moment  $\int_0^L a(x) \sin^2(\pi x/L) dx$ .

In part (I) we improve the previous works that had the stronger assumption of smallness of  $\|a\|_{L^\infty}$ ; now we may admit large values of  $|a|$  as long as  $\|a(\cdot) - \bar{a}\|_{L^2}$  is small enough; cf. the examples  $(a_\mu)_\mu$  in Section 2 with  $\|a_\mu\|_{L^\infty} = 1$  and  $\|a_\mu(\cdot) - \bar{a}_\mu\|_{L^2} \rightarrow 0$  as  $\mu \rightarrow 0$ .

Part (II) is not a contradiction to a result of Freitas in [6], saying that if (1.10) is not valid, then the solution is not exponentially decaying for sufficiently small  $\|a\|_{L^\infty}$ , because in our examples of admissible pairs  $(a, L)$ , leading to exponential decay,  $a$  (resp.  $\|a\|_{L^\infty}$ ), and  $L$  are not independent.

We remark that the results are not “perturbation results for small  $a$ ”, since for  $a = 0$  we do not have any decay.

Estimates for the decay rate are also given in terms of  $\bar{a}$ . Moreover, we show the global existence of smooth, small solutions to the corresponding nonlinear system if, additionally, the negative part of  $a$  is small enough. More precisely: If  $\alpha_0$  denotes the decay rate for the linear system,

$$\int_0^L (u_t^2 + u_x^2)(t, x) \leq c_1 e^{-2\alpha_0 t}, \quad c_1 > 0,$$

then  $a^-$  has to satisfy in particular

$$\|a^-\|_{L^\infty} < \alpha_0, \tag{1.11}$$

see Section 3.

The paper is organized as follows. In Section 2 we shall prove the exponential stability for the linearized system in either of the situations (I) or (II) above. This is the crucial part, and the method will be the spectral one characterizing exponentially stable semigroups in terms of the spectrum of the associated generator of the semigroup. It is possible to give an explicit lower bound on the decay rate which, in turn, is necessary to make (1.11) a reasonable condition in the nonlinear case.

In Section 3 the global existence of small solutions to the nonlinear system is investigated. Using the result from Section 2 and perturbation arguments, the condition (1.11) is shown to be sufficient to guarantee the global existence and also the exponential stability of the nonlinear system.

Summarizing the contributions of our paper, we present results on exponential stability for the wave equation when the function  $a$  may change sign under conditions that extend the existing results to cases with indefinite damping  $a$  with possibly large  $L^\infty$ -norm, and give examples of pairs  $(a, L)$  for which exponential stability holds but the moment  $\int_0^L a(x) \sin^2(\pi x/L) dx$  is negative. We also present an explicit description of the decay rate and of the type of the associated semigroup, and also a discussion of a corresponding nonlinear problem with global existence and stability. Finally, our approach can be applied to other one-dimensional models.

We use standard notation, e.g. for Sobolev spaces.

## 2. Exponential stability for the linearized system

We first consider the linearized system. Without loss of generality we assume  $d_0 = \sigma'(0) = 1$ .

$$u_{tt} - u_{xx} + a(x)u_t = 0 \quad \text{in } (0, \infty) \times (0, L) \tag{2.1}$$

$$u(0, \cdot) = u_0, \quad u_t(0, \cdot) = u_1 \quad \text{in } (0, L) \tag{2.2}$$

$$u(\cdot, 0) = u(\cdot, L) = 0 \quad \text{in } (0, \infty). \tag{2.3}$$

We assume that  $a \in L^\infty((0, L))$  and satisfies (1.4). The aim is to prove that the energy given in (1.9) decays to zero exponentially as time  $t$  tends to infinity in either

- Case (I):  $\|a(\cdot) - \bar{a}\|_{L^2}$  is sufficiently small – but  $\|a\|_{L^\infty}$  may be large – or
- Case (II):  $\|a\|_{L^\infty}$  is small,  $(a, L)$  satisfy certain relations — but the moment  $\int_0^L a(x) \sin^2(\pi x/L) dx$  may be negative.

We introduce the variables

$$p := u_t - u_x, \quad q := u_t + u_x$$

such that

$$p_t + p_x = -a(x)u_t, \quad q_t - q_x = -a(x)u_t. \tag{2.4}$$

Let

$$U := \begin{pmatrix} p \\ q \end{pmatrix}, \quad K := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad B := \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix},$$

and let  $\mathcal{A}$  denote the operator given by

$$\mathcal{A}U := -KU_x - \frac{a}{2}BU$$

with domain

$$D(\mathcal{A}) := \left\{ \begin{pmatrix} p \\ q \end{pmatrix} \in H^1((0, L)) \times H^1((0, L)) \mid p(0) + q(0) = p(L) + q(L) = 0, \int_0^L p(s) - q(s)ds = 0 \right\}$$

in the Hilbert space

$$\mathcal{H} := \left\{ \begin{pmatrix} p \\ q \end{pmatrix} \in L^2((0, L)) \times L^2((0, L)) \mid \int_0^L p(s) - q(s)ds = 0 \right\}$$

with the  $L^2((0, L))$  inner product.  $D(\mathcal{A})$  is dense in  $\mathcal{H}$  and  $\mathcal{A}$  is the infinitesimal generator of a  $C_0$ -semigroup  $\{e^{\mathcal{A}t}\}_{t \geq 0}$ . We can rewrite (2.1) as

$$U_t = \mathcal{A}U, \quad U(0) = U_0, \quad U \in D(\mathcal{A}). \tag{2.5}$$

In a standard way we obtain

**Lemma 2.1.**  $\mathcal{A}^{-1}$  is compact.

Lemma 2.1 implies that the spectrum  $\sigma(\mathcal{A})$  of  $\mathcal{A}$  consists of eigenvalues  $(\lambda_n)_n$  only, without any finite accumulation point.

First we consider Case (I), where finally  $\|a(\cdot) - \bar{a}\|_{L^2}$  will be chosen sufficiently small, and we will show below, by using fixed point arguments, that for  $\varepsilon > 0$

$$\Gamma_\varepsilon^I := \left\{ \varepsilon + \alpha + i\beta; \alpha > \Re \left( -\frac{\bar{a}}{2} + \sqrt{\left(\frac{\bar{a}}{2}\right)^2 - \frac{\pi^2}{L^2}} \right) \text{ and } \beta \in \mathbb{R} \right\} \subset \varrho(\mathcal{A}),$$

and also that

$$\sup_{\lambda \in \Gamma_\varepsilon^I} \|(\lambda I - \mathcal{A})^{-1}\| < \infty.$$

This will imply, see e.g. the results by Prüss [17] or [12, Thm 1.3.1], that the corresponding semigroup decays exponentially.

We shall regard  $\mathcal{A}$  as a perturbation of  $\mathcal{A}_0^I$  defined by

$$\mathcal{A}_0^I U := -KU_x - \frac{\bar{a}}{2}BU$$

on

$$D(\mathcal{A}_0^I) := D(\mathcal{A}).$$

**Lemma 2.2.** Let  $\sigma(\mathcal{A}_0^I)$  denote the spectrum of  $\mathcal{A}_0^I$ . Then we have that

$$\sigma(\mathcal{A}_0^I) = \left\{ -\frac{1}{2}\bar{a} \pm \sqrt{\left(\frac{1}{2}\bar{a}\right)^2 - \frac{k^2\pi^2}{L^2}} \mid k \in \mathbb{N} \right\}$$

**Proof.** As for  $\mathcal{A}$ , one can show that  $\mathcal{A}_0^I$  has a compact inverse. We now consider the equation

$$\lambda U - \mathcal{A}_0^I U = F \tag{2.6}$$

where  $F \in \mathcal{H}$ , which is equivalent to finding a function  $U$  such that

$$U_x + MU = KF$$

where  $M := K(\lambda I + \frac{\bar{a}}{2}B)$ . The solution to this equation is given by

$$U(x) = e^{-Mx}U_0 + \int_0^x e^{-M(x-s)}KF(s) ds. \tag{2.7}$$

This is the solution of an initial value problem. Finding the set of  $\lambda$  belonging to the resolvent set is equivalent to the problem of finding  $\lambda$  such that  $U$  satisfies the boundary conditions of the problem and can be estimated appropriately by  $F$ . Defining

$$U = \begin{pmatrix} p \\ q \end{pmatrix}, \quad U_0 = \begin{pmatrix} p_0 \\ q_0 \end{pmatrix}$$

we have to find  $U \in D(\mathcal{A}_0^I)$ . To get this we have to satisfy first  $p_0 + q_0 = 0$  which implies

$$p_0 = -q_0.$$

Now the problem reduces to finding  $p_0$  for which we have

$$p(L) + q(L) = 0. \tag{2.8}$$

Let us define

$$E(x, s) := e^{-M(x-s)} =: \begin{pmatrix} e_{11}(x, s) & e_{12}(x, s) \\ e_{21}(x, s) & e_{22}(x, s) \end{pmatrix}.$$

Note that

$$\begin{pmatrix} p \\ q \end{pmatrix} = E(x, 0) \begin{pmatrix} p_0 \\ -p_0 \end{pmatrix} + \int_0^x E(x, s) \begin{pmatrix} f(s) \\ -g(s) \end{pmatrix} ds. \tag{2.9}$$

Using the expression above to verify relation (2.8) we conclude that  $p_0$  should satisfy

$$\begin{aligned} & -\{e_{11}(L, 0) - e_{12}(L, 0) + e_{21}(L, 0) - e_{22}(L, 0)\} p_0 \\ & = \int_0^L \{e_{11}(L, s) + e_{21}(L, s)\} f(s) - \{e_{12}(L, s) + e_{22}(L, s)\} g(s) ds. \end{aligned} \tag{2.10}$$

It is not difficult to see that the above problem has a solution if and only if

$$e_{11}(L, 0) - e_{12}(L, 0) + e_{21}(L, 0) - e_{22}(L, 0) \neq 0 \tag{2.11}$$

and that therefore  $\lambda \in \rho(\mathcal{A}_0^I)$  holds if and only if condition (2.11) holds. To characterize the spectrum precisely, we need to calculate the matrix  $E(x, s)$  explicitly. To do this we note that

$$M = \begin{pmatrix} \alpha & \beta \\ -\beta & -\alpha \end{pmatrix} \quad \text{where } \alpha := \lambda + \frac{1}{2}\bar{a}, \beta := \frac{1}{2}\bar{a}. \tag{2.12}$$

To get the explicit representation of the exponential matrix  $E$  we will use the eigenvector representation. Therefore our next step is to calculate the eigenvalues and eigenvectors. Since

$$\det \begin{pmatrix} \mu - \alpha & -\beta \\ +\beta & \mu + \alpha \end{pmatrix} = \mu^2 - (\alpha^2 - \beta^2)$$

it follows that the eigenvalues are given by  $\mu = \pm\sqrt{\alpha^2 - \beta^2}$ . Let us define  $\mu_0 := \sqrt{\alpha^2 - \beta^2}$  with non-negative real part. Then we have that the eigenvectors are given by

$$w_1 = \begin{pmatrix} \mu_0 + \alpha \\ -\beta \end{pmatrix}, \quad w_2 = \begin{pmatrix} -\mu_0 + \alpha \\ -\beta \end{pmatrix}.$$

Observing  $0 \in \rho(\mathcal{A}_0^I)$  we may assume  $\lambda \neq 0$ . Letting

$$D := \begin{pmatrix} \mu_0 + \alpha & -\mu_0 + \alpha \\ -\beta & -\beta \end{pmatrix} \Rightarrow D^{-1} = -\frac{1}{2\beta\mu_0} \begin{pmatrix} -\beta & \mu_0 - \alpha \\ \beta & \mu_0 + \alpha \end{pmatrix}$$

we have the representation

$$E(x, 0) = D \begin{pmatrix} e^{-\mu_0 x} & 0 \\ 0 & e^{\mu_0 x} \end{pmatrix} D^{-1}$$

which implies

$$E(x, 0) = \begin{pmatrix} \cosh(\mu_0 x) - \frac{\alpha}{\mu_0} \sinh(\mu_0 x) & -\frac{\beta}{\mu_0} \sinh(\mu_0 x) \\ \frac{\beta}{\mu_0} \sinh(\mu_0 x) & \cosh(\mu_0 x) + \frac{\alpha}{\mu_0} \sinh(\mu_0 x) \end{pmatrix}. \tag{2.13}$$

The condition (2.11) now turns into

$$\frac{\lambda}{\mu_0} \sinh(\mu_0 L) = 0 \Rightarrow e^{2\mu_0 L} = 1.$$

Therefore we conclude that  $\mu_0 L = k\pi i$ , for integers  $k$ , or

$$\mu_0^2 L^2 = -k^2 \pi^2.$$

Recalling the definition of  $\mu_0$  we get and using (2.12) we get

$$\lambda^2 L^2 + L^2 \lambda \bar{a} + k^2 \pi^2 = 0 \Rightarrow \lambda = -\frac{1}{2} \bar{a} \pm \sqrt{\left(\frac{1}{2} \bar{a}\right)^2 - \frac{k^2 \pi^2}{L^2}}.$$

Finally, note that if  $k = 0$  then we will have  $\mu_0 = 0$  and therefore  $\lambda = 0 \in \varrho(\mathcal{A}_0^I)$ .  $\square$

**Lemma 2.3.** Let  $\lambda = \gamma + i\eta$  with  $\gamma > -\frac{\bar{a}}{2}$ , and let

$$\mu_0 = A(\eta) + iB(\eta)$$

define  $A$  and  $B$ . Then we have

$$A^2 + B^2 = \sqrt{\eta^4 + [2\gamma^2 + 2\gamma\bar{a} + \bar{a}^2]\eta^2 + (\gamma^2 + \gamma\bar{a})^2}. \tag{2.14}$$

$$\lim_{\eta \rightarrow \infty} \left| \frac{\alpha}{\mu_0} \right| = 1, \quad \lim_{\eta \rightarrow \infty} \left| \frac{\beta}{\mu_0} \right| = 0, \quad \lim_{\eta \rightarrow \infty} \left| \frac{\lambda}{\mu_0} \right| = 1, \quad \lim_{\eta \rightarrow \infty} A = \gamma + \frac{\bar{a}}{2}. \tag{2.15}$$

$$\lim_{(\gamma, \eta) \rightarrow (0, \infty)} \frac{|\sinh(\mu_0 x)|}{|\mu_0|} = x, \quad \limsup_{\eta \rightarrow \infty} |\sinh(\mu_0 x)| \leq \sqrt{1 + \sinh^2\left(\gamma + \frac{\bar{a}}{2}\right)} L. \tag{2.16}$$

**Proof.** Recalling that

$$\mu_0 = \sqrt{\alpha^2 - \beta^2} = \sqrt{\lambda^2 + \lambda\bar{a}}$$

we have

$$\mu_0 = \sqrt{\alpha^2 - \beta^2} = \sqrt{\gamma^2 + \bar{a}\gamma + i\eta(\bar{a} + 2\gamma) - \eta^2} = A + iB.$$

Squaring the above expression we get

$$\gamma^2 + \bar{a}\gamma - \eta^2 = A^2 - B^2$$

$$\eta(\bar{a} + 2\gamma) = 2AB.$$

Solving the equation for  $A$  we conclude that

$$A^2 = \frac{\gamma^2 + \bar{a}\gamma}{2} + \frac{-\eta^2 + \sqrt{\eta^4 + [2\gamma^2 + 2\gamma\bar{a} + \bar{a}^2]\eta^2 + (\gamma^2 + \gamma\bar{a})^2}}{2}$$

$$B^2 = -\frac{\gamma^2 + \bar{a}\gamma}{2} + \frac{\eta^2 + \sqrt{\eta^4 + [2\gamma^2 + 2\gamma\bar{a} + \bar{a}^2]\eta^2 + (\gamma^2 + \gamma\bar{a})^2}}{2}.$$

Summing up the above identities we get (2.14). Note that

$$\begin{aligned} \frac{\sqrt{\eta^4 + [2\gamma^2 + 2\gamma\bar{a} + \bar{a}^2]\eta^2 + (\gamma^2 + \gamma\bar{a})^2}}{2} &\leq \frac{\sqrt{(\eta^2 + \gamma^2 + \gamma\bar{a} + \frac{\bar{a}^2}{2})^2 - \bar{a}^2(\gamma + \frac{\bar{a}}{2})^2}}{2} \\ &= \frac{\eta^2 + \gamma^2 + \gamma\bar{a} + \frac{\bar{a}^2}{2}}{2}. \end{aligned}$$

On the other hand we have

$$\begin{aligned} \frac{\sqrt{\eta^4 + [2\gamma^2 + 2\gamma\bar{a} + \bar{a}^2]\eta^2 + (\gamma^2 + \gamma\bar{a})^2}}{2} &\geq \frac{\sqrt{\eta^4 + [2\gamma^2 + 2\gamma\bar{a}]\eta^2 + (\gamma^2 + \gamma\bar{a})^2}}{2} \\ &= \frac{\eta^2 + \gamma^2 + \gamma\bar{a}}{2}. \end{aligned}$$

Therefore we obtain

$$\gamma^2 + \bar{a}\gamma \leq A(\eta)^2 \leq \left(\gamma + \frac{\bar{a}}{2}\right)^2.$$

Similarly we have

$$\eta^2 \leq B(\eta)^2 \leq \eta^2 + \frac{\bar{a}^2}{4}.$$

Thus we conclude

$$\lim_{\eta \rightarrow \infty} \left| \frac{\alpha}{\mu_0} \right|^2 = \lim_{\eta \rightarrow \infty} \frac{(\gamma + \frac{1}{2}\bar{a})^2 + \eta^2}{A(\eta)^2 + B(\eta)^2} = 1$$

proving (2.15). Finally,

$$\begin{aligned} \sinh(\mu_0 x) &= \frac{1}{2} \{e^{Ax}(\cos(Bx) + i \sin(Bx)) - e^{-Ax}(\cos(Bx) - i \sin(Bx))\} \\ &= \cos(Bx) \sinh(Ax) + i \sin(Bx) \cosh(Ax). \end{aligned}$$

This implies

$$\begin{aligned} |\sinh(\mu_0 x)| &= \sqrt{\cos^2(Bx) \sinh^2(Ax) + \sin^2(Bx) \cosh^2(Ax)} \\ &= \sqrt{\sinh^2(Ax) + \sin^2(Bx)} \end{aligned}$$

and similarly we have

$$|\cosh(\mu_0 x)| = \sqrt{\sinh^2(Ax) + \cos^2(Bx)}.$$

yielding (2.16). □

**Lemma 2.4.** Let  $F = (f_1, f_2)$  and  $\lambda = \gamma + i\eta$  as above, and let

$$\begin{aligned} I(\lambda, F) &:= \frac{1}{2} \frac{\int_0^L (a - \bar{a})(\cosh((L - s)\mu_0) + \frac{\lambda + \bar{a}}{\mu_0} \sinh(\mu_0(L - s))) f_1 \, ds}{-\frac{\lambda}{\mu_0} \sinh(\mu_0 L)} \\ &\quad - \frac{1}{2} \frac{\int_0^L (a - \bar{a})(\cosh(\mu_0(L - s)) + \frac{\lambda}{\mu_0} \sinh(\mu_0(L - s))) f_2 \, dx}{-\frac{\lambda}{\mu_0} \sinh(\mu_0 L)}. \end{aligned}$$

Then we have

$$\limsup_{(\gamma, \eta) \rightarrow (0, \infty)} |I(\lambda, F)| \leq \|a - \bar{a}\|_{L^2} \|F\|_{L^2} \left( \frac{1 + \bar{a}L}{\bar{a}L} \right)$$

$$\limsup_{\eta \rightarrow \infty} |I(\lambda, F)| \leq \|a - \bar{a}\|_{L^2} \|F\|_{L^2} \left( \frac{1 + \sinh((\gamma + \frac{\bar{a}}{2})L)}{\sinh((\gamma + \frac{\bar{a}}{2})L)} \right)$$

$$\limsup_{(\gamma, \eta) \rightarrow (0, \infty)} |E(x, s)| \leq 1 + \frac{\bar{a}L}{2}$$

$$\limsup_{\eta \rightarrow \infty} |E(x, s)| \leq 2\sqrt{1 + \sinh^2\left(\left(\gamma + \frac{\bar{a}}{2}\right)L\right)}.$$

**Proof.** We have

$$|I(\lambda, F)| \leq \frac{1}{2} \frac{\|a - \bar{a}\|_{L^2} (2\|\cosh(\mu_0 x)\|_{L^\infty} + \frac{|\lambda + \bar{a}| + |\lambda|}{|\mu_0|} \|\sinh(\mu_0 x)\|_{L^\infty}) \|F\|_{L^2}}{|\frac{\lambda}{\mu_0}| |\sinh(\mu_0 L)|}$$

where  $\|\cdot\|_\infty$  denotes the sup-norm with respect to  $x$ . Our conclusion now follows from Lemma 2.3 using

$$|\sinh(\mu_0 L)| \geq |\sinh(A(\eta)L)| \rightarrow \sinh\left(\left(\gamma + \frac{\bar{a}}{2}\right)L\right) \text{ as } \eta \rightarrow \infty. \quad \square$$

**Corollary 2.5.** *There exists a positive constant  $C_0$ , depending essentially only on  $|\gamma + \frac{\bar{a}}{2}|$ , such that for any  $\eta \in \mathbb{R}$  and any  $(x, s)$  we have*

$$|I(\lambda, F)| \leq C_0 \|a - \bar{a}\|_{L^2} \|F\|_{L^2},$$

$$|E(x, s)| \leq C_0.$$

**Remark.** For the interval  $(0, 1)$  and  $0 < \mu < \frac{1}{2}$ , let the function  $a_\mu : [0, 1] \rightarrow \mathbb{R}$  be given by

$$a_\mu(x) := \begin{cases} -1 & 0 \leq x < \mu \\ 1 & \mu \leq x < 1. \end{cases}$$

Then

$$\bar{a}_\mu = 1 - 2\mu \quad \text{and} \quad \|a_\mu - \bar{a}_\mu\|_{L^2} = 2\sqrt{(1 - \mu)\mu} < 1,$$

and hence, as  $\mu \rightarrow 0$ ,

$$\|a_\mu\|_{L^\infty} = 1 \quad \text{and} \quad \|a_\mu - \bar{a}_\mu\|_{L^2} \rightarrow 0.$$

Of course, from this one can easily derive examples in  $C^\infty$ .

**Lemma 2.6.** *There exists  $\tau > 0$  such that when  $\|a - \bar{a}\|_{L^2} < \tau$ , we have*

- (i)  $\Gamma_\varepsilon^I \subset \varrho(\mathcal{A})$ ,
- (ii)  $\sup_{\lambda \in \Gamma_\varepsilon^I} \|(\lambda - \mathcal{A})^{-1}\| < \infty$ .

**Proof.** It suffices to show that for sufficiently small  $\tau > 0$  and for  $\lambda \in \Gamma_\varepsilon^I$  the equation  $(\lambda - \mathcal{A})U = F$  is solvable for any  $F \in \mathcal{H}$ , and  $\|U\|_{L^2} \leq C\|F\|_{L^2}$  with a constant  $C$  at most depending on  $\varepsilon$  and  $\tau$ . We shall use a fixed point argument to prove this. Now let  $F = (f, g)' \in \mathcal{H}$  be given as well as  $\lambda \in \Gamma_\varepsilon^I$ . Let

$$\Phi : \tilde{\mathcal{H}} \rightarrow \tilde{\mathcal{H}}, \quad V \mapsto U = \Phi V, \quad \tilde{\mathcal{H}} := L^2((0, L)) \times L^2((0, L)),$$

be defined as the solution  $U = (U_1, U_2)'$  to

$$(\lambda - \mathcal{A}_0^I)U = F - \frac{a - \bar{a}}{2}BV$$

which is well defined since  $\lambda \in \varrho(\mathcal{A}_0^I)$ . Using the explicit representation of  $U$  we have

$$U(x) = (\Phi V)(x) = E_0(x)U_0 + \int_0^x E_0(x-s) \left( KF(s) - \frac{a(s) - \bar{a}}{2}KBV(s) \right) ds \tag{2.17}$$



where  $U_0 = (p_0, -p_0)$  with  $p_0 = I(\lambda, V)$ , cf. (2.10),

$$\begin{aligned}
 -p_0 = & \left\{ \int_0^L (e_{11} + e_{21})(L - s) \left( f(s) - \frac{a(s) - \bar{a}}{2} (V_1(s) + V_2(s)) \right) - (e_{12} + e_{22})(L - s) \right. \\
 & \left. \times \left( g(s) - \frac{a(s) - \bar{a}}{2} (V_1(s) + V_2(s)) \right) ds \right\} / (e_{11}(L) - e_{12}(L) + e_{21}(L) - e_{22}(L)), \tag{2.18}
 \end{aligned}$$

where  $V = (V_1, V_2)'$ . Let

$$U^j := \Phi V^j, \quad j = 1, 2.$$

Then

$$U^1(x) - U^2(x) = E_0(x)(U_0^1 - U_0^2) + \int_0^x E_0(x - s) \left( \frac{a(s) - \bar{a}}{2} K B(V^2(s) - V^1(s)) \right) ds.$$

This implies

$$|U^1(x) - U^2(x)| \leq |E_0(x, 0)(U_0^1 - U_0^2)| + c_1 \int_0^x |(a - \bar{a})E_0(x, s)(V^1(s) - V^2(s))| ds \tag{2.19}$$

and  $c_1$  denotes here and in the sequel a positive constant at most depending on  $\tau$  and  $\varepsilon$ . We conclude from Corollary 2.5 that

$$\|U_0^1 - U_0^2\| \leq c_1 \|a - \bar{a}\|_{L^2} \|V^1 - V^2\|_{L^2}. \tag{2.20}$$

The last two inequalities yield and using Corollary 2.5 once more we have

$$\|U^1 - U^2\|_{L^2} \leq c_1 \|a - \bar{a}\|_{L^2} \|V^1 - V^2\|_{L^2}.$$

Hence  $\Phi$  is a contraction mapping if

$$c_1 \|a - \bar{a}\|_{L^2} < 1$$

determining  $\tau$  as  $\tau = \frac{1}{c_1}$ . Let  $U \equiv (p, q)'$  be the unique fixed point. It satisfies

$$\lambda U + K U_x + \frac{a}{2} B U = F. \tag{2.21}$$

By definition we have  $U \in D(\mathcal{A}_0^I) = D(\mathcal{A})$ . Thus we conclude that

$$U \in D(\mathcal{A}) \quad \text{and} \quad (\lambda - \mathcal{A})U = F.$$

Finally, we estimate the inverse  $(\lambda - \mathcal{A})^{-1}$ . Let  $U$  still be the fixed point, and let

$$\tilde{U} := \Phi(0)$$

or, in other words,

$$(\lambda - \mathcal{A}_0^I)\tilde{U} = F.$$

Then we get

$$\|U\|_{L^2} - \|\tilde{U}\|_{L^2} \leq \|U - \tilde{U}\|_{L^2} = \|\Phi U - \Phi(0)\|_{L^2} \leq d \|U\|_{L^2}$$

where  $d < 1$  describes the contraction mapping property. It follows that

$$\|U\|_{L^2} \leq \frac{1}{1 - d} \|\tilde{U}\|_{L^2}.$$

On the other hand we obtain from (2.17) and (2.18) (cf. (2.19) and (2.20))

$$\|\tilde{U}\|_{L^2} \leq c_1 \|F\|_{L^2}.$$

Hence we have proved

$$\|(\lambda - \mathcal{A})^{-1}\| \leq c_1 \frac{1}{1 - d}$$

which proves the lemma.  $\square$

As a consequence we obtain the following theorem on the exponential decay.

**Theorem 2.7.** *There exists  $\tau > 0$  such that when  $\|a - \bar{a}\|_{L^2} < \tau$ , we have that the solution to the linearized system decays exponentially, that is*

$$\exists c_0 > 0 \exists \alpha_0 > 0 \forall t \geq 0 : E_0(t) \leq c_0 e^{-2\alpha_0 t} E_0(0).$$

In particular we can take any

$$\alpha_0 > \Re \left( -\frac{\bar{a}}{2} + \sqrt{\left(\frac{\bar{a}}{2}\right)^2 - \frac{\pi^2}{L^2}} \right),$$

e.g.  $\alpha_0 = -\frac{1}{4L} \int_0^L a(x) dx$  when  $\int_0^L a(x) dx < 2\pi$ .

**Proof.** The assertion follows from Lemma 2.6 by well known characterizations of the exponential stability of semigroups; see the results given by Prüss [17], and cf. [12, Thm 1.3.1].  $\square$

Now we consider Case (II), where finally  $\|a\|_{L^\infty}$  will have to be sufficiently small, related to  $L$ , that is,  $(a, L)$  will satisfy certain relations. Contrasting these restrictions with respect to previous results for small  $a$ , e.g. those of Freitas and Zuazua [7], we shall obtain examples where the moment  $\int_0^L a(x) \sin^2(\pi x/L) dx$  is negative.

Similarly to in Case (I) above, we wish to prove that for small  $\varepsilon_1 > 0$  and any  $\varepsilon_0 > \varepsilon_1$  we can choose  $(a, L)$  with  $a$  small enough such that for any  $\varepsilon \in [\varepsilon_1, \varepsilon_0]$

$$\Gamma_\varepsilon^{II} := \left\{ -\frac{\bar{a}}{2} + \varepsilon + i\eta \mid \eta \in \mathbb{R} \right\} \subset \varrho(\mathcal{A}) \tag{2.22}$$

and that

$$\sup_{\varepsilon \in [\varepsilon_1, \varepsilon_0], \lambda \in \Gamma_\varepsilon^{II}} \|(\lambda - \mathcal{A})^{-1}\| < \infty. \tag{2.23}$$

We choose

$$\varepsilon_1 := \frac{\bar{a}}{4}, \quad \varepsilon_0 := \max\{2\varepsilon_1, 3|a|_\infty\}, \tag{2.24}$$

where  $|a|_\infty := \|a\|_{L^\infty}$ , and we observe that  $\mathcal{A} - 3|a|_\infty$  is dissipative.

Now we shall regard  $\mathcal{A}$  as a perturbation of the following operator  $\mathcal{A}_0^{II}$ , given by

$$\mathcal{A}_0^{II} U := -K U_x - \frac{a}{2} U$$

with domain

$$D(\mathcal{A}_0^{II}) := D(\mathcal{A})$$

in the same Hilbert space  $\mathcal{H}$ . For  $U \in D(\mathcal{A})$  we have

$$\mathcal{A}U = \mathcal{A}_0^{II} U - \frac{a}{2} WU$$

with

$$W := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

$(\mathcal{A}_0^{II})^{-1}$  is compact.

**Lemma 2.8.**  $\sigma(\mathcal{A}_0^{II}) = \left\{ -\frac{\bar{a}}{L} + \frac{k\pi i}{L} \mid k \in \mathbb{Z} \right\}$ .

**Proof.** We investigate the solvability of  $(\lambda - \mathcal{A}_0^{II})U = F$  for any  $F = (f, g)' \in \mathcal{H}$ . Using ideas similar to those in the proof of Lemma 2.2 we obtain

$$\lambda \in \sigma(\mathcal{A}_0^{II}) \Leftrightarrow e^{-2\lambda L - \int_0^L a(y)dy} = 1 \Leftrightarrow \lambda = \lambda_k = -\frac{\bar{a}}{L} + \frac{k\pi i}{L}, \quad k \in \mathbb{Z}. \quad \square$$

For given  $0 < \gamma_0 < \gamma_1$  we define the set of admissible  $a$  and  $L$  as

$$\mathcal{K}(\gamma_0, \gamma_1) := \left\{ (a, L) \mid \gamma_0 \leq \int_0^L a(x)dx \leq a_\infty L \leq \gamma_1 \right\}.$$

We shall prove that, after fixing  $\gamma_0$  and  $\gamma_1$ , we can determine the maximal possible  $L^\infty$ -norm of  $a$ , and the admissible values of  $L$ , that imply exponential decay.

**Lemma 2.9.** *There are pairs  $(a, L) \in \mathcal{K}(\gamma_0, \gamma_1)$  such that*

- (i)  $\forall \varepsilon \in [\varepsilon_1, \varepsilon_0] : \Gamma_\varepsilon = \{-\frac{\bar{a}}{2} + \varepsilon + i\eta \mid \eta \in \mathbb{R}\} \subset \rho(\mathcal{A})$ .
- (ii)  $\sup_{\varepsilon \in [\varepsilon_1, \varepsilon_0], \lambda \in \Gamma_\varepsilon} \|(\lambda - \mathcal{A})^{-1}\| < \infty$ .

**Proof.** We will show, for any admitted  $\varepsilon$ , that for  $\lambda \in \Gamma_\varepsilon^{II}$  the equation  $(\lambda - \mathcal{A})U = F$  is solvable for any  $F \in \mathcal{H}$ , and the postulated estimate on the inverse holds. We shall use again a fixed point argument to prove this. Now let  $F = (f, g)' \in \mathcal{H}$  be given as well as  $\lambda \in \Gamma_\varepsilon$ . Let

$$\Phi : \tilde{\mathcal{H}} \rightarrow \tilde{\mathcal{H}}, \quad V \rightarrow U = \Phi V$$

be defined as solution  $U = (U_1, U_2)'$  to

$$(\lambda - \mathcal{A}_0^{II})U = F - \frac{a}{2}WV$$

which is well defined since  $\lambda \in \rho(\mathcal{A}_0^{II})$ . Let us consider

$$U(x) = (\Phi V)(x) = E_0(x, 0, \lambda)U_0 + \int_0^x E_0(x, s, \lambda) \left( KF(s) - \frac{a(s)}{2}KWV(s) \right) ds \tag{2.25}$$

where

$$E_0(x, s, \lambda) := \begin{pmatrix} e^{-\lambda(x-s) - \int_s^x \frac{a(y)}{2}dy} & 0 \\ 0 & e^{\lambda(x-s) + \int_s^x \frac{a(y)}{2}dy} \end{pmatrix} \equiv \begin{pmatrix} e_1(x, s, \lambda) & 0 \\ 0 & e_2(x, s, \lambda) \end{pmatrix}$$

and  $U_0 = (p_0, -p_0)$  with

$$p_0 = \frac{\int_0^L e_1(L, s, \lambda) \left( f(s) - \frac{a(s)}{2}V_2(s) \right) - e_2(L, s, \lambda) \left( g(s) - \frac{a(s)}{2}V_1(s) \right) ds}{e_2(L, 0, \lambda) - e_1(L, 0, \lambda)}. \tag{2.26}$$

Let

$$U^j := \Phi V^j, \quad j = 1, 2.$$

Then

$$U^1(x) - U^2(x) = E_0(x, 0, \lambda)(U_0^1 - U_0^2) + \int_0^x E_0(x, s, \lambda) \left( \frac{a(s)}{2}KW(V^2(s) - V^1(s)) \right) ds.$$

This implies

$$|U^1(x) - U^2(x)|^2 \leq 2e^{(a_\infty + 2\varepsilon_0)L} |U_0^1 - U_0^2|^2 + e^{(a_\infty + 2\varepsilon_0)L} \frac{a_\infty^2}{2} \left( \int_0^x |V^1(s) - V^2(s)| ds \right)^2. \tag{2.27}$$

We conclude from (2.26) that

$$|U_0^1 - U_0^2|^2 \leq \frac{4e^{(a_\infty + 2\varepsilon_0)L} a_\infty^2}{(\sinh(\varepsilon_1 L))^2} \left( \int_0^L |V^1(s) - V^2(s)| ds \right)^2. \tag{2.28}$$

The last two inequalities yield

$$\|U^1 - U^2\|_{L^2} \leq a_\infty e^{(\frac{a_\infty}{2} + \varepsilon_0)L} \left( \frac{2e^{(\frac{a_\infty}{2} + \varepsilon_0)L}}{\sinh(\varepsilon_1 L)} + 1 \right) \|V^1 - V^2\|_{L^2}.$$

Hence, observing (2.24),  $\Phi$  is a contraction mapping for  $(a, L)$  in  $\mathcal{K}$  if

$$a_\infty < \frac{\sinh(\gamma_0/4)}{(\sinh(\gamma_1/4) + e^{7/2\gamma_1})e^{7/2\gamma_1}} =: c_0(\gamma_0, \gamma_1). \tag{2.29}$$

(i) Now fixing  $\gamma_0$  and  $\gamma_1$ , the last inequality determines the maximal possible  $L^\infty$ -norm of  $a$ .

(ii) Then the condition

$$\frac{\gamma_0}{a_\infty} < L \leq \frac{\gamma_1}{a_\infty} \tag{2.30}$$

determines the possible values of  $L$ .

(iii) Now fixing  $L$ , since  $a_\infty L > \gamma_0$ , we can choose  $a$  satisfying (i) as well as

$$\int_0^L a(x)dx \geq \gamma_0. \tag{2.31}$$

Altogether we have found in (i)–(iii) pairs  $(a, L) \in \mathcal{K}$  for which  $\Phi$  is a contraction.

Let  $U \equiv (p, q)'$  be the unique fixed point. It satisfies

$$\lambda U + KU_x + \frac{a}{2}BU = F. \tag{2.32}$$

By definition we have  $U \in D(\mathcal{A}_0^{II}) = D(\mathcal{A})$ . Thus we conclude

$$U \in D(\mathcal{A}) \quad \text{and} \quad (\lambda - \mathcal{A})U = F.$$

Finally, we obtain the uniform boundedness of the inverses as in Case (I); namely let  $U$  still be the fixed point, and let  $\tilde{U} := \Phi(0)$ . Using the same arguments as in the proof of Lemma 2.6 we get

$$\|U\|_{L^2} \leq c\|F\|_{L^2}$$

where  $c$  depends at most on  $a, L, \gamma_0, \gamma_1$ . This proves assertion (ii).  $\square$

As in Case (I) we conclude the exponential stability. Let

$$\mathcal{E}(t) := \frac{1}{2}\|U(t, \cdot)\|_{L^2}^2 = \frac{1}{2}\|e^{t\mathcal{A}}U_0\|_{L^2}^2$$

be the energy associated with problem (2.5). Then

$$\begin{aligned} \mathcal{E}(t) &= \frac{1}{2}\left\| \begin{pmatrix} p \\ q \end{pmatrix} (t, \cdot) \right\|_{L^2}^2 = \frac{1}{2}\left\| \begin{pmatrix} u_t - u_x \\ u_t + u_x \end{pmatrix} (t, \cdot) \right\|_{L^2}^2 \\ &= \int_0^L (u_t^2 + u_x^2)(t, x)dx. \end{aligned}$$

**Theorem 2.10.** *For sufficiently small  $\|a\|_\infty$  and admissible  $(a, L) \in \mathcal{K}(\gamma_0, \gamma_1)$ , we have*

$$\exists c_0 > 0 \exists \alpha_0 > 0 \forall t \geq 0 : \mathcal{E}(t) \leq c_0 e^{-2\alpha_0 t} \mathcal{E}(0).$$

Using Lemma 2.9 which gives information on the spectral radius  $\omega_{\sigma(\mathcal{A})} := \sup_{\lambda \in \sigma(\mathcal{A})} \Re \lambda$ , a result of Neves et al. [16] which says that the essential type  $\omega_e(\mathcal{A})$  is given by

$$\omega_e(\mathcal{A}) = -\frac{1}{2L} \int_0^L a(y)dy \tag{2.33}$$

as well as using the general characterization (see [15])

$$\omega_0(\mathcal{A}) = \max\{\omega_e(\mathcal{A}), \omega_\sigma(\mathcal{A})\},$$

where  $\omega_0(\mathcal{A})$  denotes the type of the semigroup,

$$\omega_0(\mathcal{A}) = \lim_{t \rightarrow \infty} \frac{\ln \|e^{At}\|}{t}.$$

Using this we can establish

**Theorem 2.11.** *Under the conditions of Theorem 2.10 we have*

$$\exists c_0 > 0 \exists \alpha_0 = \alpha_0 \left( \frac{1}{L} \int_0^L a(x) dx \right) > 0 \forall t \geq 0 : \mathcal{E}(t) \leq c_0 e^{-2\alpha_0 t} \mathcal{E}(0).$$

$\alpha_0$  can be chosen as any number  $-\bar{\alpha}$  with

$$0 > \bar{\alpha} > -\frac{1}{2L} \int_0^L a(x) dx + \varepsilon_1, \quad \text{e.g. } \alpha_0 := \frac{1}{4L} \int_0^L a(x) dx.$$

We can now present an example of a function  $a : [0, 1] \rightarrow \mathbb{R}$  for which exponential stability holds, but for which (1.10) is violated since we shall have

$$\int_0^1 a(x) \sin^2(\pi x) dx < 0. \tag{2.34}$$

This will not be a contradiction to a result of Freitas in [6] saying that if (1.10) is not valid, then the solution is not exponentially decaying if one replaces  $a$  by  $\varepsilon a$  for sufficiently small  $\varepsilon > 0$ , because in our example replacing  $a$  by  $\varepsilon a$  is not allowed in general because of the admissibility criteria to be observed in the construction of  $(a, L)$  in Case (II).

Let  $0 < \delta_1 < \delta_2$ , to be fixed later, and let

$$a(x) := \begin{cases} \delta_1 & \text{for } x \in [0, 1/4] \cup (3/4, 1] \\ -\delta_2 & \text{for } x \in [1/4, 3/4]. \end{cases}$$

Then  $0 < \int_0^1 a(x) dx = (\delta_1 - \delta_2)/2$  provided  $\delta_1 > \delta_2$ . Moreover, we have

$$\int_0^1 a(x) \sin^2(\pi x) dx = \frac{1}{2\pi} ((\pi - 1)\delta_1 - (\pi + 1)\delta_2) < 0$$

if and only if

$$\delta_1 < \frac{\pi + 1}{\pi - 1} \delta_2.$$

Hence choosing  $\delta_1, \delta_2$  such that

$$\delta_2 < \delta_1 < \frac{\pi + 1}{\pi - 1} \delta_2 \tag{2.35}$$

we have a function  $a$  satisfying

$$0 < \int_0^1 a(x) dx \quad \text{and} \quad \int_0^1 a(x) \sin^2(\pi x) dx < 0.$$

Choose  $\delta_2 := \alpha \delta_1$  such that (2.35) is satisfied, e.g.  $\alpha := 1/1.92$  for the rest of the exposition. Then  $\delta_1$  is still free. Let us define the function  $\hat{a}$  as

$$\hat{a}(x) := \frac{a(x/L)}{L} \quad \text{for } x \in (0, L)$$

where  $L$  will be chosen to satisfy (2.30). Since  $a_\infty = \delta_1$  we have  $\hat{a}_\infty = \delta_1/L$ . We are free to choose  $\gamma_0$  and  $\gamma_1$ . Fix  $\gamma_1$ . The condition (2.31) can be satisfied if

$$\gamma_0 \leq (\delta_1 - \delta_2)/2 = \frac{23}{96}\delta_1,$$

i.e. we choose  $\delta_1 := \frac{96}{23}\gamma_0$ , and condition (2.29) can now be satisfied if

$$\frac{\delta_1}{L} < c_0(\gamma_0, \gamma_1) = c_{\gamma_1} \sinh(\gamma_0/4)$$

where

$$c_{\gamma_1} := \frac{1}{(\sinh(\gamma_1/4) + \exp(7\gamma_1/2)) \exp(7\gamma_1/2)}.$$

Since, for small  $\gamma_0$ ,

$$\sinh(\gamma_0/4) \geq \frac{\gamma_0}{5}$$

it is hence sufficient to require

$$L > \frac{480}{23c_{\gamma_1}}. \tag{2.36}$$

The last condition (2.30) can now be satisfied if  $\frac{96}{23}\gamma_0 \leq \gamma_1$ ; e.g. take  $\gamma_0 := \chi \frac{23}{96}\gamma_1$  with appropriate  $\chi \in (0, 1]$ . As a conclusion from the discussion of Case (II) above we get that with such an  $L$  and the given  $\hat{a}$  the solution to

$$u_{tt} - u_{xx} + \hat{a}(x)u_t = 0 \quad \text{in } (0, L)$$

plus initial and boundary conditions has exponentially decaying energy. In the way we have chosen  $\hat{a}$  and  $a$ , we have now found  $u$  solving

$$u_{tt} - u_{xx} + a(x)u_t = 0 \quad \text{in } (0, 1)$$

with exponentially decaying energy for a function  $a$  satisfying

$$0 < \int_0^1 a(x)dx \quad \text{and} \quad \int_0^1 a(x) \sin^2(\pi x)dx < 0.$$

We remark that from this example one can of course also construct examples in  $C^\infty$ .

We finish this section by giving some higher norm estimates valid in both Cases (I) and (II), i.e. whenever exponential stability is given. Differentiating the differential equation (2.1) with respect to  $t$ ,

$$(\partial_t^j u)_{tt} - (\partial_t^j u)_{xx} + a(x)(\partial_t^j u)_t = 0, \quad j \in \mathbb{N},$$

and using the fact that derivatives with respect to  $x$  can be computed from the differential equation successively, we get as a consequence of  $a \in C^0([0, L], \mathbb{R})$ , and  $a \in C^{s-2}([0, L], \mathbb{R})$  if  $s \geq 2$  the following theorem.

**Theorem 2.12.** *Under the conditions of the Theorems 2.7 and 2.10, respectively, we have*

$$\forall s \in \mathbb{N} \quad \exists C_s > 0 \quad \forall t \geq 0 : \left\| \begin{pmatrix} u_t \\ u_x \end{pmatrix} (t, \cdot) \right\|_{H^s((0,L))} \leq C_s e^{-\alpha_0 t} \left\| \begin{pmatrix} u_1 \\ u_{0,x} \end{pmatrix} \right\|_{H^s(0,L)}$$

where  $\alpha_0$  is given in Theorem 2.11, and the data are assumed to be sufficiently smooth and to satisfy the usual compatibility conditions.

- Remarks.** 1. We only used  $a \in L^\infty((0, L))$ .  
 2. Without loss of generality we studied the equation

$$u_{tt} - d_0 u_{xx} + a(x)u_t = 0, \quad x \in (0, L),$$

for  $d_0 = 1$ , because if  $d_0 > 0$  is arbitrary we may define

$$v(t, y) := u(t, \sqrt{d_0}y), \quad y \in \left(0, \frac{L}{\sqrt{d_0}}\right).$$

Then  $v$  satisfies

$$v_{tt} - v_{yy} + \tilde{a}(y)v_t = 0, \quad y \in \left(0, \frac{L}{\sqrt{d_0}}\right),$$

for which **Theorem 2.11** can be applied directly replacing  $a$  by  $\tilde{a}$  and  $L$  by  $L/\sqrt{d_0}$ . The decay rate  $\tilde{\alpha}_0 = \tilde{\alpha}_0 \left(\frac{\sqrt{d_0}}{L} \int_0^{L/\sqrt{d_0}} \tilde{a}(y)dy\right)$  turns into  $\tilde{\alpha}_0 = \alpha_0 \left(\frac{1}{L} \int_0^L a(y)dy\right)$  again since  $\frac{\sqrt{d_0}}{L} \int_0^{L/\sqrt{d_0}} \tilde{a}(y)dy = \frac{1}{L} \int_0^L a(x)dx$ .

### 3. Global existence for the nonlinear system

We now return to the nonlinear system (1.1)–(1.3) assuming again the positivity of the mean value (1.4) and also the condition (1.5) on the nonlinearity, which, for example, is satisfied in the classical model for a nonlinear string, where

$$\sigma(u_x) = \frac{u_x}{\sqrt{1 + u_x^2}}.$$

After recalling the local well-posedness it is our aim to prove a global existence result for data  $(u_0, u_1)$  being sufficiently small in  $H^4((0, L))$ , and, quasi-simultaneously, to obtain the exponential stability. The method used imitates one which is well known for nonlinear evolution equations and systems; see [18] for a presentation of the general approach for Cauchy problems ( $x \in \mathbb{R}^n$ , no boundary). Here we shall have to prove so-called high energy estimates and a weighted a priori estimate – describing the expected exponential decay – for a boundary value problem and a non-dissipative problem reflected in the possible negativity of the function  $a$ . To deal with the latter the condition (1.11) on the negative part of  $a$ , i.e.

$$\|a^-\|_{L^\infty} < \alpha_0,$$

will be used.

We assume that the conditions on  $a$  and on  $(a, L)$  are satisfied which assures the exponential stability of the linearized systems as given in **Theorem 2.12**.

Observing that the term  $a(x)u_t$  is of lower order, we can recall the following local existence theorem; see for instance [4] or [8, p. 97].

**Theorem 3.1.** *There is  $T = T(\|(u_0, u_1)\|_{H^4 \times H^3}) > 0$  such that (1.1)–(1.3) has a unique local solution*

$$u \in \bigcap_{k=0}^3 C^k([0, T]), H^{4-k}((0, L)) \cap H_0^1((0, L)) \cap C^4([0, T], L^2((0, L))).$$

**Remark.** Of course  $u_0, u_1$  have to satisfy the usual compatibility conditions.

Now we turn to the high energy estimates. For this purpose it is useful to rewrite (1.1)–(1.3) as a first-order system for

$$V := (u_t, u_x)'$$

Then  $V$  satisfies

$$V_t + \underbrace{\begin{pmatrix} a & -d_0 \partial_x \\ -\partial_x & 0 \end{pmatrix}}_{=: -A} V = \begin{pmatrix} b(u_x) \partial_x u_x \\ 0 \end{pmatrix} =: F(V, V_x),$$

$$V(t = 0) = (u_1, \partial_x u_0)' =: V_0.$$

The first formally defined operator  $A$  generates a  $C_0$ -semigroup as usual; for  $F = 0$  the solution  $V$  is given by

$$V(t) = e^{tA} V_0$$

and the (local) solution to (1.1)–(1.3) satisfies

$$V(t) = e^{tA} V_0 + \int_0^t e^{(t-r)A} F(V, V_x)(r) dr. \tag{3.1}$$

We conclude from Section 2 that

$$V := (u_t, u_x)$$

as a solution of the linear system (1.1)–(1.3) written in first-order form satisfies

$$V(t) = e^{tA} V(t = 0)$$

with a  $C_0$ -semigroup  $\{e^{tA}\}_{t \geq 0}$  satisfying

$$\|V(t)\|_{H^s} \leq c_s e^{-\alpha_0 t} \|V(t = 0)\|_{H^s} \tag{3.2}$$

for  $s = 0, 1, 2$  (cf. below). This follows from Theorem 2.7 for  $s = 0$  and is obtained for  $s = 1, 2$  by differentiating Eq. (1.8) with respect to  $t$  once and then twice, as well as using the differential equation to obtain information for derivatives in  $x$ . Let

$$a_\infty^- := \|a^-\|_{L^\infty}. \tag{3.3}$$

In the sequel we assume without loss of generality that  $u_x$  is small enough a priori, i.e. such that  $\sigma'(u_x)$  remains strictly positive (near  $u_x = 0$ ; cf. (1.5)), or in other words we can assume that there is  $\eta > 0$  such that

$$d_0 + b(u_x) \geq \frac{d_0}{2} > 0 \quad \text{if } |u_x| < \eta < 1. \tag{3.4}$$

**Lemma 3.2.** *There are constants  $c_2, c_3 > 0$ , not depending on  $V_0$  or  $T$ , such that the local solution given in Theorem 3.1 satisfies, for  $t \in [0, T]$ ,*

$$\|V(t)\|_{H^3}^2 \leq c_2 \|V_0\|_{H^3}^2 e^{2a_\infty^- t} e^{c_3 \int_0^t (\|V(r)\|_{H^2} + \|V(r)\|_{H^2}^2 + \|V(r)\|_{H^2}^3) dr}.$$

**Proof.** Multiplying

$$u_{tt} - d_0 u_{xx} + au_t = b(u_x) u_{xx} \tag{3.5}$$

by  $u_t$  in  $L^2$  we obtain ( $f \equiv \int_0^L$ )

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int u_t^2 + d_0 u_x^2 dx &= - \int au_t^2 dx + \int b(u_x) u_{xx} u_t dx \\ &\leq \int a_\infty^- u_t^2 - \int (\partial_x b(u_x)) u_x u_t dx - \int b(u_x) u_x u_{xt} dx \\ &\equiv I.1 + I.2 + I.3, \end{aligned} \tag{3.6}$$

$$\begin{aligned} |I.2| &\leq \frac{1}{2} \|b'(u_x) u_{xx}\|_{L^\infty} \int u_x^2 + u_t^2 dx \\ &\leq c \|V\|_{H^2} \int u_x^2 + u_t^2 dx \end{aligned} \tag{3.7}$$

where  $c$  will denote a constant not depending on  $V^0$  or on  $T$ .

$$\begin{aligned} I.3 &= -\frac{1}{2} \frac{d}{dt} \int b(u_x) u_x^2 dx + \frac{1}{2} \int (\partial_t b(u_x)) u_x^2 dx \\ &\equiv I.3.1 + I.3.2. \end{aligned} \tag{3.8}$$



The term  $I.3.2$  can be estimated in the same way as  $I.2$  in (3.7):

$$|I.3.2| \leq c \|V\|_{H^2} \int u_x^2. \tag{3.9}$$

The term  $I.3.1$  can be incorporated into and be dominated by the left-hand side of inequality (3.6) after an integration with respect to  $t$  later on, since

$$\int_0^t I.3.1(r) dr = -\frac{1}{2} \int b(u_x) u_x^2 dx + \frac{1}{2} \int b(u_x(t=0)) u_x^2(t=0) dx. \tag{3.10}$$

Summarizing (3.6)–(3.10) we have proved

$$\|V(t)\|_{L^2}^2 \leq c \|V_0\|_{L^2}^2 + \int_0^t (2a_\infty^- + c \|V(r)\|_{H^2}) \|V(r)\|_{L^2}^2 dr. \tag{3.11}$$

In order to get estimates for the higher order derivatives of  $V$  and  $u$  we differentiate Eq. (3.5) with respect to  $t$  to get

$$u_{ttt} - d_0 u_{txx} + a u_{tt} = b'(u_x) u_{xt} u_{xx} + b(u_x) u_{txx}. \tag{3.12}$$

Multiplying by  $u_{tt}$  in  $L^2$  we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int u_{tt}^2 + d_0 u_{tx}^2 dx &\leq \int a_\infty^- u_{tt}^2 dx + \int b(u_x) u_{txx} u_{tt} dx + \int b'(u_x) u_{xt} u_{xx} u_{tt} dx \\ &\equiv I.4 + I.5 + I.6. \end{aligned} \tag{3.13}$$

The term  $I.5$  can be treated like the term  $I.2 + I.3$  from (3.6); see (3.7)–(3.11).

$$|I.6| \leq c \|V\|_{H^2} \int u_{tt}^2 + u_{tx}^2 dx. \tag{3.14}$$

Observe that the differential equation (3.5) yields the estimate

$$|u_{xx}|^2 \leq c(|u_{tt}|^2 + |u_t|^2). \tag{3.15}$$

Thus we obtain from (3.11), (3.13) and (3.14)

$$\|V(t)\|_{H^1}^2 \leq c \|V_0\|_{H^1}^2 + \int_0^t (2a_\infty^- + c \|V(r)\|_{H^2}) \|V(r)\|_{H^1}^2 dr. \tag{3.16}$$

Differentiating the differential equation (3.12) once more with respect to  $t$  we get

$$u_{tttt} - d_0 u_{ttxx} + a u_{ttt} = b''(u_x) u_{xt}^2 u_{xx} + b'(u_x) u_{xxt} u_{xx} + 2b'(u_x) u_{xt} u_{xxt} + b(u_x) u_{ttxx}. \tag{3.17}$$

Multiplying by  $u_{ttt}$  in  $L^2$  we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int u_{ttt}^2 + d_0 u_{ttx}^2 dx &\leq \int a_\infty^- u_{ttt}^2 dx + \int b''(u_x) u_{xt}^2 u_{xx} u_{ttt} dx + \int b'(u_x) u_{xxt} u_{xx} u_{ttt} dx \\ &\quad + 2 \int b'(u_x) u_{xt} u_{xxt} u_{ttt} dx + \int b(u_x) u_{ttxx} u_{ttt} dx \\ &\equiv I.7 + I.8 + I.9 + I.10 + I.11. \end{aligned} \tag{3.18}$$

The term  $I.11$  is again dealt with like  $I.2 + I.3$  in (3.7)–(3.11).

$$|I.8| + |I.9| + |I.10| \leq c(\|V\|_{H^2}^2 + \|V\|_{H^2}) \int u_{xt}^2 + u_{ttt}^2 + u_{ttx}^2 + u_{xxt}^2 + u_{xxt}^2 dx. \tag{3.19}$$

Hence we obtain from (3.16) and (3.19) using (3.12) to estimate  $u_{ttxx}$ ,

$$\|V(t)\|_{H^2}^2 \leq c \|V_0\|_{H^2}^2 + \int_0^t (2a_\infty^- + c(\|V(r)\|_{H^2} + \|V(r)\|_{H^2}^2)) \|V(r)\|_{H^2}^2 dr. \tag{3.20}$$

The final estimate is obtained after differentiating the differential equation a last time with respect to  $t$  yielding

$$\begin{aligned}
 u_{tttt} - d_0 u_{tttx} + a u_{tttt} &= b'''(u_x) u_{xt}^3 u_{xx} + 3b''(u_x) u_{xt} u_{xxt} u_{xx} \\
 &\quad + 3b''(u_x) u_{xt}^2 u_{xxt} + b'(u_x) u_{xtt} u_{xx} + 3b'(u_x) u_{xtt} u_{xxt} \\
 &\quad + 3b'(u_x) u_{xt} u_{xxt} + b(u_x) u_{tttx} \\
 &\equiv \sum_{j=12}^{18} \theta_j.
 \end{aligned} \tag{3.21}$$

**Remark.** The derivatives of order 5 are formally not defined but the estimates aimed at will only involve derivatives of order 4. A usual approximation argument with data  $V_0 \in H^4((0, L))$  and the lower semicontinuity of the norms justifies our calculation finally for  $V_0 \in H^3((0, L))$  only.

A multiplication of (3.21) by  $u_{tttt}$  in  $L^2$  yields

$$\frac{1}{2} \frac{d}{dt} \int u_{tttt}^2 + u_{tttx}^2 dx \leq \int 2a_\infty^- u_{tttt}^2 + \sum_{j=12}^{18} \int \theta_j u_{tttt} dx. \tag{3.22}$$

The term  $\int \theta_{18} u_{tttt} dx$  can be dealt with like I.2. + I.3 in (3.7)–(3.11). The terms  $\int \theta_j u_{tttx} dx$  for  $j \neq 16, 18$  can be estimated easily as before using

$$\left| \sum_{\substack{j=12 \\ j \neq 16}}^{17} \int \theta_j u_{tttt} dx \right| \leq c(\|V\|_{H^2} + \|V\|_{H^2}^2 + \|V\|_{H^2}^3) \|V\|_{H^3}^2. \tag{3.23}$$

The only more difficult term is

$$\int \theta_{16} u_{tttt} = 3 \int b'(u_x) u_{xtt} u_{xxt} u_{tttt} dx$$

involving three factors of order at least 3. This term is estimated as follows using the Gagliardo–Nirenberg inequality.

$$\begin{aligned}
 \left| \int b'(u_x) u_{ttt} u_{txx} u_{tttt} dx \right| &\leq c \|b'(u_x) u_{ttt} u_{txx}\|_{L^2} \|u_{tttt}\|_{L^2} \\
 &\leq c \|b'(u_x) u_{ttt} u_{txx}\|_{L^2} \|V\|_{H^3}.
 \end{aligned} \tag{3.24}$$

$$\begin{aligned}
 \|b'(u_x) u_{ttt} u_{txx}\|_{L^2} &\leq \|\partial_x(b'(u_x) u_{tt}) \partial_x(u_{tx})\|_{L^2} + \|b''(u_x) u_{xx} u_{tt} u_{txx}\|_{L^2} \\
 &\leq \|\partial_x(b'(u_x) u_{tt}) \partial_x(u_{tx})\|_{L^2} + c \|V\|_{H^2}^2 \|V\|_{H^3}.
 \end{aligned} \tag{3.25}$$

Using Young’s inequality and the Gagliardo–Nirenberg inequality, and observing  $L^\infty \hookrightarrow H^1$ , we conclude that

$$\begin{aligned}
 \|\partial_x(b'(u_x) u_{tt}) \partial_x(u_{tx})\|_{L^2} &\leq \|\partial_x(b'(u_x) u_{tt})\|_{L^4} \|\partial_x(u_{tx})\|_{L^4} \\
 &\leq c \|b'(u_x) u_{tt}\|_{H^2}^{1/2} \|b'(u_x) u_{tt}\|_{L^\infty}^{1/2} \|u_{tx}\|_{H^2}^{1/2} \|u_{tx}\|_{L^\infty}^{1/2} \\
 &= c (\|b'(u_x) u_{tt}\|_{L^\infty} \|u_{tx}\|_{H^2})^{1/2} (\|b'(u_x) u_{tt}\|_{H^2} \|u_{tx}\|_{L^\infty})^{1/2} \\
 &\leq c (\|b'(u_x) u_{tt}\|_{L^\infty} \|u_{tx}\|_{H^2} + \|u_{tx}\|_{L^\infty} \|b'(u_x) u_{tt}\|_{H^2}) \\
 &\leq c \|V\|_{H^2} \|V\|_{H^3}.
 \end{aligned} \tag{3.26}$$

From (3.20) and (3.22)–(3.26) we conclude that

$$\|V(t)\|_{H^3}^2 \leq c \|V_0\|_{H^3}^2 + \int_0^t (2a_\infty^- + c(\|V\|_{H^2} + \|V\|_{H^2}^2 + \|V\|_{H^2}^3))(r) \|V(r)\|_{H^3}^2 dr$$

which yields the assertion of Lemma 3.2 using Gronwall’s inequality.  $\square$

Next we want to prove a weighted a priori estimate for  $\|V(t)\|_{H^2}$ .

**Remark.** Observe that we have not yet used the assumption (1.5) requiring  $b'(0) = \sigma''(0) = 0$ . Indeed, with this assumption it would be possible to remove the linear term  $\|V(t)\|_{H^2}^1$  in the exponential in the estimate for  $\|V(t)\|_{H^3}$

in Lemma 3.2, i.e. the estimate would read

$$\|V(t)\|_{H^3}^2 \leq C \|V_0\|_{H^3}^2 e^{2a_\infty^- t} e^c \int_0^t (\|V(r)\|_{H^2}^2 + \|V(r)\|_{H^2}^3) dr \tag{3.27}$$

and without loss of generality the a priori boundedness – being proved a posteriori – of  $\|V(t)\|_{H^2}$  would be used to achieve

$$\|V(t)\|_{H^3}^2 \leq C \|V_0\|_{H^3}^2 e^{2a_\infty^- t} e^c \int_0^t (\|V(r)\|_{H^2}^2) dr. \tag{3.28}$$

Since we aim at exponential decay it will not matter whether we use (3.28), (3.27) or the statement of Lemma 3.2.

Using the representation (3.1) and Theorem 2.12 – observing that the nonlinearity satisfies the compatibility conditions for estimating the  $H^2$ -norm – we can estimate

$$\begin{aligned} \|V(t)\|_{H^2} &\leq \|e^{At} V_0\|_{H^2} + \int_0^t \|e^{(t-r)A} \mathcal{F}(V, V_x)(r)\|_{H^2} dr \\ &\leq c_1 e^{-\alpha_0 t} \|V_0\|_{H^2} + c_1 \int_0^t e^{-\alpha_0(t-r)} \|F(V, V_x)\|_{H^2} dr \end{aligned} \tag{3.29}$$

and it will be in the following estimate for  $\|F(V, V_x)\|_{H^2}$ , where we really use Assumption (1.5) to get an estimate that we need later on in the weighted a priori estimate.

**Lemma 3.3.**  $\exists c > 0 \forall W \in H^3 : \|F(W, W_x)\|_{H^2} \leq c \|W\|_{H^2}^2 \|W\|_{H^3}$ .

**Proof** (cf. [18] in  $\mathbb{R}^n$ ). Let  $u := W^1$ .

Using  $b(\tau) = \int_0^1 \int_0^1 b''(\mu\tau\nu) d\nu d\mu d\mu\tau^2$  we obtain

$$\begin{aligned} \|b(u_x)u_{xx}\|_{H^2} &\leq c(\|b(u_x)\|_\infty \|u_{xx}\|_{H^2} + \|b(u_x)\|_{H^2} \|u_{xx}\|_{L^\infty}) \\ &\leq c(\|u_x\|_{L^\infty}^2 \|u_x\|_{H^3} + \|u_x\|_{L^\infty} \|u_{xx}\|_{L^\infty} \|u_x\|_{H^3}) \\ &\leq c \|W\|_{H^2}^2 \|W\|_{H^3}. \quad \square \end{aligned}$$

Using Lemma 3.3 we conclude from (3.29) that

$$\|V(t)\|_{H^2} \leq c e^{-\alpha_0 t} \|V_0\|_{H^2} + c \int_0^t e^{-\alpha_0(t-r)} \|V(r)\|_{H^2}^2 \|V(r)\|_{H^3} dr \tag{3.30}$$

which is the starting point for proving the following weighted a priori estimate.

**Lemma 3.4.** For  $t \in [0, T)$ , the maximal interval of existence, let

$$M_2(t) := \sup_{0 \leq r \leq t} (e^{\tau_0 r} \|V(r)\|_{H^2})$$

where  $0 < \tau_0 \leq \alpha_0$ . Let (1.11) be satisfied, i.e.  $a_\infty^- < d_0$ . Then there are  $M_0 > 0$  and  $\delta > 0$  such that if  $\|V_0\|_{H^3} < \delta$  we have, for all  $0 \leq t \leq T$ ,

$$M_2(t) \leq M_0 < \infty$$

$M_0$  is independent of  $T$  (and of  $V_0$ ).

**Proof.** From (3.30) and the energy estimate in Lemma 3.2 we conclude that

$$\|V(t)\|_{H^2} \leq c \|V_0\|_{H^2} e^{-\alpha_0 t} + c \int_0^t e^{-\alpha_0(t-r)} \|V(r)\|_{H^2}^2 \|V_0\|_{H^3} e^{a_\infty^- r} e^c \int_0^r (\|V(\tau)\|_{H^2}^2 + \|V(\tau)\|_{H^2}^3 + \|V(\tau)\|_{H^2}^3) d\tau dr.$$

If  $\|V_0\|_{H^3} \leq \delta$  ( $\delta$  to be determined) we get

$$\begin{aligned} \|V(t)\|_{H^2} &\leq c\delta e^{-\alpha_0 t} + c\delta e^c \int_0^t (\|V(\tau)\|_{H^2}^2 + \|V(\tau)\|_{H^2}^3 + \|V(\tau)\|_{H^2}^3) d\tau \int_0^t e^{-\alpha_0(t-r)} e^{a_\infty^- r} \|V(r)\|_{H^2}^2 dr \\ &\leq c\delta e^{-\alpha_0 t} + c\delta e^{c(M_2(t)+M_2^2(t)+M_2^3(t))} \int_0^t e^{-\alpha_0 r} + e^{-2\alpha_0 r} + e^{-3\alpha_0 r} dr \\ &\quad \times M_2^2(t) \int_0^t e^{-\alpha_0(t-r)} e^{a_\infty^- r} e^{-2\alpha_0 r} dr \end{aligned}$$

which implies

$$M_2(t) \leq c\delta + c\delta e^{c(M_2(t)+M_2^2(t)+M_2^3(t))} M_2^2(t) \sup_{0 \leq t < \infty} e^{\alpha_0 t} \int_0^t e^{-\alpha_0(t-r)} e^{a_\infty^- r} e^{-2\alpha_0 r} dr. \tag{3.31}$$

Since by Assumption (1.11) it easily follows that

$$\sup_{0 \leq t < \infty} e^{\alpha_0 t} \int_0^t e^{-\alpha_0(t-r)} e^{a_\infty^- r} e^{-2\alpha_0 r} dr \leq c < \infty$$

we obtain from (3.31), for  $0 \leq t \leq T$ ,

$$M_2(t) \leq c\delta + c\delta M_2^2(t) e^{c(M_2(t)+M_2^2(t)+M_2^3(t))}. \tag{3.32}$$

By standard arguments (cf. e.g. [18]), considering the function

$$f(x) := c\delta(1 + cx^2 e^{c(x+x^2+x^3)}) - x$$

it follows that  $M_2(t)$  is uniformly bounded by the first zero  $M_0$  of  $f$  provided  $\delta$  and  $M_2(0)$  are sufficiently small.

This proves Lemma 3.4.  $\square$

Now we can formulate and prove the main theorem on global existence and exponential decay.

**Theorem 3.5.** *Let the Assumption (1.5) and (1.11) be satisfied. Then there exists  $\delta > 0$  such that if  $\|V_0\|_{H^3} < \delta$  there is a unique global solution  $u$  to (1.1)–(1.3) satisfying*

$$u \in \bigcap_{k=0}^3 C^k([0, \infty), H^{4-k}((0, L)) \cap H_0^1((0, L))) \cap C^4([0, \infty), L^2((0, L))).$$

Moreover there are constants  $c_0 = c_0(V_0) > 0$  and  $c_1 > 0$  such that

$$\|V(t)\|_{H^2} \leq c_0 e^{-\alpha_0 t}$$

and

$$\|V(t)\|_{H^3} \leq c_1 \|V_0\|_{H^3} e^{a_\infty^- t}, \quad t \geq 0.$$

**Proof.** From Lemmas 3.2 and 3.4 we conclude for the local solution

$$\begin{aligned} \|V(t)\|_{H^3} &\leq c \|V_0\|_{H^3} e^{a_\infty^- t} e^{c \int_0^t (\|V(r)\|_{H^2} + \|V(r)\|_{H^2}^2 + \|V(r)\|_{H^2}^3) dr} \\ &\leq c \|V_0\|_{H^3} e^{a_\infty^- t} e^{c(M_0 + M_0^2 + M_0^3)} \\ &\leq c \|V_0\|_{H^3} e^{a_\infty^- t}, \end{aligned}$$

$c$  being independent of  $t$  or  $V_0$ , from where the global existence follows by the usual continuation argument. The claim on the exponential decay of  $\|V(t)\|_{H^2}$  now is a consequence of Lemma 3.4.  $\square$

The Assumption (1.11), i.e.

$$a_\infty^- < \alpha_0,$$

together with the explicit estimates for  $\alpha_0$  from Section 2, just requires that the possibly existing negative part of  $a$  is not too large in comparison to its positive part.

### Acknowledgments

The authors thank Farid Ammar Khodja and Assia Benabdallah for discussions and for bringing the papers [3,5] to their attention, and also for mentioning the relation of [16] to Section 2.

### References

[1] A. Benaddi, B. Rao, Energy decay rate of wave equations with indefinite damping, J. Differential Equations 161 (2000) 337–357.

- [2] G. Chen, S.A Fulling, F.J. Narcowich, S. Sun, Exponential decay of energy of evolution equations with locally distributed damping, *SIAM J. Appl. Math.* 51 (1991) 266–301.
- [3] S.J. Cox, M.L. Overton, Perturbing the critically damped wave equation, *SIAM J. Appl. Math.* 56 (1994) 231–243.
- [4] C.M. Dafermos, W.J. Hrusa, Energy methods for quasilinear hyperbolic initial–boundary value problems. Applications to elastodynamics, *Arch. Ration. Mech. Anal.* 87 (1985) 267–292.
- [5] S. Cox, E. Zuazua, The rate at which energy decays in a damped string, *Comm. Partial Differential Equations* 19 (1994) 213–243.
- [6] P. Freitas, On some eigenvalue problems related to the wave equation with indefinite damping, *J. Differential Equations* 127 (1996) 320–335.
- [7] P. Freitas, E. Zuazua, Stability results for the wave equation with indefinite damping, *J. Differential Equations* 132 (1996) 338–353.
- [8] S. Jiang, R. Racke, Evolution equations in thermoelasticity, in:  $\pi$  Monographs Surveys Pure Appl. Math., vol. 112, Chapman&Hall/CRC, Boca Raton, 2000.
- [9] S. Kawashima, M. Nakao, K. Ono, On the decay property of solutions to the Cauchy problem of the semilinear wave equation with a dissipative term, *J. Math. Soc. Japan* 47 (1995) 617–653.
- [10] K. Liu, Z. Liu, R. Rao, Exponential stability of an abstract nondissipative linear system, *SIAM J. Control Optim.* 40 (2001) 149–165.
- [11] K. Liu, B. Rao, X. Zhang, Stabilization of the wave equations with potential and indefinite damping, *J. Math. Anal. Appl.* 269 (2002) 747–769.
- [12] Z. Liu, S. Zheng, Semigroups associated with dissipative systems, in:  $\pi$  Research Notes Math., vol. 398, Chapman&Hall/CRC, Boca Raton, 1999.
- [13] M. Nakao, Decay of solutions of the wave equation with a local nonlinear dissipation, *Math. Ann.* 305 (1996) 403–417.
- [14] M. Nakao, Global existence of smooth solutions to the initial boundary value problem for the quasi-linear wave equation with a localized degenerate dissipation, *Nonlinear Anal. TMA* 39 (2000) 187–205.
- [15] J. van Neerven, The asymptotic behaviour of semigroups of linear operators, in: *Operator Theory Advances Applications*, vol. 88, Birkhäuser, Basel, 1996.
- [16] A.F. Neves, H.de S. Ribeiro, O. Lopes, On the spectrum of evolution operators generated by hyperbolic systems, *J. Funct. Anal.* 67 (1986) 320–344.
- [17] J. Prüss, On the spectrum of  $C_0$ -semigroups, *Trans. Amer. Math. Soc.* 284 (1984) 847–857.
- [18] R. Racke, Lectures on nonlinear evolution equations. Initial value problems, in: *Aspects of Mathematics E19*, Friedr. Vieweg&Sohn, Braunschweig, Wiesbaden, 1992.
- [19] J. da Silva Ferreira, Exponential decay of the energy of a nonlinear system of Klein–Gordon equations with localized dampings in bounded and unbounded domains, *Asymptot. Anal.* 8 (1994) 73–92.
- [20] E. Zuazua, Exponential decay for the semilinear wave equation with locally distributed damping, *Comm. Partial Differential Equations* 15 (1990) 205–235.