Electronic communication in oligonuclear ferrocene complexes with anionic four-coordinate boron bridges†

Linda Kaufmann,^a Jens-Michael Breunig,^a Hannes Vitze,^a Frauke Schödel,^a Israel Nowik,^b Markus Pichlmaier,^c Michael Bolte,^a Hans-Wolfram Lerner,^a Rainer F. Winter,^c Rolfe H. Herber^b and Matthias Wagner^{*a}

The di- and trinuclear ferrocene species $\text{Li}[\text{Fc-BPh}_2\text{-Fc}]$ (**Li[9]**) and $\text{Li}_2[\text{Fc-BPh}_2\text{-fc-BPh}_2\text{-Fc}]$ (**Li₂[10]**) have been investigated with regard to their electrochemical properties and the degree of intervalence charge-transfer after partial oxidation. **Li[9]** shows two distinct one-electron redox waves for its chemically equivalent ferrocenyl substituents in the cyclic voltammogram ($E_{1/2} = -0.38$ V, -0.64 V; vs. FcH/FcH^+). The corresponding values of **Li₂[10]** are $E_{1/2} = -0.45$ V (two-electron process) and -1.18 V. All these redox events are reversible at r. t. on the time scale of cyclic voltammetry. X-ray crystallography on the mixed-valent $\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}$ complex $\text{Li}(\text{12-c-4})_2[\text{10}]$ reveals the centroid-centroid distance between the cyclopentadienyl rings of each of the terminal ferrocenyl substituents (3.329 Å) to be significantly smaller than in the central 1,1'-ferrocenediyl fragment (3.420 Å). This points towards a charge-localized structure (on the time scale of X-ray crystallography) with the central iron atom being in the Fe^{III} state. Mößbauer spectroscopic measurements on $\text{Li}(\text{12-c-4})_2[\text{10}]$ lend further support to this interpretation. Spectroelectrochemical measurements on **Li[9]** and **Li₂[10]** in the wavelength range between 300 – 2800 nm do not show bands interpretable as intervalence charge-transfer absorptions for the mixed-valent states. All data accumulated so far lead to the conclusion that electronic interaction between the individual Fe atoms in **Li[9]** and **Li₂[10]** occurs *via* a through-space pathway and/or is electrostatic in nature.

Introduction

Poly(ferrocenylene)s **A** (Fig. 1) represent an important class of processable metal-containing polymers with applications ranging from molecular electronics to the preparation of magnetic ceramics.^{1–4} The materials properties of macromolecules **A** depend to a large extent on the nature of the bridging element ER_x , which can be varied over a broad range (e.g. $\text{ER}_x = \text{SiMe}_2$,⁵ Sn^+Bu_2 ,⁶ PPh ,⁷ S^8). For the following reasons, our group is particularly interested in boron-bridged poly(ferrocenylene)s:⁹ (i) Three-coordinate boron atoms ($\text{ER}_x = \text{BR}'$) possess an empty *p*-orbital that is well-suited for π -conjugation with the cyclopentadienyl rings and thus able to act as an efficient transmitter of electronic interactions between the individual 1,1'-ferrocenediyl moieties.^{10,11} (ii) Four-coordinate boron atoms ($\text{ER}_x = \text{BR}'_2$) can bear a positive, a negative, or zero charge which provides a powerful set-screw for tuning the $\text{Fe}^{\text{II}}/\text{Fe}^{\text{III}}$ redox potentials of corresponding ferrocene oligomers by electrostatic means.^{12,13} (iii) The formation of B–L

adducts between poly(ferrocenylene)s with three coordinate boron bridges and Lewis bases (L) influences the degree of electronic communication along the polymer backbone, thereby offering an opportunity to design novel sensor compounds and switchable nanowires.

Given this background, a thorough evaluation of the degree of ferrocene–ferrocene interaction *via* three- and four-coordinate boron linkers and a comprehension of the underlying transmission pathways is essential for further rational developments in this area. We have already reported evidence for pronounced electronic communication along the chains of **A**-type polymers in which ER_x equals BMes^{10} or $\text{BO}(\text{CH}_2)_4\text{Br}^{11}$ ($\text{Mes} = \text{mesityl}$). There is also a strong indication for charge-transfer interactions between ferrocene and π^* -orbitals of the 4,4'-bipyridyl units in polymeric B–N adducts **B** (Fig. 1) even though the boron atoms are four-coordinate.^{14–17} The electronic structure, photophysics, and relaxation dynamics of charge-transfer excited states in soluble model systems of type $[\text{C}](\text{PF}_6)_2$ (Fig. 1) have been investigated using cyclic voltammetry, spectroelectrochemistry and laser spectroscopy.¹⁸ These results led to the conclusion that a four-coordinate boron atom is not necessarily an insurmountable barrier to electronic interactions between its redox-active substituents.

This conclusion is further substantiated by the fact that two different redox potentials are observed for the $\text{Fe}^{\text{II}}/\text{Fe}^{\text{III}}$ transitions in dinuclear complexes like **Li[D]** (Fig. 1) even though the two ferrocenyl substituents are chemically equivalent.^{12,13}

Most importantly in this context, a broad band with a maximum near $\lambda = 2200$ nm has been reported to appear in the UV/vis/NIR spectrum of the related ferricenyltris(ferrocenyl)borate zwitterion

^aInstitut für Anorganische Chemie, Goethe-Universität Frankfurt, Max-von-Laue-Strasse 7, 60438, Frankfurt (Main), Germany. E-mail: Matthias.Wagner@chemie.uni-frankfurt.de

^bRacah Institute of Physics, The Hebrew University of Jerusalem, 91904, Jerusalem, Israel

^cInstitut für Anorganische Chemie, Universität Regensburg, Universitätsstrasse 31, 93040, Regensburg, Germany

† Electronic supplementary information (ESI) available: Experimental details. CCDC reference numbers 699649 (3), 699647 (6), 699646 (7), 699650 ($[\text{Li}(\text{12-c-4})(\text{THF})_2][\text{10}]$), 699648 ($[\text{Li}(\text{12-c-4})_2][\text{10}]$) and 710783 ($[\text{Li}(\text{THF})_4][\text{11}]$). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/b821406a

Fig. 1 General representation of ER_x-bridged poly(ferrocenylene)s **A**; coordination polymer **B** showing charge-transfer interactions between the ferrocenylene fragments and the 4,4'-bipyridyl bridge; [C](PF₆)₂, a soluble model system of **B**; dinuclear BMe₂-bridged complex **Li[D]** possessing two different Fe^{II}/Fe^{III} redox potentials.

([Fc^{III}(Fc^{II})₃B]) and was interpreted as intervalence charge-transfer absorption (Fc = (C₅H₅)Fe(C₅H₄)).¹⁹ The question thus arises whether the electrochemical behaviour of **Li[D]**-type oligoferrocenes is merely governed by electrostatic interactions or whether charge delocalization also plays a significant role. In addition to **Li[D]**, our group has recently published the synthesis and structural characterization of its trinuclear congener Li₂[Fc-BMe₂-fc-BMe₂-Fc] (fc = (C₅H₄)₂Fe).¹³ Unfortunately, both compounds tend to decompose upon iron oxidation, so that cyclic voltammograms had to be recorded at -78 °C and all efforts regarding the isolation of mixed-valent species were unsuccessful.

Given the apparent stability of [Fc^{III}(Fc^{II})₃B], we postulated that replacement of the methyl substituents by phenyl groups in Li[Fc-BMe₂-Fc] (**Li[D]**) and Li₂[Fc-BMe₂-fc-BMe₂-Fc] will lead to increased stability and thus allow us to assess the level of electronic communication between the redox-active sites in greater detail. The results of our studies on Li[Fc-BPh₂-Fc] (**Li[9]**; Scheme 2)

and Li₂[Fc-BPh₂-fc-BPh₂-Fc] (**Li[10]**; Scheme 2) are outlined in this paper.

Results and discussion

Synthesis and NMR spectroscopy

Compounds Li[Fc-BMe₂-Fc] (**Li[D]**) and Li₂[Fc-BMe₂-fc-BMe₂-Fc] are readily accessible *via* B-C adduct formation between FcBMe₂ and FcLi or 1,1'-fcLi₂ × 2/3 TMEDA.¹³ We therefore considered the corresponding diphenylborylferrocenes **3** and **6** (Scheme 1) useful building blocks for the synthesis of BPh₂-bridged oligo- and polyferrocenes. FcBPh₂ (**3**) has already been prepared in the form of a red oil by Herberhold and Wrackmeyer, who treated FcLi with MeOBPh₂.²⁰ In our hands, the reaction between FcHgCl (**1**)²¹ and BrBPh₂ (**2**)²² resulted in better yields and higher purity of **3**.

The 1,1'-diborylated derivative **6** is not known in the literature so far, and its preparation turned out to be cumbersome. A synthesis approach analogous to the preparation of **3** is not practical due to the poor solubility of the doubly mercurated ferrocene 1,1'-fc(HgCl)₂. The following potential alternative routes resulted in inseparable product mixtures rather than in the formation of pure **6**: (i) 1,1'-fc(BBr₂)₂ and PhLi or SnPh₄ or PhSiMe₃, (ii) 1,1'-fc(B(OMe)₂)₂ and PhLi, (iii) 1,1'-fcLi₂ × 2/3 TMEDA and XBPh₂ (X = Br, O'Pr), (iv) Li₂[1,1'-fc(BPh₃)₂] and ClSiMe₃. We were finally able to synthesize **6** from 1,1'-fcLi₂ × 2/3 TMEDA (**4**) and 2 equiv. of MeOBPh₂, however, the target compound was contaminated with substantial amounts of Li(TMEDA)[MeOBPh₃] (**7**; Scheme 1). Both components precipitated from hexane as single crystals. Since it was not possible to completely separate the two products by fractional crystallization, we had to rely on manual crystal selection in order to obtain samples of reasonable purity for NMR-spectroscopic characterization.

Scheme 1 Synthesis of the borylated ferrocenes **3** and **6**. (i) hexane, -78 °C to r. t.

The dinuclear BPh_2 -bridged compound $\text{Li}[9]$ is readily accessible from Fc_2BBr (**8**) and 2 equiv. of PhLi (Scheme 2). Details of the synthesis protocol as well as an X-ray crystal structure determination of $\text{Li}(\text{OBU}_2)[9]$ have already been published elsewhere.²³ Attempts at the synthesis of the $\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}$ mixed-valent species **9** by reaction of $\text{Li}[9]$ with AgBF_4 , AgPF_6 or I_2 , always resulted in the formation of a red-brown oil that could not be transformed into an analytically pure solid. However, the UV/vis spectra of chemically oxidized $\text{Li}[9]$ proved to be identical to the spectrum obtained during electrochemical oxidation of this compound (see below). Due to its paramagnetic nature, interpretable NMR spectra of **9** could not be acquired.

Scheme 2 Synthesis of the di- and trinuclear BPh_2 -bridged ferrocene aggregates $\text{Li}[9]$ and $\text{Li}_2[10]$; oxidation of $\text{Li}_2[10]$ to its mixed-valent state $\text{Li}[10]$. (i) toluene/ OBU_2 , -78°C to r. t.; (ii) THF, -78°C to r. t.; (iii) THF, r. t.

The trinuclear species $\text{Li}_2[10]$ is formed from 2 equiv. of FcBPh_2 (**3**) and $1,1'\text{-fcLi}_2 \times 2/3 \text{ TMEDA}$ (**4**)^{24,25} under mild conditions and in good yields. In the presence of trace amounts of oxygen, $\text{Li}_2[10]$ is immediately transformed into its mixed-valent state $\text{Li}[10]$ (Scheme 2). $\text{Li}_2[10]$ and $\text{Li}[10]$ crystallized from THF/hexane in the presence of crown ether (12-c-4) as ether adducts ($\text{Li}(12\text{-c-4})(\text{THF})_2[10]$ and $\text{Li}(12\text{-c-4})_2[10]$), respectively.

For reasons of comparison, we decided to revisit the ferricenyltris(ferrocenyl)borate inner salt $[\text{Fc}^{\text{III}}(\text{Fc}^{\text{II}})_3\text{B}]^{19}$ (**11**; Scheme 3) and

Scheme 3 Synthesis of $\text{Li}[11]$; oxidation of $\text{Li}[11]$ to its mixed-valent state **11**. (i) THF/pentane, r. t.; (ii) CH_2Cl_2 , r. t.

also to synthesize and structurally characterize the fully reduced form $\text{Li}[11]$. $\text{Li}[11]$ is accessible from BF_3OEt_2 and excess FcLi , provided that strictly anaerobic conditions are maintained. Single crystals of $\text{Li}(\text{THF})_4[11]$ were grown from THF/pentane. The targeted oxidation of $\text{Li}(\text{THF})_4[11]$ to **11** was performed essentially as described in the literature (Scheme 3).¹⁹

The NMR data of **3**²⁰ and $\text{Li}[9]$ ²³ are in accord with published values. The ^{11}B NMR resonance of **6** is broadened beyond detection, probably as a result of slow intramolecular motion (note that already the monosubstituted analog **3** shows a very broad signal ($h_{1/2} = 600 \text{ Hz}$; $\delta(^{11}\text{B}) = 63.3$). All ^1H and ^{13}C NMR resonances of **6** are similar to those of **3** and therefore do not merit further discussion. The ^{11}B NMR spectrum of $\text{Li}_2[10]$ is characterized by a signal at -11.7 ppm which lies in a range typical of four-coordinate boron nuclei.²⁶ All proton resonances are broadened at r. t. A hump lacking any fine structure appears in the region of the C_5H_4 and C_5H_5 signals; the phenyl resonances are better resolved and appear at 6.63 ppm , 6.79 ppm , and 7.35 ppm . Similar to $\text{Li}_2[10]$, $\text{Li}[11]$ gives rise to a signal at -15.3 ppm in the ^{11}B NMR spectrum. All four ferrocenyl substituents are chemically equivalent ($\delta(^1\text{H}) = 3.69 (\text{C}_5\text{H}_5)$, 3.87 , $4.34 (\text{C}_5\text{H}_4)$). In the ^{13}C NMR spectrum, both C_5H_4 resonances of $\text{Li}[11]$ show a multiplet structure due to partially resolved ^{11}B coupling (the *ipso*-carbon signal is not observed due to quadrupolar broadening). Similar to **9**, the paramagnetic nature of $\text{Li}[10]$ and **11** precluded their characterization by NMR spectroscopy.

Crystal structure determinations

Selected crystallographic data of **3**, **6**, $\text{Li}(12\text{-c-4})(\text{THF})_2[10]$, $\text{Li}(12\text{-c-4})_2[10]$, and $\text{Li}(\text{THF})_4[11]$ are summarized in Tables 1

Table 1 Crystallographic data of **3** and **6**

compound	3	6
formula	C ₂₂ H ₁₉ BFe	C ₃₄ H ₂₈ B ₂ Fe
fw	350.03	514.03
colour, shape	red, needle	red, needle
temperature (K)	173(2)	173(2)
crystal system	tetragonal	orthorhombic
space group	<i>P</i> 4 ₃	<i>Fdd</i> 2
<i>a</i> (Å)	11.2142(7)	18.425(4)
<i>b</i> (Å)	11.2142(7)	30.690(6)
<i>c</i> (Å)	13.5550(11)	9.2970(19)
α (°)	90	90
β (°)	90	90
γ (°)	90	90
<i>V</i> (Å ³)	1704.7(2)	5257.1(18)
<i>Z</i>	4	8
<i>D</i> _{calcd.} (g cm ⁻³)	1.364	1.299
<i>F</i> (000)	728	2144
μ (mm ⁻¹)	0.883	0.595
crystal size (mm ³)	0.24 × 0.06 × 0.05	0.24 × 0.11 × 0.10
no. of rflns collected	21782	7049
no. of indep rflns (<i>R</i> _{int})	3214 (0.0716)	2204 (0.0758)
data/restraints/parameters	3214/1/217	2204/1/168
GOOF on <i>F</i> ²	1.025	1.010
<i>R</i> 1, <i>wR</i> 2 (<i>I</i> > 2σ(<i>I</i>))	0.0433, 0.1033	0.0372, 0.0710
<i>R</i> 1, <i>wR</i> 2 (all data)	0.0486, 0.1069	0.0521, 0.0788
largest diff peak and hole (e Å ⁻³)	0.461, -0.476	0.196, -0.236

and **2**; details of the crystal structure analysis of **7** are compiled in the ESI (Fig. 1S, Table 1S†).

3 (Fig. 2) and **6** (Fig. 3) feature planar three-coordinate boron atoms. In both compounds, the B–Cp bonds are shorter by about 0.04 Å than the B–Ph bonds, thereby indicating that the ferrocene fragment acts as a stronger π -donor than a phenyl ring. The resulting borafulvene character, which also results in a distinctive

C–C bond length alternation within the cyclopentadienyl rings of **3** and **6**, contributes to the electronic saturation of the electron-deficient boron atoms. In addition to that, there is a second type of interaction in borylated ferrocenes that involves filled *d*-type orbitals at iron and the empty *p*-orbital at boron and manifests itself by a bending of the boryl substituent out of the plane of the cyclopentadienyl ring towards the iron atom.^{27,28} In the cases of **3** and **6**, the corresponding dip angles α^* amount to 13.0° in **3** and 10.8° in the C₂-symmetric molecule **6** ($\alpha^* = 180^\circ - \alpha$, $\alpha = \text{COG}(\text{C}_5\text{H}_4) - \text{C}_{\text{ipso}} - \text{B}$; COG(C₅H₄): centroid of a cyclopentadienyl ring). As usual, the degree of bending is higher in the monoborylated than in the diborylated species. The α^* value of **3** is the same as in FcBMe₂ ($\alpha^* = 13.0^\circ$),²⁸ but it is smaller than the dip angle of FcBBr₂ ($\alpha^* = 18.3^\circ$)²⁷ and FcB(C₆F₅)₂ ($\alpha^* = 16^\circ$).²⁹ No X-ray crystal structure analyses of 1,1'-fc(BMe₂)₂ or 1,1'-fc(B(C₆F₅)₂)₂ are available to date. We therefore compare **6** with 1,1'-fc(BBrMe)₂ ($\alpha^* = 9.4^\circ$)²⁸ and 1,1'-fc(BBr₂)₂ ($\alpha^* = 9.1^\circ$)³⁰ which reveal a similar degree of ligand bending. The ferrocenyl substituent in **3** as well as the 1,1'-ferrocenediyl backbone in **6** adopt an eclipsed conformation (**3**: C(1)–COG(1)–COG(11)–C(11) = 7.2°, **6**: C(1)–COG(1)–COG(4A)–C(4A) = -4.0°; COG(X): centroid of the cyclopentadienyl ring containing the carbon atom C(X)). As a consequence, the torsion angle between the two boryl substituents of **6** has a value of B(1)–COG(1)–COG(1A)–B(1A) = 140.9°.

In the solid state, both trinuclear aggregates (Li(12-c-4)(THF))₂[**10**] (Fig. 4) and Li(12-c-4)₂[**10**] (Fig. 5) possess an inversion centre located at Fe(1). The Li⁺ ions are wrapped by ether ligands and do not establish short contacts with the anionic molecules. (Li(12-c-4)(THF))₂[**10**] contains two Li⁺ ions per oligoferrocene moiety, whereas in Li(12-c-4)₂[**10**] the cation : anion ratio is 1 : 1. This is in accord with the presence of three Fe^{II} ions in (Li(12-c-4)(THF))₂[**10**] but points towards two Fe^{II} ions and one Fe^{III} centre in Li(12-c-4)₂[**10**]. This conclusion is

Table 2 Crystallographic data of (Li(12-c-4)(THF))₂[**10**], Li(12-c-4)₂[**10**] and Li(THF)₄[**11**]

compound	(Li(12-c-4)(THF)) ₂ [10]	Li(12-c-4) ₂ [10]	Li(THF) ₄ [11]
formula	C ₇₈ H ₉₄ B ₂ Fe ₃ Li ₂ O ₁₀	C ₇₀ H ₇₈ B ₂ Fe ₃ LiO ₈ × C ₄ H ₈ O	C ₅₆ H ₆₈ BFe ₄ LiO ₄
fw	1394.58	1315.54	1046.25
colour, shape	orange, block	brown, needle	red, block
temperature (K)	173(2)	173(2)	173(2)
crystal system	monoclinic	orthorhombic	monoclinic
space group	<i>P</i> 2 ₁ / <i>n</i>	<i>Pnma</i>	<i>P</i> 2 ₁ / <i>n</i>
<i>a</i> (Å)	14.9703(10)	17.4879(19)	14.9829(9)
<i>b</i> (Å)	10.3522(6)	27.436(3)	20.0854(10)
<i>c</i> (Å)	22.9360(18)	15.5563(17)	16.7066(9)
α (°)	90	90	90
β (°)	105.526(5)	90	100.813(5)
γ (°)	90	90	90
<i>V</i> (Å ³)	3424.8(4)	7463.9(14)	4938.4(5)
<i>Z</i>	2	4	4
<i>D</i> _{calcd.} (g cm ⁻³)	1.352	1.171	1.407
<i>F</i> (000)	1472	2772	2192
μ (mm ⁻¹)	0.687	0.626	1.198
crystal size (mm ³)	0.23 × 0.21 × 0.18	0.30 × 0.12 × 0.11	0.37 × 0.33 × 0.32
no. of rflns collected	29466	33604	30721
no. of indep rflns (<i>R</i> _{int})	6418 (0.1120)	6709 (0.1866)	9233 (0.0628)
data/restraints/parameters	6418/48/539	6709/33/378	9233/0/595
GOOF on <i>F</i> ²	0.971	0.979	0.961
<i>R</i> 1, <i>wR</i> 2 (<i>I</i> > 2σ(<i>I</i>))	0.0510, 0.0971	0.1029, 0.1944	0.0394, 0.0901
<i>R</i> 1, <i>wR</i> 2 (all data)	0.0973, 0.1107	0.2222, 0.2438	0.0586, 0.0965
largest diff peak and hole (e Å ⁻³)	0.720, -0.433	0.718, -0.539	0.490, -0.521

Fig. 2 Structure of **3** in the crystal. Displacement ellipsoids are drawn at the 30% probability level. H atoms omitted for clarity. Selected bond lengths (Å), bond angles (°), and the dip angle α^* (°): B(1)–C(1) = 1.542(5), B(1)–C(21) = 1.579(5), B(1)–C(31) = 1.581(5); C(1)–B(1)–C(21) = 122.4(3), C(1)–B(1)–C(31) = 121.1(3), C(21)–B(1)–C(31) = 116.5(3); α^* = 13.0.

Fig. 3 Structure of **6** in the crystal. Displacement ellipsoids are drawn at the 50% probability level. H atoms omitted for clarity. Selected bond lengths (Å), bond angles (°), and the dip angle α^* (°): B(1)–C(1) = 1.542(5), B(1)–C(11) = 1.585(6), B(1)–C(21) = 1.584(6); C(1)–B(1)–C(11) = 119.2(3), C(1)–B(1)–C(21) = 123.1(3), C(11)–B(1)–C(21) = 117.6(3); α^* = 10.8.

further supported by the observation of almost equal distances between the cyclopentadienyl rings of the three ferrocene fragments in $(\text{Li}(12\text{-c-4})(\text{THF}))_2[\mathbf{10}]$ (COG(1)–COG(1A) = 3.330 Å, COG(11)–COG(21) = 3.325 Å), with values characteristic of ferrocenes containing Fe^{II} ions.^{19,31} In contrast, $\text{Li}(12\text{-c-4})_2[\mathbf{10}]$ exhibits similar centroid–centroid distances only for the terminal ferrocenyl substituents (COG(21)–COG(31) = 3.329 Å), while the central moiety is significantly expanded (COG(11)–COG(11A) = 3.420 Å). We take this as evidence that Fe(1) in $\text{Li}(12\text{-c-4})_2[\mathbf{10}]$ is an Fe^{III} centre.^{19,31} The B–Cp and B–Ph bond lengths are

the same in $(\text{Li}(12\text{-c-4})(\text{THF}))_2[\mathbf{10}]$ and in $\text{Li}(12\text{-c-4})_2[\mathbf{10}]$ (*cf.* Fig. 4 and 5). This is in agreement with *a priori* expectations, because differences in the π -donor strengths of both aromatic substituents are only relevant for three-coordinate boranes like **3** and **6** but not for compounds containing four-coordinate boron atoms. The central 1,1'-ferrocenediyl fragments in $(\text{Li}(12\text{-c-4})(\text{THF}))_2[\mathbf{10}]$ and $\text{Li}(12\text{-c-4})_2[\mathbf{10}]$ adopt staggered conformations with boron substituents pointing in opposite directions. The major difference in the overall conformations of both oligomers lies in the position of the terminal ferrocenyl substituents

Fig. 4 Structure of $(\text{Li}(12\text{-c-4})(\text{THF}))_2[\mathbf{10}]$ in the crystal. Displacement ellipsoids are drawn at the 50% probability level. H atoms and $\text{Li}(12\text{-c-4})(\text{THF})^+$ counterions omitted for clarity. Selected bond lengths (Å), atom...atom distances (Å), and bond angles ($^\circ$): $\text{B}(1)\text{--C}(1) = 1.658(4)$, $\text{B}(1)\text{--C}(11) = 1.646(5)$, $\text{B}(1)\text{--C}(31) = 1.647(5)$, $\text{B}(1)\text{--C}(41) = 1.663(4)$, $\text{COG}(1)\text{--COG}(1A) = 3.330$, $\text{COG}(11)\text{--COG}(21) = 3.325$, $\text{Fe}(1)\cdots\text{Fe}(2) = 6.112(1)$; $\text{C}(1)\text{--B}(1)\text{--C}(11) = 105.3(3)$, $\text{C}(31)\text{--B}(1)\text{--C}(41) = 106.9(3)$. $\text{COG}(X)$: centroid of the cyclopentadienyl ring containing the carbon atom $\text{C}(X)$. Symmetry transformations used to generate equivalent atoms: $A: -x + 1, -y, -z + 1$.

Fig. 5 Structure of $\text{Li}(12\text{-c-4})_2[\mathbf{10}]$ in the crystal. Displacement ellipsoids are drawn at the 50% probability level. H atoms, the $\text{Li}(12\text{-crown-4})^+$ counterion and the THF molecule omitted for clarity. Selected bond lengths (Å), atom...atom distances (Å), and bond angles ($^\circ$): $\text{B}(1)\text{--C}(11) = 1.646(12)$, $\text{B}(1)\text{--C}(21) = 1.644(12)$, $\text{B}(1)\text{--C}(41) = 1.649(13)$, $\text{B}(1)\text{--C}(51) = 1.649(12)$, $\text{COG}(11)\text{--COG}(11A) = 3.420$, $\text{COG}(21)\text{--COG}(31) = 3.329$, $\text{Fe}(1)\cdots\text{Fe}(2) = 5.477(1)$; $\text{C}(11)\text{--B}(1)\text{--C}(21) = 112.4(6)$, $\text{C}(41)\text{--B}(1)\text{--C}(51) = 111.3(6)$. $\text{COG}(X)$: centroid of the cyclopentadienyl ring containing the carbon atom $\text{C}(X)$. Symmetry transformations used to generate equivalent atoms: $A: -x + 1, -y + 1, -z$.

with respect to the central ferrocene moiety (dihedral angles: $\text{Li}(12\text{-c-4})(\text{THF})_2[\mathbf{10}]$: $\text{Cp}(\text{C}(1))//\text{Cp}(\text{C}(11)) = 83.0^\circ$; $\text{Li}(12\text{-c-4})_2[\mathbf{10}]$: $\text{Cp}(\text{C}(11))//\text{Cp}(\text{C}(21)) = 41.1^\circ$).

The crystal lattice of the tetrakis(ferrocenyl)borate salt $\text{Li}(\text{THF})_4[\mathbf{11}]$ contains tetrahedrally coordinated cations and anions; the molecular structure of the $[\text{BFc}_4]^-$ ion is shown in Fig. 6. We note a nearly threefold axis normal to the plane defined by $\text{Fe}(1)$, $\text{Fe}(2)$, and $\text{Fe}(3)$. As a result, the fourth ferrocenyl unit, *i.e.* that labeled $\text{Fe}(4)$ in Fig. 6, is distinct from the others. Thus, the conformation of $[\text{BFc}_4]^-$ in $\text{Li}(\text{THF})_4[\mathbf{11}]$ is strikingly similar to that of the neutral mixed-valent species $\mathbf{11}$.¹⁹ All B–C bond lengths of $\text{Li}(\text{THF})_4[\mathbf{11}]$ fall in the interval between $\text{B}(1)\text{--C}(21) = 1.634(4)$ Å and $\text{B}(1)\text{--C}(11) = 1.653(4)$ Å; the smallest C–B–C angle is $\text{C}(11)\text{--B}(1)\text{--C}(31) = 103.2(2)^\circ$, the largest $\text{C}(21)\text{--B}(1)\text{--C}(31) = 113.0(2)^\circ$. Most importantly, all four centroid-to-centroid distances within the ferrocenyl substituents lie between 3.312 Å and 3.318 Å and are thus characteristic of Fe^{II} states. This is in contrast to the molecular structure reported for the mixed-valent species $\mathbf{11}$, which contains three ferrocenyl groups with short centroid-to-centroid distances (3.291 Å to 3.320 Å; Fe^{II} states), and one substituent with a significantly elongated centroid-to-centroid distance (3.428 Å; Fe^{III} state).¹⁹

Electrochemical and spectroelectrochemical investigations

The electrochemical parameters of the redox events exhibited by $\text{Li}[\mathbf{9}]$,²³ $\text{Li}_2[\mathbf{10}]$, $\text{Li}[\mathbf{10}]$ and the related complexes $\text{Li}[\text{Fc-BMe}_2\text{-Fc}]^{13}$ and $\text{Li}_2[\text{Fc-BMe}_2\text{-fc-BMe}_2\text{-Fc}]^{13}$ are summarized in Table 3.

Each of the dinuclear complexes $\text{Li}[\mathbf{9}]$ and $\text{Li}[\text{Fc-BMe}_2\text{-Fc}]$ displays two oxidation processes of relative intensity 1:1, assignable as successive one-electron transitions at the two ferrocenyl moieties. In the case of the phenyl derivative $\text{Li}[\mathbf{9}]$, both electron transitions are reversible at r. t. on the cyclic voltammetric timescale, whereas the methyl derivative $\text{Li}[\text{Fc-BMe}_2\text{-Fc}]$ has to be examined at -78°C in order to obtain cyclic voltammograms showing features

Table 3 Formal electrode potentials $E_{1/2}$ (vs. FcH/FcH⁺) and peak-to-peak separations ΔE (at 0.1 V s⁻¹) for the Fe^{II}/Fe^{III} redox processes exhibited by compounds Li[9],²³ Li₂[10], Li[10], Li[Fc-BMe₂-Fc],¹³ and Li₂[Fc-BMe₂-fc-BMe₂-Fc]¹³

	$E_{1/2}$ [V]	ΔE [mV]	$\Delta E_{1/2}$ [mV]	solvent	$\Delta E(\text{FcH})$ [mV]
Li[9]	-0.38/-0.64	99/97	260	CH ₂ Cl ₂	103
Li ₂ [10]	-0.45/-1.18	125/82	730	CH ₂ Cl ₂	122
Li[10]	-0.45/-1.18	119/85	730	CH ₂ Cl ₂	108
Li[Fc-BMe ₂ -Fc] ^a	-0.43/-0.64	90/100	210	CH ₂ Cl ₂	100
Li ₂ [Fc-BMe ₂ -fc-BMe ₂ -Fc] ^a	-0.51/-1.21	330/210	700	CH ₂ Cl ₂	250

^a recorded at a scan rate of 0.2 V s⁻¹ and at a temperature of -78 °C.

Fig. 6 Structure of Li(THF)₄[11] in the crystal. Displacement ellipsoids are drawn at the 50% probability level. H atoms and the Li(THF)₄⁺ counterion omitted for clarity. Selected bond lengths (Å), atom...atom distances (Å), and bond angles (°): B(1)–C(1) = 1.648(4), B(1)–C(11) = 1.653(4), B(1)–C(21) = 1.634(4), B(1)–C(31) = 1.644(4), COG(1)–COG(6) = 3.312, COG(11)–COG(16) = 3.318, COG(21)–COG(26) = 3.314, COG(31)–COG(36) = 3.314, av. Fe...Fe = 5.675(1); C(1)–B(1)–C(11) = 109.6(2), C(1)–B(1)–C(21) = 109.6(2), C(1)–B(1)–C(31) = 111.8(2), C(11)–B(1)–C(21) = 109.4(2), C(11)–B(1)–C(31) = 103.2(2), C(21)–B(1)–C(31) = 113.0(2). COG(X): centroid of the cyclopentadienyl ring containing the carbon atom C(X).

of chemical reversibility. Moreover, when also determined at a temperature of -78 °C, both $E_{1/2}$ values of Li[9] (-0.35 V/-0.57 V) are anodically shifted by about 0.08 V with respect to the redox potentials of Li[Fc-BMe₂-Fc] (-0.43 V/-0.64 V), which can easily be explained by the greater electronegativity of the phenyl rings as compared to methyl substituents.

The cyclic voltammograms of the trinuclear complexes Li₂[10] and Li[10] are congruent to each other and reveal two redox events with an intensity ratio of 1 : 2 (Fig. 7; note that the different oxidation states of Li₂[10] and Li[10] have been confirmed by linear sweep voltammetry).

Both these processes are chemically reversible as evidenced by the following criteria: the current ratios i_{pc}/i_{pa} are constantly equal to 1, the current functions $i_{pa}/v^{1/2}$ remain constant, and the peak-to-peak separations (ΔE) do not depart appreciably from the value

Fig. 7 Cyclic voltammogram of Li₂[10] (CH₂Cl₂, [NBu₄][PF₆] as supporting electrolyte (0.1 M), scan rate 0.1 V s⁻¹; vs. FcH/FcH⁺).

found for the internal ferrocene standard ($\Delta E(\text{FcH})$, Table 3; theoretically expected value for a chemically and electrochemically reversible one-electron step: 59 mV). The more anodic two-electron transfer at $E_{1/2} = -0.45$ V can be attributed to the terminal ferrocenyl moieties, while the less anodic one-electron redox event at $E_{1/2} = -1.18$ V takes place at the interior iron centre. These data agree with the results obtained from electrochemical investigations of Li₂[Fc-BMe₂-fc-BMe₂-Fc]. However, as in the case of Li[9] and Li[Fc-BMe₂-Fc], the $E_{1/2}$ values of Li₂[10]/Li[10] are slightly shifted to the anodic regime compared to the redox potentials of Li₂[Fc-BMe₂-fc-BMe₂-Fc] (Table 3).

Oxidation of the 1,1'-ferrocenylene unit in Li₂[10] takes place at a much more cathodic redox potential than oxidation of the terminal ferrocenyl groups ($\Delta E_{1/2} = 730$ mV) which is clearly due to the fact that the former has two negatively charged substituents, while each of the latter bears only one such group. The Fc moieties of Li[9] are, however, chemically equivalent. The comparatively large differences of more than 200 mV between the redox potentials of the two Fe^{II}/Fe^{III} transitions in Li[9] as well as Li[Fc-BMe₂-Fc] indicate that the two Fc subunits are mutually interacting. The question thus arises whether this interaction is entirely electrostatic in nature or whether there is a certain degree of charge delocalization *via* four-coordinate boron linkers or even through-space. The latter assumption is supported by the finding of a very broad band with a maximum near 2200 nm in the electronic spectrum of the Fe^{II}₃Fe^{III} species 11.¹⁹ It has been suggested that this absorption is due to intervalence charge-transfer processes which may proceed by a through-space mechanism.

In view of this background, we decided to carry out spectroelectrochemical measurements on the dinuclear compound Li[9] and to look for intervalence charge-transfer (IVCT) bands. To this end, we have performed a coulometrically controlled

one-electron oxidation at a working potential of -0.5 V (CH_2Cl_2 , $[\text{NBu}_4][\text{PF}_6]$ (0.1 M)) and simultaneously recorded the changes in the UV/vis/NIR spectrum of the solution. Prior to oxidation, we observe the lowest-energy band at $\lambda_{\text{max}} = 470$ nm which compares reasonably well to the absorption band at $\lambda_{\text{max}} = 442$ nm exhibited by parent ferrocene under the same conditions. After exhaustive one-electron oxidation, the resulting $\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}$ species **9** still showed a band at $\lambda_{\text{max}} = 471$ nm and, in addition, an absorption at $\lambda_{\text{max}} = 698$ nm (*cf.* $[(\text{C}_5\text{H}_5)_2\text{Fe}]\text{PF}_6$: $\lambda_{\text{max}} = 635$ nm). After exhaustive two-electron oxidation at an applied voltage of 0.1 V, only a very broad band at $\lambda_{\text{max}} = 619$ nm remained in this region of the spectrum. Most importantly, at no stage did we observe any feature at longer wavelengths than 900 nm that might be interpretable as $\text{Fe}^{\text{II}}/\text{Fe}^{\text{III}}$ IVCT band. An IVCT absorption is also absent in the electronic spectrum of the $\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}$ mixed-valent complex **Li[10]**. In order to test our spectroelectrochemical setup we also re-investigated the tetraferrocenylborate **Li[11]** and have fully reproduced the published spectral data¹⁹ including the IVCT band. We have moreover observed that upon further oxidation of mixed-valent **11** ($\text{Fe}^{\text{II}}_3\text{Fe}^{\text{III}}$) to **[11]⁺** ($\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}_2$) and to **[11]²⁺** ($\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}_3$) the broad featureless NIR band first intensifies and then decreases in intensity (*cf.* Fig. 2S of the ESI†). Upon potential reversal and subsequent reduction these changes are reversible.

⁵⁷Fe Mößbauer spectroscopy

Since there are no analytically pure and solid samples of the mixed-valent $\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}$ species **9** available, we had to restrict our Mößbauer spectroscopic studies to the trinuclear $\text{Fe}^{\text{II}}_2\text{Fe}^{\text{III}}$ compound **Li[10]**.

Mößbauer data were acquired on single crystals of $\text{Li}(\text{12-c-4})_2[\text{10}]$ in the temperature range $97.5 \text{ K} \leq T \leq 304 \text{ K}$. We first consider the spectrum taken at 97.5 K which is shown in the lower trace of Fig. 8. The spectrum consists of two distinct iron sites, Fe and Fe', and the relative area under the resonance curve is $\text{Fe} : \text{Fe}' \approx 2 : 1$. This already suggests Fe to correspond to the Fe^{II} sites and Fe' to the unique Fe^{III} site of $\text{Li}(\text{12-c-4})_2[\text{10}]$. At 90 K , the isomer shifts (IS) are $0.529 \pm 0.002 \text{ mm s}^{-1}$ (Fe) and $0.50 \pm 0.04 \text{ mm s}^{-1}$ (Fe') and the corresponding quadrupole splittings (QS) are $2.379 \pm 0.002 \text{ mm s}^{-1}$ (Fe) and $-0.24 \pm 0.02 \text{ mm s}^{-1}$ (Fe'; Note: It is not always possible to record Mößbauer spectra at exactly 90 K . To effect intersample comparison, the hyperfine parameter data have been linearly extrapolated to 90 K and those values are reported herein). The QS of the Fe' site is negative, as has previously been observed for ferricinium centres in related compounds.³²

The qualitative picture remains the same irrespective of the temperature applied during measurements. This leads to the conclusion that there is no electron delocalization over the three iron centres in the entire interval $97.5 \text{ K} \leq T \leq 304 \text{ K}$. However, it should be noted that at higher temperatures the Fe^{III} resonance sharpens and becomes a well-defined doublet, as shown in the upper trace of Fig. 8. This observation is consistent with a spin-lattice relaxation process (*vide infra*).

The temperature dependence of the IS for the Fe^{II} site Fe can be fitted by a linear regression with a correlation coefficient of 0.992 for 11 data points (slope = $-(4.12 \pm 0.15) \cdot 10^{-4} \text{ mm s}^{-1} \text{ K}^{-1}$). From this temperature dependence, the effective vibrating mass of the metal centre is calculated to be $M_{\text{eff}} = 101 \pm 3$ Daltons. Likewise, the ln of the temperature dependence of the recoil-free fraction

Fig. 8 ^{57}Fe Mößbauer spectrum of $\text{Li}(\text{12-c-4})_2[\text{10}]$ at 97.5 K (lower trace). The velocity scale is with respect to the centroid of a r. t. α -Fe spectrum. The corresponding spectrum at 304 K is shown in the upper trace and confirms the absence of electron delocalization between the iron sites, as discussed in the text.

averaged over both types of iron sites (as determined from the areas under the resonance curves) is linear over the whole temperature range (slope = $-(9.89 \pm 0.44) \cdot 10^{-3} \text{ K}^{-1}$; correlation coefficient = 0.99 for 7 data points). From these data it is possible to calculate the parameter $F_{\text{M173}} = k^2 \langle x_{\text{ave}}^2 \rangle$ (k : wave vector of the Mößbauer γ -ray; x_{ave}^2 : average root-mean-square amplitude of vibration of the iron atom), which can be compared to the parameter F_{X173} as it has been extracted from the U_{ij} value determined by X-ray crystallography on $\text{Li}(\text{12-c-4})_2[\text{10}]$ at 173 K . This comparison reveals the two parameters, $F_{\text{M173}} = 1.70$ and $F_{\text{X173}} = 1.73$, to be in excellent agreement with each other. The root-mean-square amplitudes of vibration of the Fe atoms in $\text{Li}(\text{12-c-4})_2[\text{10}]$ derived from the Mößbauer data are 0.193 Å , 0.215 Å , and 0.236 Å at 200 K , 250 K , and 300 K , respectively.

As mentioned above, the paramagnetic Fe^{III} centre Fe' relaxes by spin-lattice relaxation and obeys a fifth order power law, indicative of a Raman process, as has previously been reported for other $S = 5/2$ spin systems.³³ The relaxation is slow (on the Mößbauer time scale) below 150 K , but becomes rapid at higher temperatures (*cf.* Fig. 9 for a plot of the temperature dependence of the relaxation rate).

Conclusion

We have shown that the molecular framework of BPh_2 -bridged oligoferrocenes $\text{Li}[\text{Fc-BPh}_2\text{-Fc}]$ (**Li[9]**) and $\text{Li}_2[\text{Fc-BPh}_2\text{-fc-BPh}_2\text{-Fc}]$ (**Li[10]**) remains intact when the iron atoms are oxidized (Fc:

Fig. 9 Temperature dependence of the spin-lattice relaxation process for the Fe^{III} site in $\text{Li}(\text{12-c-4})_2[\text{10}]$. The solid line represents a fifth order temperature dependence indicative of a Raman-type relaxation process.

$(\text{C}_5\text{H}_5)\text{Fe}(\text{C}_5\text{H}_4)$). A corresponding $\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}$ mixed-valent species $\text{Li}[\text{Fc-BPh}_2\text{-fc-BPh}_2\text{-Fc}]$ ($\text{Li}[\text{10}]$) has been structurally characterized by X-ray crystallography. Both the crystallographical data and ^{57}Fe Mößbauer spectroscopy on $\text{Li}[\text{10}]$ point towards a largely localized electronic structure with the central iron atom adopting an oxidation state of +III. Moreover, spectroelectrochemical measurements on $\text{Li}[\text{9}]$ and $\text{Li}_2[\text{10}]$ in the UV/vis/NIR region do not reveal absorption bands assignable to intervalence charge-transfer processes after partial oxidation of the compounds. This is in striking contrast to published data (which have been confirmed by our own measurements) on the $\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}$ mixed-valent complex Fc_4B (**11**) for which electron delocalization has been observed. To account for this different behaviour, we offer two explanations: (i) In the dinuclear molecule $[\text{Fe}^{\text{III}}\text{Fe}^{\text{II}}\text{BPh}_2]$ (**9**), $\text{Fe}^{\text{II}} \rightarrow \text{Fe}^{\text{III}}$ charge transfer is three times less likely than in the tetranuclear compound $[\text{Fe}^{\text{III}}(\text{Fc}^{\text{II}})_3\text{B}]$ (**11**). As a result, the absorbance of a hypothetical IVCT band of **9** should be considerably smaller than the corresponding value of the IVCT band of **11**, which is already rather low ($\epsilon = 300$). (ii) If electron transfer in **11** proceeds via a through-space mechanism (as has been suggested by Cowan *et al.*¹⁹) the average $\text{Fe} \cdots \text{Fe}$ distance becomes a decisive factor. Since the ferrocenyl substituents in **11** are much more densely packed than in **9**, a through-space charge-transfer operative in **11** may well no longer be possible in **9** (*cf.* the solid state structure of $\text{Li}(\text{OBu}_2)[\text{9}]^{23}$ shows that conformations of this molecule are possible in which the $\text{Fe} \cdots \text{Fe}$ distance is as long as 6.679(1) Å; in contrast, the average $\text{Fe} \cdots \text{Fe}$ distance in the less flexible molecule $\text{Li}(\text{THF})_4[\text{11}]$ is only 5.675(1) Å).

We therefore conclude that electronic interaction between the individual iron sites in $\text{Li}[\text{9}]$, **9**, $\text{Li}_2[\text{10}]$, and $\text{Li}[\text{10}]$ does probably not occur by charge delocalization via the BPh_2 -bridge, but is either a through-space process or of an electrostatic nature. In this context, it is interesting to compare the results of Curtis *et al.* on the degree of electronic communication within mixed-valent poly(ferrocenylenearylene)s.³⁴ These authors find very similar behaviour to our oligomers (*e.g.* significant ferrocene-ferrocene interaction as measured by cyclic voltammetry, but little electron transfer as measured by NIR and Mößbauer spectroscopy), even though the polymer chains in poly(ferrocenylenearylene)s are

conjugated, whereas in **9** and **10** there is no π -conjugation between the redox-active moieties.

Experimental

General considerations

All reactions were carried out under a nitrogen atmosphere using Schlenk tube techniques. Solvents were freshly distilled under argon from Na/benzophenone (diethyl ether, THF, d_8 -THF), Na/Pb alloy (pentane, hexane) or CaH_2 (CH_2Cl_2 , CDCl_3) prior to use. NMR spectrometers: Bruker AM 250, AV 300 and AMX 400. Chemical shifts are referenced to residual solvent peaks (^1H , $^{13}\text{C}\{^1\text{H}\}$) or external $\text{BF}_3 \cdot \text{Et}_2\text{O}$ ($^{11}\text{B}\{^1\text{H}\}$). Abbreviations: s = singlet, d = doublet, tr = triplet, mult = multiplet, br = broad, o = ortho, m = meta, p = para, n.o. = not observed, n.r. = not resolved. All NMR spectra were run at r. t. Electrochemical measurements: Potentiostat EG&G Princeton Applied Research 263 A. Compounds FcHgCl (**1**),²¹ BrBPh_2 (**2**),²² FcLi ,³⁵ and $1,1'\text{-fcLi}_2 \times 2/3$ TMEDA (**4**)^{24,25} were synthesized according to literature procedures. The synthesis of MeOBPh_2 (**5**) is described in the ESI.†

Synthesis of 3. A solution of BrBPh_2 **2** (0.53 g, 2.16 mmol) in hexane (12 mL) was added dropwise with stirring at -78°C to a suspension of FcHgCl **1** (0.91 g, 2.16 mmol) in hexane (35 mL). The reaction mixture was slowly warmed to r. t. and stirred overnight, whereupon a grey solid precipitated. After filtration, the filtrate was slowly evaporated *in vacuo* to a volume of 5 mL whereupon single crystals of **3** formed. Yield: 0.60 g (80%). $^{11}\text{B}\{^1\text{H}\}$ NMR (128.4 MHz, CDCl_3): δ 63.3 ($h_{1/2} = 600$ Hz). ^1H NMR (400.1 MHz, CDCl_3): δ 4.18 (s, 5H, C_5H_5), 4.54, 4.82 ($2 \times$ n.r., $2 \times 2\text{H}$, C_5H_4), 7.40–7.46 (mult, 6H, *m*-Ph, *p*-Ph), 7.75 (d, 4H, $^3J_{\text{HH}} = 7.6$ Hz, *o*-Ph). $^{13}\text{C}\{^1\text{H}\}$ NMR (100.6 MHz, CDCl_3): δ 69.4 (C_5H_5), 75.8, 78.4 (C_5H_4), 127.2 (*m*-Ph), 129.4 (*p*-Ph), 135.4 (*o*-Ph), n.o. (BC). Elemental analysis: Calcd. for $\text{C}_{22}\text{H}_{19}\text{BFe}$ (350.03): C, 75.49; H, 5.47. Found: C, 75.23; H, 5.50%.

Synthesis of 6 and 7. A solution of MeOBPh_2 **5** (0.36 g, 1.86 mmol) in hexane (10 mL) was added dropwise with stirring at -78°C to a suspension of $1,1'\text{-fcLi}_2 \times 2/3$ TMEDA **4** (0.26 g, 0.93 mmol) in hexane (15 mL). The reaction mixture was slowly warmed to r. t. and stirred overnight. The resulting orange suspension was filtered and the filtrate was evaporated to a volume of 5 mL. Red needles of **6** and colourless plates of **7** crystallized after the solution had been stored for several months at -35°C . The crystals were separated by manual selection in a glovebox. NMR data of **6**: $^{11}\text{B}\{^1\text{H}\}$ NMR (128.4 MHz, CDCl_3): δ n.o. ^1H NMR (400.1 MHz, CDCl_3): δ 4.54, 4.75 ($2 \times$ n.r., $2 \times 4\text{H}$, C_5H_4), 7.39 (mult, 8H, *m*-Ph), 7.47 (mult, 4H, *p*-Ph), 7.68 (d, 8H, $^3J_{\text{HH}} = 7.2$ Hz, *o*-Ph). $^{13}\text{C}\{^1\text{H}\}$ NMR (100.6 MHz, CDCl_3): δ 76.6, 79.4 (C_5H_4), 127.3 (*m*-Ph), 129.9 (*p*-Ph), 135.5 (*o*-Ph), n.o. (BC).

NMR data of **7**: $^{11}\text{B}\{^1\text{H}\}$ NMR (128.4 MHz, d_8 -THF): δ 2.5 ($h_{1/2} = 150$ Hz). ^1H NMR (400.1 MHz, d_8 -THF): δ 2.15 (s, 12H, NMe), 2.31 (s, 4H, NCH_2), 3.11 (s, 3H, OMe), 6.86 (tr, 3H, $^3J_{\text{HH}} = 7.6$ Hz, *p*-Ph), 7.00 (mult, 6H, *m*-Ph), 7.41 (d, 6H, $^3J_{\text{HH}} = 7.2$ Hz, *o*-Ph). $^{13}\text{C}\{^1\text{H}\}$ NMR (100.6 MHz, d_8 -THF): δ 46.4 (NMe), 59.1 (NCH_2), 52.2 (OMe), 124.0 (*p*-Ph), 126.9 (*m*-Ph), 135.3 (*o*-Ph), n.o. (BC).

Synthesis of Li₂[10] and Li[10]. A solution of FcBPh₂ **3** (0.15 g, 0.43 mmol) in THF (8 mL) was added dropwise with stirring at -78 °C to a solution of 1,1'-FcLi₂ × 2/3 TMEDA **4** (0.06 g, 0.22 mmol) in THF (12 mL). The reaction mixture was slowly warmed to r. t. and stirred overnight. The volume of the solution was first reduced to 4 mL *in vacuo* and then 12-crown-4 (0.27 mL) and hexane (10 mL) were added. The resulting red precipitate was extracted with hexane (3 × 10 mL). Single crystals of (Li(12-c-4)(THF))₂[10] were grown by gas-phase diffusion of hexane into a THF solution of the crude product under strict exclusion of air. Single crystals of Li(12-c-4)₂[10] were grown under similar conditions but without strict exclusion of air. Yield of (Li(12-c-4)(THF))₂[10]: 0.10 g (33%). Yield of Li(12-c-4)₂[10]: 0.030 g (10%). NMR data of (Li(12-c-4)(THF))₂[10]: ¹B{¹H} NMR (128.4 MHz, *d*₈-THF): δ -11.7 (*h*_{1/2} = 70 Hz). ¹H NMR (400.1 MHz, *d*₈-THF): δ 3.59 (s, 32H, 12-c-4), 3.68 (very br, C₅H₄/C₅H₅), 6.63 (br, 4H, *p*-Ph), 6.79, 7.35 (2 × br, 2 × 8H, *o,m*-Ph). Elemental analysis: Calcd. for C₇₈H₉₄B₂Fe₃Li₂O₁₀ (1394.58): C, 67.18; H, 6.79. Found: C, 67.40; H, 6.89%. Elemental analysis: Calcd. for C₇₀H₇₈B₂Fe₃LiO₈ (1243.43) × C₄H₈O (72.11): C, 67.56; H, 6.59. Found: C, 67.71; H, 6.62%.

Synthesis of Li[11]. Ferrocene (5.00 g, 26.88 mmol) was treated with THF (25 mL) and the mixture cooled to 0 °C. *tert*-BuLi in pentane (1.6 M; 14 mL, 22.4 mmol) was added dropwise with stirring to the amber coloured slurry, whereupon the colour changed to red. The reaction mixture was stirred at 0 °C for another 15 min and allowed to warm to r. t. BF₃·OEt₂ (0.37 g, 2.61 mmol) in THF (5 mL) was added slowly over a period of 1 h. The mixture was stirred for 15 h, the solvents were removed under vacuum and the residue extracted with Et₂O (40 mL). The remaining solid was kept under vacuum overnight to remove residual ferrocene. Single crystals of Li(THF)₄[11] were grown by gas-phase diffusion of pentane into a THF solution of the crude product. Yield of Li(THF)₄[11]: 0.70 g (26%). ¹B{¹H} NMR (96.3 MHz, *d*₈-THF): δ -15.3 (*h*_{1/2} = 10 Hz). ¹H NMR (300.0 MHz, *d*₈-THF): δ 3.69 (s, 20H, C₅H₅), 3.87, 4.34 (2 × n.r., 2 × 8H, C₅H₄). ¹³C{¹H} NMR (75.5 MHz, *d*₈-THF): δ 66.0 (mult, C₅H₄), 68.1 (C₅H₅), 74.7 (mult, C₅H₄), n.o. (BC). Elemental analysis: Calcd. for C₅₆H₆₈BF₄LiO₄ (1046.25): C, 64.29; H, 6.55. Found: C, 63.88; H, 6.54%.

Crystal structure determinations of **3**, **6**, **7**, (Li(12-c-4)(THF))₂[10], Li(12-c-4)₂[10] and Li(THF)₄[11]

Single crystals of **3**, **6**, **7**, (Li(12-c-4)(THF))₂[10], Li(12-c-4)₂[10] and Li(THF)₄[11] were analyzed with a STOE IPDS II two-circle diffractometer with graphite-monochromated MoK_α radiation. Empirical absorption corrections were performed using the MULABS³⁶ option in PLATON.³⁷ The structures were solved by direct methods using the program SHELXS³⁸ and refined against *F*² with full-matrix least-squares techniques using the program SHELXL-97.³⁹ All non-hydrogen atoms (except disordered atoms in Li(12-c-4)₂) were refined with anisotropic displacement parameters. Hydrogen atoms were refined using a riding model. The crown ether molecules of (Li(12-c-4)(THF))₂[10] are disordered over two positions (occupancy factors 0.663(5) and 0.337(5)). Li(12-c-4)₂[10] contains one equivalent of non-coordinating THF in the crystal lattice. One of the two crown ether molecules of Li(12-c-4)₂[10] is disordered over two positions (occupancy factors

0.51(1) and 0.49(1)). The Flack-*x*-parameters for structures **3** and **6** are 0.00(2) and -0.02(3), respectively.

CCDC reference numbers: 699649 (**3**), 699647 (**6**), 699646 (**7**), 699650 ((Li(12-c-4)(THF))₂[10]), 699648 (Li(12-c-4)₂[10]) and 710783 (Li(THF)₄[11]).†

Electrochemical measurements. All electrochemical measurements were performed by using an EG&G Princeton Applied Research 263A potentiostat with glassy carbon or platinum disc working electrode. Carefully dried (CaH₂) and degassed CH₂Cl₂ was used as the solvent and [NBu₄][PF₆] as the supporting electrolyte (0.1 M). All potential values are referenced against the FcH/FcH⁺ couple. Spectroelectrochemical measurements were performed in a home-built optically transparent thin-layer electrolysis (ÖTTLE) cell following the design of Hartl *et al.*⁴⁰ with a Bruins Instruments Omega 20 UV/vis/NIR spectrometer.

Möbßbauer spectra. The details of ⁵⁷Fe temperature-dependent Möbßbauer spectroscopy have been described earlier.^{41–43} Due to the air- and moisture sensitivity of the compounds, sample transfer to perspex sample holders, lubricated with high-temperature silicone grease and sealed with O-rings, was effected in an inert-atmosphere glove box (VAC model DLX-001-S-P) having an oxygen partial pressure of less than 0.5 ppm and less than 1 ppm H₂O. The filled and sealed sample holders were removed from the glove box, immediately cooled to liquid nitrogen temperature, and then placed into the Möbßbauer spectrometer pre-cooled to ≈ 90 K. Data accumulation (in the first instance) was effected in a warming mode as discussed above. All isomer shifts are reported with respect to the centroid of a room temperature α-Fe absorber spectrum which was also used for spectrometer calibration.

Acknowledgements

M. W. is grateful to the "Deutsche Forschungsgemeinschaft" (DFG) and the "Fonds der Chemischen Industrie" (FCI) for financial support. L. K. wishes to thank the "Hessisches Ministerium für Wissenschaft und Kunst" for a Ph. D. grant. The authors are also indebted to A. Aharoni for effecting the careful sample transfers in an inert atmosphere glovebox to avoid oxidative degradation prior to spectral examination of the samples referred to herein.

References

- I. Manners, *Adv. Organomet. Chem.*, 1995, **37**, 131.
- P. Nguyen, P. Gómez-Elipe and I. Manners, *Chem. Rev.*, 1999, **99**, 1515.
- I. Manners, *Science*, 2001, **294**, 1664.
- I. Manners, *Synthetic Metal-Containing Polymers*, Wiley-VCH, Weinheim, 2004.
- D. A. Foucher, B.-Z. Tang and I. Manners, *J. Am. Chem. Soc.*, 1992, **114**, 6246.
- F. Jäkle, R. Rulkens, G. Zech, D. A. Foucher, A. J. Lough and I. Manners, *Chem. Eur. J.*, 1998, **4**, 2117.
- T. J. Peckham, J. A. Massey, C. H. Honeyman and I. Manners, *Macromolecules*, 1999, **32**, 2830.
- R. Rulkens, D. P. Gates, D. Balaishis, J. K. Pudelski, D. F. McIntosh, A. J. Lough and I. Manners, *J. Am. Chem. Soc.*, 1997, **119**, 10976.
- K. Ma, M. Scheibitz, S. Scholz and M. Wagner, *J. Organomet. Chem.*, 2002, **652**, 11.
- J. B. Heilmann, M. Scheibitz, Y. Qin, A. Sundaraman, F. Jäkle, T. Kretz, M. Bolte, H.-W. Lerner, M. C. Holthausen and M. Wagner, *Angew. Chem. Int. Ed.*, 2006, **45**, 920.

- 11 J. B. Heilmann, Y. Qin, F. Jäkle, H.-W. Lerner and M. Wagner, *Inorg. Chim. Acta*, 2006, **359**, 4802.
- 12 M. Scheibitz, R. F. Winter, M. Bolte, H.-W. Lerner and M. Wagner, *Angew. Chem. Int. Ed.*, 2003, **42**, 924.
- 13 M. Scheibitz, J. B. Heilmann, R. F. Winter, M. Bolte, J. W. Bats and M. Wagner, *Dalton Trans.*, 2005, 159.
- 14 M. Fontani, F. Peters, W. Scherer, W. Wachter, M. Wagner and P. Zanello, *Eur. J. Inorg. Chem.*, 1998, 1453.
- 15 M. Fontani, F. Peters, W. Scherer, W. Wachter, M. Wagner and P. Zanello, *Eur. J. Inorg. Chem.*, 1998, 2087.
- 16 M. Grosche, E. Herdtweck, F. Peters and M. Wagner, *Organometallics*, 1999, **18**, 4669.
- 17 R. E. Dinnebier, M. Wagner, F. Peters, K. Shankland and W. I. F. David, *Z. Anorg. Allg. Chem.*, 2000, **626**, 1400.
- 18 M. D. Thomson, M. Novosel, H. G. Roskos, T. Müller, M. Scheibitz, M. Wagner, F. Fabrizi de Biani and P. Zanello, *J. Phys. Chem. A*, 2004, **108**, 3281.
- 19 D. O. Cowan, P. Shu, F. L. Hedberg, M. Rossi and T. J. Kistenmacher, *J. Am. Chem. Soc.*, 1979, **101**, 1304.
- 20 B. Wrackmeyer, U. Dörfler, W. Milius and M. Herberhold, *Z. Naturforsch. B*, 1996, **51**, 851. Note that also the corresponding mesityl derivative FcBMes₂ has recently been published: A. E. J. Broomsgrove, D. A. Addy, C. Bresner, I. A. Fallis, A. L. Thompson and S. Aldridge, *Chem. Eur. J.*, 2008, **14**, 7525.
- 21 R. W. Fish and M. Rosenblum, *J. Org. Chem.*, 1965, **30**, 1253.
- 22 W. Haubold, J. Herdtle, W. Gollinger and W. Einholz, *J. Organomet. Chem.*, 1986, **315**, 1.
- 23 L. Kaufmann, H. Vitze, M. Bolte, H.-W. Lerner and M. Wagner, *Organometallics*, 2007, **26**, 1771.
- 24 M. D. Rausch and D. J. Ciappenelli, *J. Organomet. Chem.*, 1967, **10**, 127.
- 25 I. R. Butler, W. R. Cullen, J. Ni and S. J. Rettig, *Organometallics*, 1985, **4**, 2196.
- 26 H. Nöth and B. Wrackmeyer, Nuclear Magnetic Resonance Spectroscopy of Boron Compounds, in *NMR Basic Principles and Progress*, ed. P. Diehl, E. Fluck and R. Kosfeld, Springer, Berlin, 1978.
- 27 A. Appel, F. Jäkle, T. Priermeier, R. Schmid and M. Wagner, *Organometallics*, 1996, **15**, 1188.
- 28 M. Scheibitz, M. Bolte, J. W. Bats, H.-W. Lerner, I. Nowik, R. H. Herber, A. Krapp, M. Lein, M. C. Holthausen and M. Wagner, *Chem. Eur. J.*, 2005, **11**, 584.
- 29 B. E. Carpenter, W. E. Piers, M. Parvez, G. P. A. Yap and S. J. Rettig, *Can. J. Chem.*, 2001, **79**, 857.
- 30 B. Wrackmeyer, U. Dörfler, W. Milius and M. Herberhold, *Polyhedron*, 1995, **14**, 1425.
- 31 K. Venkatasubbaiah, I. Nowik, R. H. Herber and F. Jäkle, *Chem. Commun.*, 2007, 2154.
- 32 I. Nowik and R. H. Herber, *Eur. J. Inorg. Chem.*, 2006, 5069.
- 33 R. H. Herber, I. Nowik, D. A. Loginov, Z. A. Starikova and A. R. Kudinov, *Eur. J. Inorg. Chem.*, 2004, 3476.
- 34 G. E. Southard and M. D. Curtis, *Organometallics*, 2001, **20**, 508. For detailed studies on IVCT interactions in a 1,2-diborylated ferrocene dimer see: (a) K. Venkatasubbaiah, L. N. Zakharov, W. S. Kassel, A. L. Rheingold and F. Jäkle, *Angew. Chem., Int. Ed.*, 2005, **44**, 5428; (b) K. Venkatasubbaiah, A. Doshi, I. Navik, R. H. Herber, A. L. Rheingold and F. Jäkle, *Chem.-Eur. J.*, **14**, 444; (c) ref. 31.
- 35 F. Rebiere, O. Samuel and H. B. Kagan, *Tetrahedron Lett.*, 1990, **31**, 3121.
- 36 R. H. Blessing, *Acta Cryst.*, 1995, **A51**, 33.
- 37 A. L. Spek, *J. Appl. Cryst.*, 2003, **36**, 7.
- 38 G. M. Sheldrick, *Acta Cryst.*, 1990, **A46**, 467.
- 39 G. M. Sheldrick, *SHELXL-97. A Program for the Refinement of Crystal Structures*, Universität Göttingen, 1997.
- 40 M. Krejčík, M. Daněk and F. Hartl, *J. Electroanal. Chem.*, 1991, **317**, 179.
- 41 I. Nowik and R. H. Herber, *Inorg. Chim. Acta*, 2000, **310**, 191.
- 42 R. H. Herber and I. Nowik, *Hyperfine Interact.*, 2000, **126**, 127.
- 43 R. H. Herber and I. Nowik, *Hyperfine Interact.*, 2001, **136/137**, 699.