Principles of Intracellular Signaling in Ciliated Protozoa—A Brief Outline

Helmut Plattner

Abstract Ciliates have available most of the intracellular signaling mechanisms known from metazoans. Long-range signals are represented by firmly installed microtubules serving as gliding rails aiming at specific targets. Many components are distinctly arranged to guarantee locally restricted effects. Short-range signals include Ca^{2+} , provided from different sources, and proteins for membrane recognition and fusion, such as SNAREs, GTPases and high affinity Ca^{2+} -binding proteins (still to be defined). A battery of ion conductances serves for electric coupling from the outside medium to specific responses, notably ciliary activity, which also underlies gravitaxis responses. Eventually cyclic nucleotides are involved, e.g. in ciliary signaling. Furthermore, an elaborate system of protein kinases and phosphatases exerts signaling mechanisms in widely different processes.

2.1 Introduction—Basic Aspects of Signaling in Ciliates

As for every eukaryotic cell one may ask also for ciliates which cellular processes require signaling, how signaling is executed and over which distances, whether principles are shared with metazoans and plants, whether mechanisms are maintained during evolution, abolished or newly invented. Together with *Dictyostelium*, the ciliates *Paramecium* and *Tetrahymena* represent the protozoa which, at this time, are best analyzed with regard to signaling. It is useful to differentiate between long- and short-range signaling, e.g. by microtubules or electrical signals and by molecular interactions or spatially restricted Ca^{2+} signals (Plattner and Klauke 2001), respectively.

H. Plattner (✉)

Department of Biology, University of Konstanz, P.O. Box M625, 78457 Konstanz, Germany
e mail: helmut.plattner@uni-konstanz.de

2.1.1 Basic Phenomena Applicable to Ciliates

Signaling pertinent to ciliary activity in ciliated protozoa is as elaborate, or even more than in metazoan (Machemer 1988a) as these cells are highly mobile and capable of reacting to various environmental stimuli (Machemer 1988b; Bell et al. 2007). To achieve this, mechanical, electrical, biochemical and molecular signals, i.e. long range and short-range signals, can be combined in some variation to the basic theme.

Ciliates have at their disposal a highly sophisticated vesicle trafficking system, as illustrated (<http://www5.pbrc.hawaii.edu/allen/>) and summarized (Allen and Fok 2000) for *Paramecium* and *Tetrahymena* (Frankel 2000). The routes have to be addressed here. (i) Endoplasmic reticulum (ER) → Golgi apparatus → lysosomes + dense core-secretory organelles (trichocysts in *Paramecium* and mucocysts in *Tetrahymena*). (ii) Constitutive exocytosis of surface coat materials (Flötenmeyer et al. 1999) and dense core-secretory organelle exocytosis (Plattner et al. 1985; Plattner and Kissmehl 2003a). (iii) Phagocytosis, from cytostome → phagosome → endosomal and lysosomal input → phagolysosome (called “food [digesting] vacuole”) → discharge of spent vacuoles at the cytoproct. (iv) Endocytosis via early endosomes → links to phagosomes + lysosomes. (v) Vesicle recycling from the cytoproct to the nascent phagosome. (vi) In addition, the contractile vacuole complex impresses not only by its dynamic activity (Allen and Naitoh 2002) in the context of ongoing osmoregulation (Allen et al. 2009), but it also represents a site endowed with the machinery typical of vesicle trafficking (Plattner 2015b) although vesicle trafficking within the organelle is less obvious. Steps (iii) to (v) have been documented in detail for *Paramecium* (Allen and Fok 2000) as well as for *Tetrahymena* (Frankel 2000). Beyond short-range signaling, steps (i), (iii) and (v) include long-range signaling. All these pathways serve for proper delivery and positioning of signaling elements so that they can execute their signaling function at distinct sites of the cell.

2.1.2 Molecular Key Players

Recent availability of a macronuclear genome database for the most frequently used species, *P. tetraurelia* and *T. thermophila*, has enabled the identification, localization and assessment of the functional relevance of key players. In *Paramecium* such work has included mainly SNARE (soluble *N*-ethylmaleimide sensitive factor [NSF] attachment protein receptors) proteins, actin and H⁺-ATPase, as summarized previously (Plattner 2010) as well as Ca²⁺-release channels (CRC) of the type inositol 1,4,5-trisphosphate receptors (InsP₃R) and ryanodine receptor-like proteins (RyR-LP) (Ladenburger and Plattner 2011; Ladenburger et al. 2006, 2009; Plattner 2015a), as summarized recently (Plattner and Verkhatsky 2015). This is complemented by monomeric GTP (guanosine trisphosphate) binding proteins

(G-proteins), the GTPases, not only in higher eukaryotes (Zhen and Stenmark 2015) but also in ciliates (Bright et al. 2010). Isoforms, i.e. paralogs or ohnologs (in case of diversification following whole genome duplications, particularly in *P. tetraurelia*), can be assigned to different steps and routes of vesicle trafficking and, thus, mirror the high complexity of the ciliate cell.

2.1.3 Long- and Short-Range Signals

The distinction between short-range and long-range signals has been extensively elaborated elsewhere (Plattner 2016a). A typical long-range signal is the docking of trichocysts (Aufderheide 1978) along microtubules which emanate from ciliary basal bodies and, thus, serve as transport rails (Plattner et al. 1982). This has to be complemented by short-range signals. For instance SNAREs and G-proteins are important for vesicle docking and finally membrane fusion. Local Ca^{2+} increase is another signal which has to arise from a nearby source since Ca^{2+} signals decay rapidly (Neher 1998). This also guarantees selective activation of distinct sites and also avoids cytotoxicity (Plattner and Verkhatsky 2015). Local restriction of Ca^{2+} signaling is most obvious, for instance, by the assignment of different CRCs types to different trafficking organelles, from the cell surface to deep inside, in *Paramecium* (Plattner 2015a). Moreover, ciliates fascinate particularly by their highly regular design that predetermines their vesicle trafficking routes and signaling sites based on epigenetic phenomena (Frankel 2000; Beisson 2008). Accordingly, cilia and secretory organelles are arranged in a strikingly regular surface pattern.

2.2 Overview of Trafficking Regulation Along Different Signaling Pathways

Basic trafficking pathways in ciliates are outlined in Fig. 2.1a. Box 1 outlines different kinds of cytoplasmic signaling operating in ciliates. Despite the old evolutionary age of ciliates, signaling mechanisms are quite similar to those in animals and—with exceptions—in plants. Similarities encompass the role of monomeric GTP-binding proteins (G-proteins acting as GTPases) (Bright et al. 2010), H^+ -ATPase, SNARE proteins and their chaperone, NSF, as well as the regulation of membrane fusion by a local Ca^{2+} signal (Plattner 2010). The importance of luminal acidification of trafficking vesicles is derived from the observation that a trans-membrane signal generated by the conformational change of H^+ -ATPase intramembranous V0 part causes binding of GTPase modulators (Hurtado-Lorenzo et al. 2006), thus facilitating docking and membrane fusion. Specificity of vesicle interaction is finally mediated by SNAREs (Plattner 2010) and GTPases (Bright et al. 2010). Sequences encoding GTPases and GTPase modulators, such as GAP

◀ **Fig. 2.1** Signaling pathways in the *Paramecium* cell. **a** Vesicle trafficking pathways encompass different main streams, such as the exocytotic, the endocytotic, the phagocytotic pathway and less overt trafficking in the contractile vacuole complex. *Dotted arrows* are less well established, particularly membrane input into this organelle via acidosomes, as derived from various recent papers about other protists. Also for proteins passing or bypassing the Golgi apparatus has not yet been sufficiently specified in detail. **b** Cortical organelles, such as cilia and exocytosis sites are regulated separately. Depolarization induces ciliary beat reversal by Ca^{2+} influx via ciliary voltage dependent Ca^{2+} channels, abolished via negative feedback ($\ominus$) by intraciliary $[\text{Ca}^{2+}]$ increase. CRCs in alveolar sacs, type RyR LPs, are facing the plasmamembrane, opposite to the SERCA pump. Alveolar sacs contain a calsequestrin like high capacity/low affinity CaBP. Trichocyst exocytosis is governed by a SOCE mechanism (store operated Ca^{2+} entry), i.e. Ca^{2+} release from alveolar sacs in a first step, followed by Ca^{2+} influx via somatic (non ciliary) channels in a tightly coupled second step. **c** Summary of events during trichocyst exocytosis. *Top* Freeze fracture images of fusion/resealing stages and their estimated duration, derived from synchronous stimulation/quenched flow/rapid freezing analysis. Note decay of rosette particle aggregates and rapid formation of a fusion pore which expands and, thus, allows Ca^{2+} access to the secretory contents which triggers their explosive discharge by descondensation (stretching). *Below* Parallel situations seen on ultrathin sections. **a** Data pertinent to trichocyst processing are based on previous reviews (Plattner et al. 1993; Plattner 2014), those for endo /phagocytotic trafficking are mainly derived from Allen and Fok (2000) and **c** trafficking in context of the contractile vacuole complex is based on recent reviews (Plattner 2015b, 2016a) **b** is modified from Plattner (2014), **c** is modified from Plattner et al. (1993, 1997)

(guanine nucleotide activation protein) and GEF (guanine nucleotide exchange factor), also occur in the *P. tetraurelia* database (Plattner and Kissmehl 2003b).

The Ca^{2+} signal is generated by intracellular CRCs of which different types are assigned to different organelles (Ladenburger and Plattner 2011; Plattner and Verkhratsky 2013). The Ca^{2+} -sensor causing fusion, as known from higher eukaryotes, is a low capacity/high affinity Ca^{2+} -binding protein (CaBP) which usually contains two high affinity Ca^{2+} -binding C2 domains (β -barrels with a Ca^{2+} -binding loop), such as synaptotagmin (Rizo et al. 2006; Südhof 2014). Although such CaBPs have not yet been specified in ciliates, equivalents of synaptotagmin occur in the *P. tetraurelia* database (Farrell et al. 2012). Extended synaptotagmins (e-syntag) with more than two C2 domains are known from some mammalian cells (Min et al. 2007), but they also occur in the *Paramecium* database (H. Plattner and R. Kissmehl, unpublished observations). Calmodulin (CaM) is another low capacity/high affinity CaBP, with four EF-hand type loops, each with high affinity Ca^{2+} -binding capacity. CaM operates at many sites also in ciliates. In the CaM molecule, the extensive conformational change upon hierarchical Ca^{2+} binding in the EF-hand loops I to IV represents the transduction of a chemical to a molecular-mechanical signal (Park et al. 2008). Thus, CaM can regulate a variety of surface influx channels (Saimi and Kung 2002), phagocytosis (Gonda et al. 2000) and probably endocytosis, also in ciliates.

Box 1 also shows that Ca^{2+} for activation may eventually also come from the outside medium for some specific effects, e.g. for activating some nucleotide cyclases, kinases and phosphatases, in the context of ciliary activity. This includes the signaling function of cyclic nucleotides, such as cyclic adenosine monophosphate (cAMP) and cyclic guanosine monophosphate (cGMP) and activation of the

Box 1. Kinds of cytoplasmic signals operating in ciliates – a survey

Electrical signals: depolarization, hyperpolarization → change of ciliary beat

Acidification of organelle lumen

H^+ -ATPase/pump → binding of GTPases + their modulators → membrane docking and fusion by interference of SNARE proteins (SNARE = soluble N-ethylmaleimide sensitive factor [NSF] attachment protein receptors) and low capacity/high affinity Ca^{2+} -binding proteins (CaBPs) with C2-domains

Ca^{2+} , from outside or from intracellular stores (Ca^{2+}_o , Ca^{2+}_i)

→ activation of some nucleotide cyclases, some protein kinases and phosphatases, of CaBPs and of membrane fusion and fission (synaptotagmin with two C2 domains, extended synaptotagmin with more C2 domains): to be settled for ciliates

Nucleotides

cyclic adenosine monophosphate (cAMP) → protein kinase A (PKA)

cyclic guanosine monophosphate (cGMP) → protein kinase G (PKG)

→ protein phosphorylation, also in ciliates

cyclic adenosinediphosphoribose (cADPR) from NAD (nicotinamide adeninedinucleotide) → activation of some (ryanodine receptor-type) Ca^{2+} -release channels (CRCs) in mammalian cells: activators effective in ciliates, but not yet assigned to specific channels

nicotinic acid adenine dinucleotidephosphate (NAADP), formed from NADP,

→ activation of two-pore channels, in acidic compartments: also in ciliates?

Protein phosphorylation

PKA, PKG: as defined above

Ca^{2+} -dependent protein kinases (CDPKs) with integrated calmodulin- (CaM)-like motifs

Protein dephosphorylation by protein phosphatases (PPs) type PP1, PP2A, PP2B

PP2B (calcineurin)

with subunit (SU) A (catalytic SU, containing SU-B- and CaM-binding domain)

and SU-B (regulatory SU, with Ca^{2+} -binding domain) also in ciliates

respective protein kinases, protein kinase A (PKA) and protein kinase G (PKG) (Bonini and Nelson 1990); for review see Plattner (2016a). Also some metabolic Ca^{2+} channel activators, such as cyclic adenosinediphosphoribose (cADPR) and nicotinic acid adenine dinucleotidephosphate (NAADP) are derived

from nucleotides, i.e. nicotinamide adeninedinucleotide (NAD) and nicotinic acid adenine dinucleotidephosphate (NAADP), respectively, as known from vertebrates (Lee 2012). For cADPR and NAADP effects there is only circumstantial evidence in *Paramecium* (Plattner et al. 2012).

A total of 2600 kinases has been found in the *P. tetraurelia* genome (Bemm et al. 2009), thus contributing by 7 % to the macronuclear genome. In *T. thermophila* the proportion is 3.8 % (Tian et al. 2014). Both values stress their importance for signal transduction. The difference between the two genera may originate from whole genome duplication in *Paramecium*. A considerable difference between protein kinases in animal cells and in ciliates is the absence in the latter of a “CaM kinase”, i.e. a kinase activated by a complex of calmodulin (CaM) and Ca^{2+} . Whereas such CaM-kinases in metazoans contribute to the regulation of neuronal activity, they are replaced in ciliates by “ Ca^{2+} -dependent protein kinases” (CDPKs). These contain CaM-like sequences integrated in the kinase molecule (Kim et al. 1998).

Box 1 also indicates the occurrence in ciliates of protein phosphatases (PPs), e.g. PP1, PP2A and PP2B. PP2B, which is identical with calcineurin, encompasses two subunits, catalytical subunit A and regulatory subunit B, from ciliates (Fraga et al. 2010) to man where it regulates immune-response and long term potentiation, i.e. learning. In ciliates, multiple roles can be expected for calcineurin, including exo-/endocytosis regulation (Momayezi et al. 1987; Fraga et al. 2010).

2.3 Subcompartmentalization of Signaling Including Signaling in Cilia

Signals can be rather precisely restricted to subcompartments, e.g. cilia (Box 2), for which Box 3 shows details. Mechanical stimulation of a ciliated protozoan cell causes depolarization or hyperpolarization, depending on whether stimulation occurs at the anterior or posterior part of the cell (Eckert and Brehm 1979; Machemer 1988a, b). This is enabled by a graded differential distribution of specific ion channels over the somatic (non-ciliary) cell membrane. The respective receptor potential formed by different ion conductances activates different mechanisms in cilia. For instance, depolarization activates voltage-dependent Ca^{2+} -channels selectively occurring in cilia (Machemer and Ogura 1979) and, thus, a Ca^{2+} -carried action potential. (This signaling occurs no more in metazoans beyond Ctenophores.) Increased intraciliary Ca^{2+} shuts off this Ca^{2+} influx (Brehm and Eckert 1978). Hyperpolarization accelerates forward swimming (Preston et al. 1992).

During de- and hyperpolarization, different cyclic nucleotides are formed, activating PKG and PKA, respectively (Box 3). Ciliary activation mechanisms are independent of Ca^{2+} -activated processes during exocytosis, except when massive exocytosis stimulation entails an exuberant Ca^{2+} signal (Husser et al. 2004). In

Box 2. Examples of subcompartmentalization of signals in ciliates

External mechanical stimulation

→ Ca^{2+} signaling in cortical domains, e.g. for receptor potential formation
 anterior stimulation → depolarization/action potential → ciliary reversal
 posterior stimulation → hyperpolarization → accelerated forward swimming

for details of ciliary activity, see Box 3

Activation of non-ciliary (somatic) membrane phenomena for exocytosis: see Box 5

Constitutive local intracellular Ca^{2+} signaling via organelle-specific Ca^{2+} -release channels (CRCs) in all trafficking vesicles

Organelle specific protein phosphorylation processes for other activation mechanisms: see Box 8

Box 3. Ciliary beat activity in ciliates

(A) Anterior mechanical stimulus (hitting an obstacle)

somatic cell membrane → receptor potential: depolarization by activation of anterior mechanosensitive Ca^{2+} -channels, repolarization by K^{+} -efflux

Effect of depolarization on cilia: action potential by

activation of voltage-dependent (ΔV) Ca^{2+} influx channels in ciliary membrane
 → guanylate cyclase activation → cGMP formation → PKG activation
 → phosphorylation of target proteins in ciliary axoneme;
 in parallel: activation of axonemal calmodulin (CaM) → different effects

No Ca^{2+} spillover into cell soma

Inactivation of ciliary reversal by closing ciliary ΔV -channels by a Ca^{2+} /CaM complex and binding of excessive Ca^{2+} to immobile buffer (CaM, centrin)

(B) Posterior mechanical stimulus

Hyperpolarization by K^{+} efflux (somatic cell membrane)
 → hyperpolarization-activated Ca^{2+} -channels (somatic cell membrane),
 adenylate cyclase activation → cAMP formation → PKA activation →
 phosphorylation of target proteins in ciliary axoneme

(C) Gravikinesis/gravitaxis

Positive gravitaxis: “statocyst”-mediated intracellular signal perception (*Loxodes*)

Negative gravitaxis: very much predominant form of gravitaxis (*Paramecium*, *Tetrahymena*)

activation by hyperpolarization via posteriorly enriched K^{+} -channels; postulated link to cortical F-actin → upward movement

summary, a mechanical signal is transformed into a long-range electrical signal generated at the somatic cell membrane that is transduced into cilia where it causes short-range Ca^{2+} signaling and a mechanical response in ciliary activity.

Mechanisms described for basic ciliary activity (Fig. 2.1b) also apply to chemotaxis and to gravitaxis/gravikinesis (Box 3). Chemotaxis requires the activation of distinct ion conductances to achieve specific behavioral responses (Saimi and Kung 2002; Bell et al. 2007; Yano et al. 2015). Positive gravitaxis is rare in ciliates where negative gravitaxis, causing upward swimming in the gravity field, by far prevails. For this, *Paramecium* is the best analyzed example (Machemer et al. 1998; Hemmersbach and Braun 2006; Machemer 2014). Accordingly cAMP and PKA are assumed to be involved in negative gravitaxis (Hemmersbach et al. 2002). Investigators assume that, for sensing gravity, channels have to be linked to filamentous actin (F-actin) (Machemer 2014). In fact, actin has been localized to the cell cortex (Sehring et al. 2007) and, even more precisely, to the narrow space between cell membrane and alveolar sacs (Kissmehl et al. 2004).

2.4 Organelle Trafficking Signals

2.4.1 Molecular Background

Long-range signals, such as firmly installed microtubules, can guide vesicles to their target sites (Box 4). This is true of trichocysts (Aufderheide 1978; Plattner et al. 1982) and organelles of the phagocytotic cycle (Allen and Fok 2000). Short-range signals involved are GTPases, SNAREs, H^+ -ATPase, as outlined in Sect. 2.1, together with actin. For GTPases (Bright et al. 2010) and the other key players, organelle specific isoforms are available (Plattner 2010). The multimeric H^+ -ATPase molecule is composed of an intramembranous V0 basepiece and a catalytic head, V1, which may dis- and re-assemble by interaction with an elongate, variable a-SU (Sun-Wada and Wada 2015). Considering the key role of H^+ -ATPase (Sect. 2.2), the unsurpassed number of 17 a-subunits in *Paramecium* may mediate adjustment to local requirements (Wassmer et al. 2005, 2006, 2009). Among SNAREs, longin-type sequences in *Paramecium*'s "synaptobrevins" may contribute to organelle specificity, in addition to the usual domain sequences (Schilde et al. 2006, 2010). In *P. tetraurelia*, plasmalemmal Syntaxin 1 (PtSyx1) is engaged in trichocyst exocytosis (Kissmehl et al. 2007). For more details, see Plattner (2010, 2016a).

Vesicles undergoing trafficking are endowed with CRCs identical with, or related to InsP_3 Rs and RyRs (Ladenburger and Plattner 2011; Plattner and Verkhatsky 2013); see Box 5. An exception are trichocysts which seem to be devoid of luminal Ca^{2+} , in contrast to what is known from some other dense core-secretory vesicles, endosomes and phagocytotic organelles of higher eukaryotes (Hay 2007). The presence of the key players mentioned above, including

Box 4. Organellar trafficking signals

Long-range signals

microtubular “rails” as firmly established guidelines (emanating from oral cavity [for phagocytosis] and from ciliary basal bodies [for trichocyst docking], respectively)

Short-range signals/molecular recognition sites

cooperative role of H^+ -ATPase molecules (acidification of organelle lumen) → binding of organelle-specific small GTPases (+ modifying proteins, e.g. GAP = guanosine nucleotide activation protein, GEF = guanosine nucleotide exchange factor, as known from higher eukaryotes)

vesicle docking: SNAREs + GTPases

→ docking at target organelle/membrane: organelle-specific GTPase-binding proteins yet to be identified

local Ca^{2+} signal and Ca^{2+} sensor activation → membrane fusion

Vesicle budding: coatamer proteins (COPs), clathrin, adaptor proteins

Golgi apparatus: ill-defined molecular cues and signals in ciliates awaiting scrutiny

CRCs, in the endo-/phagocytotic cycle of *Paramecium* may reflect the intensity and multitude of vesicle trafficking known from ultrastructural studies (Allen and Fok 2000) [In *Paramecium*, not all of these vesicles are acidic (Wassmer et al. 2009), and not all lysosomal enzymes have an acidic pH-optimum (Fok and Paeste 1982; Fok 1983)]. Appropriate CRCs may drive membrane interactions in concert with, or independently from other key players. The importance of local availability and regulation of Ca^{2+} during membrane docking and fusion is discussed in the accompanying paper (Plattner 2016b). The numerous members of the six CRC subfamilies found in *Paramecium* may fine tune Ca^{2+} signals and membrane interactions depending on local requirements.

2.4.2 Dense Core-Secretory Vesicle Exocytosis

Ca^{2+} regulation of trichocyst exocytosis involves three steps (Box 5, Fig. 2.1b, c): (i) Ca^{2+} release from alveolar sacs via RyR-like proteins and (ii) immediately superimposed Ca^{2+} influx from the outside medium (Klauke and Plattner 1997; Ladenburger and Plattner 2011; Plattner 2014). Both mechanisms acting in concert are called store-operated Ca^{2+} entry, SOCE—a mechanism maintained up to mammals. A large excess of Ca^{2+} , much more than seen by fluorochromes, has to flood trichocyst exocytosis sites to become activated, just as in some

Box 5. Why a multiplicity of Ca^{2+} -release channels?

Paramecium contains 34 genes for Ca^{2+} -release channels (CRCs, 6 subfamilies), to be assigned to the superfamily of InsP_3R /Ryanodine receptor (RyR) type CRCs, distributed over different trafficking organelles for local signaling

Example A - RyR-like channels: alveolar sacs (cortical Ca^{2+} stores)

activation by RyR activators caffeine, 4-chloro-m-cresol or by polyamines (AED)

→ Ca^{2+} release from alveolar sacs by RyR-LP of CRC-IV subfamily, superimposed by Ca^{2+} -entry from the outside medium (store-operated Ca^{2+} -entry, SOCE) → trichocyst exocytosis based on

(i) membrane fusion (depending on $[\text{Ca}^{2+}]_i$ increase by SOCE) and →

(ii) Ca^{2+}_o access to trichocyst contents for inducing decondensation

(vigorous ejection)

membrane resealing and ghost detachment (exocytosis-coupled endocytosis), also driven by $[\text{Ca}^{2+}]_i$ increase

Example B - InsP_3Rs : occurring in the contractile vacuole complex; serving

(i) for fine-tuning of $[\text{Ca}^{2+}]$ in the cytosol and

(ii) for internal membrane restructuring (hypothetic) during contraction cycles

InsP_3R /RyR- type channels also include mixed types, in compartments undergoing trafficking

Additional Ca^{2+} -release channels in ciliates

two-pore channels NAADP-activated, in acidic stores? Occurrence likely

TRP-type and mechanosensitive channels: also not yet specified in ciliates

neuroendocrine cells (for details, see Plattner 2016a). (iii) Discharge of contents follows formation of an exocytotic opening and requires the entry of Ca^{2+} from the outside and binding to some secretory components, thus causing decondensation by conformational change (Plattner et al. 1997; Klauke et al. 1998; Plattner 2014). This in turn depends on proper processing of secretory protein precursors (Pouphile et al. 1986; Bowman et al. 2005).

2.4.3 The Phagocytotic Cycle

This aspect is reviewed here in more detail, as it demonstrates the complex sequence of interacting signaling molecules although these are only partially known.

The phagocytotic cycle in *Paramecium* requires multiple signaling (Allen and Fok 2000), including firmly established microtubules as long-range signals and variable stage-dependent short-range signals. In detail the sequence is as follows. (i) At the cytopharynx, at the bottom of the cytostome, vesicles recycling from advanced stages of food vacuoles, together with vesicles from the cytoproct, deliver membrane material for a bulging nascent food vacuole. Thus, a phagosome is formed at converging microtubular rails, the “postoral fibers”. (ii) After detachment, acidosomes (late endosomes) fuse with the phagosome, thus endowing it with H⁺-ATPase molecules for luminal acidification. (iii) This is followed by fusion with lysosomes, thus forming phagolysosomes. (iv) Lysosomal enzymes are retrieved later on during cyclosis, (v) as are parts of the membrane for delivery to the cytopharynx. (vi) The contents of spent food vacuoles are released by exocytosis at the cytoproct and membranes are recycled as indicated for step (i) (Allen and Fok 2000).

In *Paramecium tetraurelia*, key players for signaling in the different stages (Box 6) encompass exchanging sets of SNAREs (Schilde et al. 2006, 2010; Kissmehl et al. 2007), subunits (SU) of H⁺-ATPase (Wassmer et al. 2005, 2006), and actin, as outlined in a separate chapter (Plattner 2016b). In *Tetrahymena*, different types of GTPases are exchanged during cyclosis (Bright et al. 2010). In *Paramecium*, the exchange of numerous actin isoforms, types 1, 3, 6, 8, 11 14 and 16 as well as their patchy or unilateral arrangement in some stages is a most striking phenomenon (Sehring et al. 2007). This may serve propulsion of the organelle and/or regulation of accessibility to fusion and/or budding of vesicles. All this documents a series of interacting long- and short-range signaling during cyclosis.

Box 6. The phagocytotic cycle in ciliates

At cytopharynx

cell membrane enlargement by fusion of recycling vesicles → association with actin
→ nascent food vacuole pinches off to form a phagosome

During cyclosis

→ fusion with acidosomes (late endosomes) providing H⁺-ATPase → luminal acidification → fusion with lysosomes to form a phagolysosome (mature food vacuole)
endowment with varying SNAREs, small GTPases, and
actin coats (for details see text), whereas Ca²⁺-release channels are throughout
of type CRC-III (InsP₃R-type)
selective membrane input and retrieval, contents digestion
pH gradually increasing to ~7

At cytoproct

contents discharge by exocytosis and formation of recycling (“discoidal”) vesicles

2.4.4 The Contractile Vacuole Complex

Surprisingly, the contractile vacuole complex contains all components relevant for vesicle trafficking, except actin, in even higher variability and with strict localization to specific substructures, such as the vacuole, the pore and the meshwork of the smooth spongiome (Box 7). The organelle has a very complex design (Allen and Naitoh 2002). It not only can expel fluid with an excess of Ca^{2+} and other ions (Stock et al. 2002), but it also shows some reflux of Ca^{2+} into the cytosol via constitutively active InsP_3Rs (Ladenburger et al. 2006). This may serve not only for fine tuning of cytosolic Ca^{2+} but also to drive the extensive membrane fusion and fission processes within the organelle during systole/diastole cycles (Plattner 2015b).

Box 7. Signaling in the contractile vacuole complex

Signals assumed for self-assembly of new anlagen: centrin, CaM, γ -tubulin, NIMA kinase, as discussed elsewhere (Plattner 2015b)

Signals assumed for organelle growth: delivery of vesicle with specific v-/R- and t-/Q SNAREs

Local tubularization (spongiome) and reversible planar-tubular transitions: F-BAR proteins (hypothetic), as discussed elsewhere (Plattner 2015b)

Acidification by V-type H^+ -ATPase

ΔH^+ exploited for Ca^{2+} sequestration (hypothetic $\text{X}^+/\text{H}^+/\text{Ca}^{2+}$ exchanger, in addition to Ca^{2+} -ATPase (see Plattner 2016a) and expulsion of H_2O , Ca^{2+} and other ions by periodic exocytotic release at the pore

CRCs of type InsP_3R for constitutive partial Ca^{2+} reflux into cytosol: for $[\text{Ca}^{2+}]_i$ fine tuning and probably for spongiome restructuring

Pore for periodic contents release by exocytosis: with specific SNAREs and CRCs at the pore
periodic signal for vacuole contents release: mechanosensitive channels (suggested by occurrence of stomatin [Reuter et al. 2013] and in agreement with other systems [Plattner 2015b]), in conjunction with pore-specific SNAREs and CRCs

2.4.5 Additional Signals

Little is known about other types of Ca^{2+} release channels in ciliates, such as two pore-channels (TPC) and transient receptor potential-channels (TRPC) and their activators (Box 5). Particularly metabolic CRC activators (Lee 2012), such as cADPR, NAADP, remain to be assigned to different channels and organelles in

ciliates. Such channels have to be expected also in ciliates, based on microinjection studies (Plattner et al. 2012).

Vesicle budding at the Golgi apparatus and other organelles as well as at the plasmamembrane requires a set of additional proteins, such as coatamer proteins (COPs) and clathrin, together with their adaptor proteins known from higher eukaryotes up to mammals (Rothman 2014). In ciliate cells, coatamer coats are suggested to occur by electron microscopy in the cis- and trans-side of the Golgi apparatus (Allen and Fok 1993; Garreau De Loubresse 1993) and clathrin coats in addition by molecular biology according to Elde et al. (2005) who also reported the expression of adaptor proteins, AP-1, AP-2, AP-3 and AP-4 in *T. thermophila*. While none of them appear important for lysosome biogenesis (Briguglio et al. 2013), AP-2 is important for endocytosis via coated pits (Elde et al. 2005). Sequences encoding all these adaptor proteins have also been found in the *P. tetraurelia* database, in addition to the ARF/SAR-type G-protein known as a target of the drug, brefeldin A (Plattner and Kissmehl 2003b). The same is true of clathrin heavy chains and of COPs.

In summary, for vesicle trafficking ciliates have at their disposal most of the signaling components known from multicellular organisms. Note, however, that $\text{InsP}_3\text{R/RyR}$ -like molecules are absent from higher plants (Plattner and Verkhatsky 2015), whereas they occur in some green algae (Wheeler and Brownlee 2008). Globally a ciliate's signaling machinery closely resembles that of metazoans.

2.5 Protein Phosphorylation for Activation and Deactivation of Signaling Processes

2.5.1 Phosphorylation Processes

As mentioned in Sect. 2.3, signaling in cilia includes PKA and PKG activity for enhanced forward and backward swimming, respectively (Kim et al. 1998; Kutomi et al. 2012). Activating cyclic nucleotides are generated within one ciliary stroke (Yang et al. 1997). Together with CDPKs they belong to the superfamily of Seryl/Threonyl kinases (Box 8). Phosphoproteins are substrates of the different phosphatases. Among them, PP1 dephosphorylates a ciliary phosphoprotein formed during ciliary reversal in *Paramecium* (Klumpp et al. 1990). PP2B/calcineurin probably has a broad spectrum of activity, depending on its A-subunit, whereas the two genes for the B-SU in *Paramecium* result in an identical translation product, with a well conserved binding domain in the A-SU (Fraga et al. 2010).

As indicated in Box 8 and discussed in more detail somewhere else (Plattner 2016a), the occurrence of Tyrosyl phosphorylation may be largely restricted in ciliates to cell cycle and mitosis regulation. Work with mammalian cells exposed to

Box 8. Protein phosphorylation for signaling and activation processes in ciliates

(A) Seryl/Threonyl phosphorylation

Protein kinases (PK)

protein kinase A, PKA (cAMP-activated)

protein kinase G, PKG (cGMP-activated)

CDPK (Ca²⁺-dependent protein kinase, with a CaM-like domain) substituting for CaM-kinase (activated by a separate Ca²⁺/CaM-complex) in animal cells

Protein phosphatases (PP): PP1, PP2A, PP2B (Ca²⁺/CaM-dependent PP = calcineurin)

(B) Dedicated Tyrosyl phosphorylation

Some predicted for ciliates from proteomic analysis (still to be confirmed)

mainly concerning cell cycle and mitosis regulation (MAPKs = mitogen-activated protein kinases)

new aspects emerging from phosphoproteomic analysis

Euplotes gamones indicates signaling via a mitogen-activated protein kinase (MAPK) cascade with Tyrosine phosphorylation (Vallesi et al. 2010; Cervia et al. 2013). See also chapter by Luporini.

2.5.2 Signal Downregulation

Also ciliates possess different ways to downregulate signals (Box 9). Cyclic nucleotides are deactivated by diesterases and phosphoproteins are dephosphorylated by protein phosphatases. For instance, the association of calcineurin with parasomal sacs (Momayezi et al. 2000), the clathrin-coated pits in ciliates, is compatible with dynamin dephosphorylation known from mammalian coated pits.

Ca²⁺ signals are downregulated by different mechanisms with different kinetics (Box 9). The most rapid is binding to centrin (Sehring et al. 2009) a CaBP with high capacity/low affinity (in addition to low capacity/high affinity) binding sites localized in the cell cortex of *Paramecium* (see Plattner 2016a). This is orders of magnitude more rapid than downregulation by Ca²⁺-ATPases/pumps (Plattner 2016a) of which type Sarcoplasmic/Endoplasmic Reticulum Ca²⁺-ATPase (SERCA) (Hauser et al. 1998) or plasmamembrane Ca²⁺-ATPase (PMCA) (Elwess and Van Houten 1997) have been analyzed in *Paramecium*. PMCA also occurs in cilia of *Tetrahymena* (Dentler 1988) and *Paramecium* (Yano et al. 2013). These two are P-type ATPases because they autocatalytically form a phospho-intermediate which then dephosphorylates itself. Ca²⁺ exchangers, though not yet identified, show up in ciliate databases; they are driven by a H⁺-gradient formed by a H⁺-ATPase (V-type, in vesicles) operating without a phospho-intermediate formation. Although such

Box 9. Shut-down of signalling in ciliatesInactivation of Ca^{2+}

- binding to high capacity/low affinity Ca^{2+} -binding proteins, e.g. centrin

- reduction by pumps and transporters

- Ca^{2+} extrusion and sequestration by Ca^{2+} -ATPases/pumps

- PMCA (plasmamembrane Ca^{2+} -ATPase)

- SERCA (sarcoplasmic/endoplasmic reticulum Ca^{2+} -ATPase)

- hypothetical: $\text{X}^+/\text{Ca}^{2+}$ exchangers, e.g. $\text{H}^+/\text{Ca}^{2+}$ antiporter (postulated specifically for contractile vacuole complex)

Inactivation of cyclic nucleotides by diesterases

Reversion of phosphorylation state

- protein phosphatase PP1, possibly also PP2C, for deactivation of ciliary reversal

- PP2B/calcineurin: pleiotropic effects to be expected, e.g. dynamin activation for vesicle budding and regulation of Ca^{2+} stores, e.g. by effects on CRCs

exchangers urgently call for scrutiny in ciliates it appears that they are much more efficient in signal downregulation than the pumps (Ladenburger et al. 2006; Plattner 2016a).

2.6 Signaling by Surface Receptors

These aspects are summarized in Box 10. The occurrence of trimeric GTP-binding proteins (G-proteins) is likely (De Ondarza et al. 2003; Lampert et al. 2011), but not firmly established in protozoa in general (Krishnan et al. 2015) and in ciliates in particular since important details have not been examined yet, as discussed in more detail elsewhere (Plattner 2016a). The same is true of G-protein-coupled receptors (GPCRs). All this also applies to the secretagogue, aminoethyldextran, which, in *Paramecium*, is most efficient in activating highly synchronous exocytosis (Plattner et al. 1985; Plattner and Kissmehl 2003; Knoll et al. 1991) by a SOCE mechanism for trichocyst exocytosis (Hardt and Plattner 2000; Plattner 2014). For hints to MAPK activity and Tyrosyl phosphorylation, see Sect. 2.5.

Purinergic receptors can be assumed to occur in *Paramecium* as these cells, upon exposure to $\geq 10 \mu\text{M}$ GTP, perform periodic back- and forward swimming accompanied by depolarization (Clark et al. 1993) and Ca^{2+} waves oscillating with the same periodicity (Sehring and Plattner 2004). This is unusual insofar as purinergic receptors normally respond to ATP or, less common, to UTP. We assume a function in keeping cells from dispersal to low density which is known to inhibit cell division and maintenance of the population.

Box 10. Surface receptor signalling in ciliates

Trimeric GTP-binding proteins (G-proteins) and G-protein-coupled receptors (GPCRs) existence in ciliates under considerable debate, fragmentary information

Mitogen-activated protein kinase (MAP kinase; MAPK): related activities are currently assumed also for ciliates

Effects of exogenous GTP ($\geq 10 \mu\text{M}$): Ca^{2+} oscillations in parallel to de-/repolarizations

the first $[\text{Ca}^{2+}]_i$ peak (larger than subsequent periodic peaks) requires Ca^{2+} .

→ further (smaller) cyclic activity peaks in $\sim 8 \text{ s}$ oscillations supported by Ca^{2+} from internal stores (type of store for internal Ca^{2+} mobilization during GTP activation: unknown) → ongoing periodic activation → desensitization and signal downregulation (mechanism unknown) → decaying Ca^{2+} signal

Unknown: purinergic receptors, Ca^{2+} /polyvalent cation-sensing receptor (a GPCR?)

Chemotaxis chemoreceptors

Operating via specific ion conductances (see text)

2.7 Conclusions

Intracellular signaling by pheromones (gamones) in ciliates (Luporini et al. 2014) is summarized separately in this volume. Epigenetic signaling is also covered separately in this volume by Nowacki; for surveys, also see Chalker et al. (2013) and Simon and Plattner (2014). Most of the other signaling mechanisms described here seem to be evolutionarily old and maintained from protozoa on, particularly ciliates, up to top-ranking metazoans. The impressive complexity of ciliate cells and their elaborate trafficking system may have required a complex signaling system—an old heritage from early eukaryotic ancestors (Dacks and Field 2007; Plattner and Verkhatsky 2015).

Acknowledgements Experimental work by the author cited herein has been supported by the German Research Council.

References

- Allen RD, Fok AK (1993) Nonclathrin vesicle coats and filament networks in the transition zone and trans Golgi region of the Golgi complex of *Paramecium*. *J Struct Biol* 110:215–226
- Allen RD, Fok AK (2000) Membrane trafficking and processing in *Paramecium*. *Int Rev Cytol* 198:277–318
- Allen RD, Naitoh Y (2002) Osmoregulation and contractile vacuoles of protozoa. *Int Rev Cytol* 215:351–394

- Allen PD, Tominaga T, Naitoh Y (2009) The contractile vacuole complex and cell volume control in protozoa. In: Evans DH (ed) Osmotic and ionic regulation: cells and animals. CRC Press, Taylor and Francis Group, Boca Raton, FL, pp 69–106
- Aufderheide KJ (1978) Motility events of trichocyst insertion in *Paramecium tetraurelia*. *J Protozool* 25:362–365
- Beisson J (2008) Preformed cell structure and cell heredity. *Prion* 2:1–8
- Bell WE, Preston RR, Yano J, Van Houten JL (2007) Genetic dissection of attractant induced conductances in *Paramecium*. *J Exp Biol* 210:357–365
- Bemm F, Schwarz R, Forster F, Schultz J (2009) A kinome of 2600 in the ciliate *Paramecium tetraurelia*. *FEBS Lett* 583:3589–3592
- Bonini NM, Nelson DL (1990) Phosphoproteins associated with cyclic nucleotide stimulation of ciliary motility in *Paramecium*. *J Cell Sci* 95:219–230
- Bowman GR, Elde NC, Morgan G, Winey M, Turkewitz AP (2005) Core formation and the acquisition of fusion competence are linked during secretory granule maturation in *Tetrahymena*. *Traffic* 6:303–323
- Brehm P, Eckert R (1978) Calcium entry leads to inactivation of calcium channel in *Paramecium*. *Science* 202:1203–1206
- Bright LJ, Kambesis N, Nelson SB, Jeong B, Turkewitz AP (2010) Comprehensive analysis reveals dynamic and evolutionary plasticity of Rab GTPases and membrane traffic in *Tetrahymena thermophila*. *PLoS Genet* 6:e1001155
- Briguglio JS, Kumar S, Turkewitz AP (2013) Lysosomal sorting receptors are essential for secretory granule biogenesis in *Tetrahymena*. *J Cell Biol* 203:537–550
- Cervia D, Catalani E, Belardinelli MC, Perrotta C, Picchietti S, Alimenti C, Casini G, Fausto AM, Vallesi A (2013) The protein pheromone Er 1 of the ciliate *Euplotes raikovi* stimulates human T cell activity: involvement of interleukin 2 system. *Exp Cell Res* 319:56–67
- Chalker DL, Meyer E, Mochizuki K (2013) Epigenetics of ciliates. *Cold Spring Harb Perspect Biol* 5:a017764
- Clark KD, Hennessey TM, Nelson DL (1993) External GTP alters the motility and elicits an oscillating membrane depolarization in *Paramecium tetraurelia*. *Proc Natl Acad Sci USA* 90:3782–3786
- Dacks JB, Field MC (2007) Evolution of the eukaryotic membrane trafficking system: origin, tempo and mode. *J Cell Sci* 120:2977–2985
- Dentler WL (1988) Fractionation of *Tetrahymena* ciliary membranes with triton X 114 and the identification of a ciliary membrane ATPase. *J Cell Biol* 107:2679–2688
- De Oндarza J, Symington SB, Van Houten JL, Clark JM (2003) G protein modulators alter the swimming behavior and calcium influx of *Paramecium tetraurelia*. *J Eukaryot Microbiol* 50:349–355
- Eckert R, Brehm P (1979) Ionic mechanisms of excitation in *Paramecium*. *Ann Rev Biophys Bioeng* 8:353–383
- Elde NC, Morgan G, Winey M, Sperling L, Turkewitz AP (2005) Elucidation of clathrin mediated endocytosis in *Tetrahymena* reveals an evolutionarily convergent recruitment of dynamin. *PLoS Genet* 1:e52
- Elwess NL, Van Houten JL (1997) Cloning and molecular analysis of the plasma membrane Ca^{2+} ATPase gene in *Paramecium tetraurelia*. *J Eukaryot Microbiol* 44:250–257
- Farrell A, Thirugnanam S, Lorestani A, Dvorin JD, Eidell KP, Ferguson DJP, Anderson White BR, Duraisingh MT, Marth GT, Gubbels MJ (2012) A DOC2 protein identified by mutational profiling is essential for apicomplexan parasite exocytosis. *Science* 335:218–221
- Flötenmeyer M, Momayezi M, Plattner H (1999) Immunolabeling analysis of biosynthetic and degradative pathways of cell surface components (glycocalyx) in *Paramecium* cells. *Eur J Cell Biol* 78:67–77
- Fok AK (1983) An inhibition and kinetic study of acid phosphatase in *Paramecium caudatum* and *Paramecium tetraurelia*. *J Protozool* 30:14–20
- Fok AK, Paeste RM (1982) Lysosomal enzymes of *Paramecium caudatum* and *Paramecium tetraurelia*. *Exp Cell Res* 139:159–169

- Fraga D, Sehring IM, Kissmehl R, Reiss M, Gaines R, Hinrichsen R, Plattner H (2010) Protein phosphatase 2B (PP2B, calcineurin) in *Paramecium*: partial characterization reveals that two members of the unusually large catalytic subunit family have distinct roles in calcium dependent processes. *Eukaryot Cell* 9:1049–1063
- Frankel J (2000) Cell biology of *Tetrahymena thermophila*. *Meth Cell Biol* 62:27–125
- Garreau de Loubresse N (1993) Early steps of the secretory pathway in *Paramecium*: Ultrastructural, immunocytochemical, and genetic analysis of trichocyst biogenesis. In: Plattner H (ed) *Membrane traffic in protozoa*. JAI Press, Greenwich, CT, USA, pp 27–59
- Gonda K, Komatsu M, Numata O (2000) Calmodulin and Ca^{2+} /calmodulin binding proteins are involved in *Tetrahymena thermophila* phagocytosis. *Cell Struct Funct* 25:243–251
- Hardt M, Plattner H (2000) Sub second quenched flow/X ray microanalysis shows rapid Ca^{2+} mobilization from cortical stores paralleled by Ca^{2+} influx during synchronous exocytosis in *Paramecium* cells. *Eur J Cell Biol* 79:642–652
- Hauser K, Pavlovic N, Kissmehl R, Plattner H (1998) Molecular characterization of a sarco (endo)plasmic reticulum Ca^{2+} ATPase gene from *Paramecium tetraurelia* and localization of its gene product to sub plasmalemmal calcium stores. *Biochem J* 334:31–38
- Hay JC (2007) Calcium: a fundamental regulator of intracellular membrane fusion? *EMBO Rep* 8:236–240
- Hemmersbach R, Braun M (2006) Gravity sensing and gravity related signaling pathways in unicellular model systems of protists and plants. *Sign Transduct* 6:432–442
- Hemmersbach R, Wilczek M, Stieber C, Bräucker R, Ivanova K (2002) Variable acceleration influences cyclic AMP levels in *Paramecium biaurelia*. *J Gravit Physiol* 9:P267–P268
- Hurtado Lorenzo A, Skinner M, El Annan J, Futai M, Sun Wada GH, Bourgoin S, Casanova J, Wildeman A, Bechoua S, Ausiello DA, Brown D, Marshansky V (2006) V ATPase interacts with ARNO and Arf6 in early endosomes and regulates the protein degradative pathway. *Nat Cell Biol* 8:124–136
- Husser MR, Hardt M, Blanchard MP, Hentschel J, Klauke N, Plattner H (2004) One way calcium spill over during signal transduction in *Paramecium* cells: from the cell cortex into cilia, but not in the reverse direction. *Cell Calcium* 36:349–358
- Kim K, Messinger LA, Nelson DL (1998) Ca^{2+} dependent protein kinases of *Paramecium* cloning provides evidence of a multigene family. *Eur J Biochem* 251:605–612
- Kissmehl R, Sehring IM, Wagner E, Plattner H (2004) Immunolocalization of actin in *Paramecium* cells. *J Histochem Cytochem* 52:1543–1559
- Kissmehl R, Schilde C, Wassmer T, Danzer C, Nuehse K, Lutter K, Plattner H (2007) Molecular identification of 26 syntaxin genes and their assignment to the different trafficking pathways in *Paramecium*. *Traffic* 8:523–542
- Klauke N, Plattner H (1997) Imaging of Ca^{2+} transients induced in *Paramecium* cells by a polyamine secretagogue. *J Cell Sci* 110:975–983
- Klauke N, Kissmehl R, Plattner H, Haga N, Watanabe T (1998) An exocytotic mutant of *Paramecium caudatum*: membrane fusion without secretory contents release. *Cell Calcium* 23:349–360
- Klumpp S, Cohen P, Schultz JE (1990) Okadaic acid, an inhibitor of protein phosphatase 1 in *Paramecium*, causes sustained Ca^{2+} dependent backward swimming in response to depolarizing stimuli. *EMBO J* 9:685–689
- Knoll G, Braun C, Plattner H (1991) Quenched flow analysis of exocytosis in *Paramecium* cells: time course, changes in membrane structure, and calcium requirements revealed after rapid mixing and rapid freezing of intact cells. *J Cell Biol* 113:1295–1304
- Krishnan A, Mustafa A, Sällman Almén M, Fredriksson R, Williams MJ, Schiöth HB (2015) Evolutionary hierarchy of vertebrate like heterotrimeric G protein families. *Mol Phylogenet Evol* 91:27–40
- Kutomi O, Hori M, Ishida M, Tominaga T, Kamachi H, Koll F, Cohen J, Yamada N, Noguchi M (2012) Outer dynein arm light chain 1 is essential for controlling the ciliary response to cyclic AMP in *Paramecium tetraurelia*. *Eukaryot Cell* 11:645–653

- Ladenburger EM, Plattner H (2011) Calcium release channels in *Paramecium*. Genomic expansion, differential positioning and partial transcriptional elimination. PLoS ONE 6:e27111
- Ladenburger EM, Korn I, Kasielke N, Wassmer T, Plattner H (2006) An Ins(1,4,5)P₃ receptor in *Paramecium* is associated with the osmoregulatory system. J Cell Sci 119:3705–3717
- Ladenburger EM, Sehring IM, Korn I, Plattner H (2009) Novel types of Ca²⁺ release channels participate in the secretory cycle of *Paramecium* cells. Mol Cell Biol 29:3605–3622
- Lampert TJ, Coleman KD, Hennessey TM (2011) A knockout mutation of a constitutive GPCR in *Tetrahymena* decreases both G protein activity and chemoattraction. PLoS ONE 6:e28022
- Lee HC (2012) Cyclic ADP ribose and nicotinic acid adenine dinucleotide phosphate (NAADP) as messengers for calcium mobilization. J Biol Chem 287:31633–31640
- Luporini P, Alimenti C, Vallesi A (2014) Ciliate mating types and pheromones. In: Hausmann K, Radek R (eds) Cilia and flagella, ciliates and flagellates. Schweizerbart Science Publishers, Stuttgart, pp 95–118
- Machemer H (1988a) Electrophysiology. In: Görtz HD (ed) *Paramecium*. Springer, Berlin, Heidelberg, pp 185–215
- Machemer H (1988b) Motor control of cilia. In: Görtz HD (ed) *Paramecium*. Springer, Berlin, Heidelberg, pp 216–235
- Machemer H (2014) How do protists keep up? In: Hausmann K, Radek R (eds) Cilia and flagella, ciliates and flagellates. Schweizerbart Science Publishers, Stuttgart, pp 133–146
- Machemer H, Ogura A (1979) Ionic conductances of membranes in ciliated and deciliated *Paramecium*. J Physiol 296:49–60
- Machemer H, Bräucker R, Machemer Röhnisch S, Nagel U, Neugebauer DC, Weskamp M (1998) The linking of extrinsic stimuli to behaviour: roles of cilia in ciliates. Eur J Protistol 34:254–261
- Min SW, Chang WP, Südhof TC (2007) E Syts, a family of membranous Ca²⁺ sensor proteins with multiple C2 domains. Proc Natl Acad Sci USA 104:3823–3828
- Momayezi M, Lumpert CJ, Kersken H, Gras U, Plattner H, Krinks MH, Klee CB (1987) Exocytosis induction in *Paramecium tetraurelia* cells by exogenous phosphoprotein phosphatase in vivo and in vitro: possible involvement of calcineurin in exocytotic membrane fusion. J Cell Biol 105:181–189
- Momayezi M, Kissmehl R, Plattner H (2000) Quantitative immunogold localization of protein phosphatase 2B (calcineurin) in *Paramecium* cells. J Histochem Cytochem 48:1269–1281
- Neher E (1998) Vesicle pools and Ca²⁺ microdomains: new tools for understanding their roles in neurotransmitter release. Neuron 20:389–399
- Park HY, Kim SA, Korlach J, Rhoades E, Kwok LW, Zipfle WR, Waxham MN, Webb WW, Pollack L (2008) Conformational changes of calmodulin upon Ca²⁺ binding studied with a microfluidic mixer. Proc Natl Acad Sci USA 105:542–547
- Plattner H (2010) Membrane trafficking in protozoa SNARE proteins, H⁺ ATPase, actin, and other key players in ciliates. Int Rev Cell Mol Biol 280:79–184
- Plattner H (2014) Calcium regulation in the protozoan model, *Paramecium tetraurelia*. J Eukaryot Microbiol 61:95–114
- Plattner H (2015a) Molecular aspects of calcium signalling at the crossroads of unikont and bikont eukaryote evolution – the ciliated protozoan *Paramecium* in focus. Cell Calcium 57:174–185
- Plattner H (2015b) The contractile vacuole complex of protists – new cues to function and biogenesis. Crit Rev Microbiol 41:218–227
- Plattner H (2016a) Signalling in ciliates: Long and short range signals and molecular determinants for cellular dynamics. Biol Rev (in press). doi: [10.1111/brv.12218](https://doi.org/10.1111/brv.12218)
- Plattner H (2016b) Signals regulating vesicle trafficking in *Paramecium* cells. This volume
- Plattner H, Klauke N (2001) Calcium in ciliated protozoa: sources, regulation, and calcium regulated cell functions. Int Rev Cytol 201:115–208
- Plattner H, Kissmehl R (2003a) Dense core secretory vesicle docking and exocytotic membrane fusion in *Paramecium* cells. Biochim Biophys Acta (Mol Cell Res) 1641:183–193
- Plattner H, Kissmehl R (2003b) Molecular aspects of membrane trafficking in *Paramecium*. Int Rev Cytol 232:185–216

- Plattner H, Verkhatsky A (2013) Ca^{2+} signalling early in evolution – all but primitive. *J Cell Sci* 126:2141–2150
- Plattner H, Verkhatsky A (2015) The ancient roots of calcium signalling evolutionary tree. *Cell Calcium* 57:123–132
- Plattner H, Westphal C, Tiggemann R (1982) Cytoskeleton secretory vesicle interactions during the docking of secretory vesicles at the cell membrane in *Paramecium tetraurelia* cells. *J Cell Biol* 92:368–377
- Plattner H, Stürzl R, Matt H (1985) Synchronous exocytosis in *Paramecium* cells. IV. Polyamino compounds as potent trigger agents for repeatable trigger redocking cycles. *Eur J Cell Biol* 36:32–37
- Plattner H, Knoll G, Pape R (1993) Synchronization of different steps of the secretory cycle in *Paramecium tetraurelia*: trichocyst exocytosis, exocytosis coupled endocytosis, and intracellular transport. In: Plattner H (ed) *Membrane traffic in protozoa*. JAI Press, Greenwich (CT) and London, pp 123–148
- Plattner H, Braun C, Hentschel J (1997) Facilitation of membrane fusion during exocytosis and exocytosis coupled endocytosis and acceleration of “ghost” detachment in *Paramecium* by extracellular calcium. A quenched flow/freeze fracture analysis. *J Membr Biol* 158:197–208
- Plattner H, Sehring IM, Mohamed IK, Miranda K, De Souza W, Billington R, Genazzani A, Ladenburger EM (2012) Calcium signaling in closely related protozoan groups (Alveolata): non parasitic ciliates (*Paramecium*, *Tetrahymena*) vs. parasitic Apicomplexa (*Plasmodium*, *Toxoplasma*). *Cell Calcium* 51:351–382
- Pouphile M, Lefort Tran M, Plattner H, Rossignol M, Beisson J (1986) Genetic dissection of the morphogenesis of exocytosis sites in *Paramecium*. *Biol Cell* 56:151–162
- Preston RR, Saimi Y, Kung C (1992) Calcium current activated upon hyperpolarization of *Paramecium tetraurelia*. *J Gen Physiol* 100:233–251
- Reuter AT, Stuermer CAO, Plattner H (2013) Identification, localization, and functional implications of the microdomain forming stomatin family in the ciliated protozoan *Paramecium tetraurelia*. *Eukaryot Cell* 12:529–544
- Rizo J, Chen X, Arac D (2006) Unraveling the mechanisms of synaptotagmin and SNARE function in neurotransmitter release. *Trends Cell Biol* 16:339–350
- Rothman JE (2014) The principle of membrane fusion in the cell (Nobel lecture). *Angew Chemie Int Ed* 53:12676–12694
- Saimi Y, Kung C (2002) Calmodulin as an ion channel subunit. *Annu Rev Physiol* 64:289–311
- Schilde C, Wassmer T, Mansfeld J, Plattner H, Kissmehl R (2006) A multigene family encoding R SNAREs in the ciliate *Paramecium tetraurelia*. *Traffic* 7:440–455
- Schilde C, Schönmeyer B, Sehring IM, Plattner H (2010) Distinct subcellular localization of a group of synaptobrevin like SNAREs in *Paramecium tetraurelia* and effects of silencing SNARE specific chaperone NSF. *Eukaryot Cell* 9:288–305
- Sehring IM, Plattner H (2004) Ca^{2+} oscillations mediated by exogenous GTP in *Paramecium* cells: assessment of possible Ca^{2+} sources. *Cell Calcium* 36:409–420
- Sehring IM, Reiner C, Mansfeld J, Plattner H, Kissmehl R (2007) A broad spectrum of actin paralogs in *Paramecium tetraurelia* cells display differential localization and function. *J Cell Sci* 120:177–190
- Sehring IM, Klotz C, Beisson J, Plattner H (2009) Rapid downregulation of the Ca^{2+} signal after exocytosis stimulation in *Paramecium* cells: essential role of a centrion rich filamentous cortical network, the infraciliary lattice. *Cell Calcium* 45:89–97
- Simon M, Plattner H (2014) Unicellular eukaryotes as models in cell and molecular biology: critical appraisal of their past and future value. *Int Rev Cell Mol Biol* 309:141–198
- Stock C, Grønlien HK, Allen RD (2002) The ionic composition of the contractile vacuole fluid of *Paramecium* mirrors ion transport across the plasma membrane. *Eur J Cell Biol* 81:505–515
- Südhof TC (2014) The molecular machinery of neurotransmitter release (Nobel lecture). *Angew Chemie Int Ed* 53:12696–12717
- Sun Wada GH, Wada Y (2015) Role of vacuolar type proton ATPase in signal transduction. *Biochim Biophys Acta* 1847:1166–1172

- Tian M, Chen X, Xiong Q, Xiong J, Xiao C, Ge F, Yang F, Miao W (2014) Phosphoproteomic analysis of protein phosphorylation networks in *Tetrahymena thermophila*, a model single celled organism. *Mol Cell Proteom* 13:503–519
- Vallesi A, Di Pretoro B, Ballarini P, Apone F, Luporini P (2010) A novel protein kinase from the ciliate *Euplotes raikovi* with close structural identity to the mammalian intestinal and male germ cell kinases: characterization and functional implications in the autocrine pheromone signaling loop. *Protist* 161:250–263
- Wassmer T, Froissard M, Plattner H, Kissmehl R, Cohen J (2005) The vacuolar proton ATPase plays a major role in several membrane bounded organelles in *Paramecium*. *J Cell Sci* 118:2813–2825
- Wassmer T, Kissmehl R, Cohen J, Plattner H (2006) Seventeen a subunit isoforms of *Paramecium* V ATPase provide high specialization in localization and function. *Mol Biol Cell* 17:917–930
- Wassmer T, Sehring IM, Kissmehl R, Plattner H (2009) The V ATPase in *Paramecium*: functional specialization by multiple gene isoforms. *Eur J Physiol* 457:599–607
- Wheeler GL, Brownlee C (2008) Ca^{2+} signalling in plants and green algae – changing channels. *Trends Plant Sci* 13:506–514
- Yang WQ, Braun C, Plattner H, Purvee J, Van Houten JL (1997) Cyclic nucleotides in glutamate chemosensory signal transduction of *Paramecium*. *J Cell Sci* 110:2567–2572
- Yano J, Rajendran A, Valentine MS, Saha M, Ballif BA, Van Houten JL (2013) Proteomic analysis of the cilia membrane of *Paramecium tetraurelia*. *J Proteom* 78:113–122
- Yano JY, Valentine MS, Van Houten JL (2015) Novel insights into the development and function of cilia using the advantages of the *Paramecium* cell and its many cilia. *Cells* 4:297–314
- Zhen Y, Stenmark H (2015) Cellular functions of Rab GTPases at a glance. *J Cell Sci* 128:3171–3176