Towards a General Understanding of Bacterial Interactions

Zhijie Zhang,¹
Mark van Kleunen,^{1,2}
Lutz Becks,³ and
Madhav P. Thakur ^{1,*}

Understanding the general rules of microbial interactions is central for advancing microbial ecology. Recent studies show that interaction range, interaction strength, and community context determine bacterial interactions and the coexistence and evolution of bacteria. We highlight how these factors could contribute to a general understanding of bacterial interactions.

Microbial Community Ecology

A central tenet of community ecology is to understand the rules of species interactions and their consequences for community structure, such as species diversity. Historically, microbial interactions and their outcomes were studied to test theories of community ecology (e.g., [1,2]) which were developed for plants and animals [3,4]. This trend has changed remarkably in the last few years with rapid progress in microbial community ecology due mainly to the continued growth of molecular technologies [4,5]. Today, microbial community ecology has become an integral branch of ecology and microbiology and it continues to advance our understanding of the implications of microbial interactions for disease control, food production, climate change mitigation, and biodiversity conservation.

Bacteria are among the most studied microorganisms in microbial community ecology, particularly when it comes to experimental assessments of microbial

interactions [4]. Yet, we still lack a consensus for how bacterial species interact, and how those interactions may regulate the diversity of bacteria [6]. This is due to several key challenges in studying bacterial interactions, such as delineating the spatial and temporal scale at which bacteria interact, quantifying the strength of their interactions, dependence of their interactions on abiotic conditions, and constantly varying interactions owing to rapid evolution and changes in bacterial abundance. Three recent studies provide insight into overcoming these challenges [7–9]. These studies independently show interaction range [7], interaction strength [8], and community context [9] as three factors determining bacterial species interaction and the coexistence and evolution of bacteria. We highlight how simultaneous consideration of these three factors can help to achieve a more general understanding of bacterial interactions (Figure 1) and their relevance for building predictive models for microbial community ecology.

Interaction Range, Interaction Strength, and Community Context

Interaction range is the spatial domain in which an individual can interact with other individuals of either the same species or a different species. Usually, the interaction ranges of several bacterial cells are limited to their close surroundings, such as biofilms [10]. However, even within biofilms, some cells are close neighbors while others are distant. Whether bacterial cells in spatially structured biofilms interact with close and distant neighbors is not well understood given the complexity in empirically quantifying the distances between bacteria during their interactions. Dal Co *et al.* [7] used an elegant approach of measuring the interaction range of bacterial cells as the size of the neighborhood from which one bacterial cell can retrieve the amino acids produced by neighboring cells. They highlight that the interaction range of bacteria is short, that is,

individuals interact only with their immediate neighbors. More specifically, they used two genotypes of *Escherichia coli* that exchange essential amino acids for their growth. They showed that the growth of one genotype was promoted when most of its neighbors in its interaction range were of the other genotype [7]. This study contributes to a general understanding of how the range of interaction matters for facilitation and coexistence at the cellular level.

While the size of the interaction range of a bacterial cell indicates the distance over which it can affect another cell (Figure 1), the direction (mutualistic or antagonistic) of these bacterial interactions is likely to depend on how bacterial cells modify their immediate environment [11]. Towards this end, Ratzke *et al.* [8] showed that increased nutrient availability promoted the ability of soil bacterial species to modify their environment (e.g., pH values), which subsequently magnified the negative interactions between soil bacterial species. Interestingly, this study highlighted that the strength of negative interactions between bacterial species was mediated mainly by the production of toxic metabolites and could not be attenuated by refreshing nutrients. These strong competitive interactions between bacterial species impeded species coexistence and decreased the stability of the bacterial communities.

There is a growing interest in understanding the roles of evolutionary processes in driving species coexistence [12]. An experimental study by Scheuerl *et al.* [9] demonstrated that bacterial species (obtained from the rainwater in tree holes) had greater evolutionary responses to environmental changes (low pH) when the bacterial communities had a low diversity, presumably owing to weaker competitive interactions. These authors show that strong competitive interactions, as found in the diverse bacterial communities, limit

Trends in Microbiology

Figure 1. Some General Rules and Outcomes of Bacterial Interactions. Interaction range (short and long), interaction strength (weak and strong), and community context (low and high diversity) affect bacterial coexistence and evolution. We discuss several possibilities for bacterial coexistence using an example of a focal bacterial species (gray filled) and its interaction with neighboring bacterial species. For simplicity, we focus on competitive interactions, but the framework can be adapted to include both competitive and facilitative interactions. (A) Low-diversity, short-interaction range. In a low-diversity scenario, when the gray bacterial species has short interaction range (indicated by the gray circle) and interaction strength (indicated by the thicker double-sided arrows), it excludes its nearest neighbors, like the yellow bacterial species, by its ability to modify the environment (e.g., the pH). However, when considering evolution, such competitive interactions could also lead to diversification (indicated by the emergence of the orange bacteria) when the yellow bacteria can adapt to the new environment modified by the gray bacteria. (B) Low-diversity, long-interaction range. When the same gray bacterial species has a longer interaction range it excludes more yellow bacteria in the absence of evolution, whereas a greater diversification of the yellow bacteria takes place in the presence of evolution. (C) High-diversity, short-interaction range. In this scenario, we illustrate the example of five bacterial species living next to the gray bacteria, but only one of them (the yellow one) falls within the short interaction range. The results are identical to the outcomes shown in scenario (A) despite the difference in species diversity. (D) High-diversity, long-interaction range. When the same gray bacterial species has a longer interaction range it excludes the four neighboring species within its interaction range in the absence of evolution. When considering evolution, Scheuerl *et al.* [9] showed that evolution is constrained when multiple species are interacting (presumably owing to strong interaction strength – not shown in the figure), implying that evolution might not be rapid enough to prevent exclusion of the four neighboring species.

the capacity of bacterial species to evolve, thereby highlighting the importance of 'community context' as a key predictor of bacterial evolutionary dynamics. Taken together, these three recent independent studies [7–9] have provided us with potential general rules on how bacteria interact, and further highlight the often overlooked implications of bacterial interactions for evolutionary dynamics (Figure 1).

Outlook

We suggest that interaction range, interaction strength, and community context should be considered simultaneously for a better understanding of bacterial interactions. In particular, considering evolution will enhance our ability to predict variations in a bacterial community in a given environment (some scenarios are shown in Figure 1). Microbial ecologists have con-

sistently emphasized the importance of making microbial ecology a predictive science rather than just a descriptive science as most research in microbial ecology still continues to be on microbial inventories in various environments [3,6]. We believe that, to make microbial ecology more predictive, we need to integrate general rules of microbial interactions to study microbial community diversity, functioning, and

evolution. The rules highlighted by the three studies discussed earlier [7–9] are an important step towards this goal. Future studies can further explore how these rules may affect each other (Figure 1) – for instance, how the interaction range of bacteria and their ability to modify the environment relate to each other. Moreover, it will be important to test the significance of interaction range, interaction strength, and community context for regulating bacterial community structure in the context of bacterial interactions with other microorganisms (e.g., fungi) and their predators (e.g., protists), and also in the context of different and novel environments, such as those created by climate warming and drought.

Acknowledgments

We thank two anonymous reviewers for helpful comments. Z.Z. acknowledges funding from the

China Scholarship Council (201606100049). M.P.T. acknowledges funding from the German Research Foundation (DFG, TH 2307/1-1, 2-1).

¹Ecology, Department of Biology, University of Konstanz, 78464, Konstanz, Germany

²Zhejiang Provincial Key Laboratory of Plant Evolutionary Ecology and Conservation, Taizhou University, Taizhou 318000, China

³Aquatic Ecology and Evolution, Department of Biology, University of Konstanz, 78464, Konstanz, Germany

References

1. Gause, G.F. (1934) Experimental analysis of Vito Volterra's mathematical theory of the struggle for existence. *Science* 79, 16–17
2. Kassen, R. et al. (2000) Diversity peaks at intermediate productivity in a laboratory microcosm. *Nature* 406, 508–512
3. Little, A.E.F. et al. (2008) Rules of engagement: interspecies interactions that regulate microbial communities. *Annu. Rev. Microbiol.* 62, 375–401
4. Konopka, A. (2009) What is microbial community ecology. *ISME J.* 3, 1223–1230
5. Faust, K. and Raes, J. (2012) Microbial interactions: From networks to models. *Nat. Rev. Microbiol.* 10, 538–550
6. Prosser, J.I. and Martiny, J.B.H. (2020) Conceptual challenges in microbial community ecology. *Philos. Trans. R. Soc. B Biol. Sci.* 375, 20190241
7. Dial Co, A. et al. (2020) Short-range interactions govern the dynamics and functions of microbial communities. *Nat. Ecol. Evol.* 4, 366–375
8. Ratzke, C. et al. (2020) Strength of species interactions determines biodiversity and stability in microbial communities. *Nat. Ecol. Evol.* 4, 376–383
9. Scheuerl, T. et al. (2020) Bacterial adaptation is constrained in complex communities. *Nat. Commun.* 11, 754
10. Nadeil, C.D. et al. (2016) Spatial structure, cooperation and competition in biofilms. *Nat. Rev. Microbiol.* 14, 589–600
11. Goldford, J.E. et al. (2018) Emergent simplicity in microbial community assembly. *Science* 361, 469–474
12. Turcotte, M.M. and Levine, J.M. (2016) Phenotypic plasticity and species coexistence. *Trends Ecol. Evol.* 31, 803–813