PAIREF: paired refinement also for *Phenix* users

Martin Malý,^{a,b} Kay Diederichs,^c Jan Dohnálek^b and Petr Kolenko^{a,b*}

^aFaculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Břehová 7, 115 19 Prague 1, Czech Republic, ^bInstitute of Biotechnology of the Czech Academy of Sciences, BIOCEV, Průmyslová 595, 252 50 Vestec, Czech Republic, and ^cUniversity of Konstanz, Box M647, 78457 Konstanz, Germany. *Correspondence e-mail: petr.kolenko@jfifi.cvut.cz

Keywords: macromolecular crystallography; PAIREF; Phenix; X-ray diffraction; paired refinement; high-resolution limit.

In macromolecular crystallography, paired refinement is generally accepted to be the optimal approach for the determination of the high-resolution cutoff. The software tool *PAIREF* provides automation of the protocol and associated analysis. Support for *phenix.refine* as a refinement engine has recently been implemented in the program. This feature is presented here using previously published data for thermolysin. The results demonstrate the importance of the complete cross-validation procedure to obtain a thorough and unbiased insight into the quality of high-resolution data.

1. Introduction

During diffraction data processing, a cutoff is usually applied to reject high-resolution data that do not improve the model during structure refinement. In recent decades, a number of criteria have been used to decide on this cutoff (Karplus & Diederichs, 2015). Nowadays, paired refinement is considered to be the optimal approach for the determination of this parameter (Karplus & Diederichs, 2012; Diederichs & Karplus, 2013). In brief, a conservative cutoff has to be chosen first to secure reliable data. Data from higher resolution shells are added in a stepwise process to the model refinement and their impact on the model quality is verified. Such an approach is time-consuming and prone to errors when no automation is available. Paired refinement is particularly sensible in the later stages of structure refinement when the decision on the refinement program has already been made.

Recently, we developed an automatic tool (Malý *et al.*, 2020) for paired refinement that uses *REFMAC5* (Murshudov *et al.*, 2011) for structure refinement. *PAIREF* provides comprehensive data analysis together with merging statistics, correlation coefficients (for example CC_{work} and CC^*) and indicators of the stability of structure refinement. Here, we present a new feature of *PAIREF*: a module that performs structure refinement with *phenix.refine* (Afonine *et al.*, 2012). The algorithm does not differ from that relying on *REFMAC5*. Structure-refinement parameters for *phenix.refine* can be specified in detail through a definition file (option `--def setting.def`). Besides the current implementation of paired refinement in the *Phenix* package (Liebschner *et al.*, 2019), our tool provides additional features, for example a complete cross-validation procedure (Brünger, 1993; Jiang & Brünger, 1994). For each set of test reflections, the paired refinement protocol is run in parallel, and averaged data-quality indicators are reported. Both the standard and the complete cross-validation procedure are shown for test data. Moreover, a graphical user interface (Fig. 1) has recently been developed to simplify job execution.

2. Materials and methods

One of the previously reported cases where paired refinement using *REFMAC5* proved to be helpful was the crystal structure of thermolysin. In the previous report (Winter *et al.*, 2018), the initial high-resolution cutoff was set to 1.8 Å and reflections were added in thin shells with a width of 0.01 Å. A decrease in R_{gap} ($R_{\text{gap}} = R_{\text{free}} - R_{\text{work}}$) was observed up to a resolution of 1.56 Å. Similarly, we carried out paired refinement while adding shells with a width of 0.10 Å, referred to as run 0 in this manuscript (Malý *et al.*, 2020). The R_{free} values systematically decreased up to a resolution of 1.5 Å, which indicates model improvement. Thus, we suggested cutting the data at this resolution level. Here, we show the results from paired refinement carried out with *PAIREF* using *phenix.refine* (*Phenix* version 1.16-3546) instead of *REFMAC5*.

We performed three distinct runs of *PAIREF* using the previously reported diffraction data for thermolysin. The diffraction images were processed with *xia2* (Winter, 2010) employing *DIALS* (Winter *et al.*, 2018) and *AIMLESS* (Evans & Murshudov, 2013) at 1.5 Å resolution in space group *P6₁22*. The input model for all runs originated from the structure of thermolysin (PDB entry 3n21; Behnen *et al.*, 2012) with water molecules. To remove model bias, the atomic coordinates were perturbed by an average of 0.25 Å and all ADPs were set to their average value with *phenix.pdbtools* (Liebschner *et al.*, 2019). Subsequently, restrained refinement was performed with *phenix.refine* at a resolution of 1.8 Å, converging sufficiently in 12 cycles. We performed three *PAIREF* runs: run 1, a standard run with the addition of high-resolution shells with a width of 0.10 Å; run 2, a fine-sliced standard run with a width of 0.01 Å; and run 3, the complete cross-validation procedure

using all 20 sets of test reflections with a shell width of 0.10 Å. As an example, the command to launch run 1 (in the Unix shell or *Phenix* Command Prompt) is `cctbx.python -m pairef --XYZIN 3n21_edit05_shaken.pdb --HKLIN thermc_merged.mtz -u thermc_unmerged.mtz --phenix --project thermc_step0-10A_phenix --prerefinement-ncyc 12 -i 1.8 -r 1.7,1.6,1.5,1.4`; the execution of this run using the graphical interface is also shown in Fig. 1.

The results are shown in Fig. 2. The related merging statistics and details of run 0 have previously been published (Malý *et al.*, 2020).

3. Results and discussion

Paired refinements using the thermolysin data demonstrate the differences that may appear using various refinement engines together with the importance of the complete cross-validation procedure (Fig. 2). Results from the individual runs, run 0 (*REFMAC5*, 0.10 Å step), run 1 (*phenix.refine*, 0.10 Å step), run 2 (*phenix.refine*, 0.01 Å step) and run 3 (*phenix.refine*, 0.10 Å step, complete cross-validation), vary in the suggestion of cutoff choice: 1.5, 1.6, 1.52 and 1.5 Å, respectively. For instance, we obtain different suggestions with *phenix.refine* and *REFMAC5* (runs 0 and 1) while using the standard scenario, *i.e.* a 0.10 Å step and exclusion of the original set of test reflections (test flag equals 0). Moreover, the complete cross-validation procedure (run 3) suggests a higher cutoff, at 1.5 Å, than the standard run 1. As the former provides more general and meaningful insight into the quality of high-resolution data (Figs. 2c and 2d), it leads us to a final decision on the high-resolution cutoff at 1.5 Å resolution. This choice is in agreement with the fine-sliced run 2, where the R_{gap} value is minimal at 1.52 Å (Fig. 2b). The last resolution shell (1.6–1.5 Å) merging statistics are as follows: $I/\sigma(I) = 0.8$, $R_{\text{p.i.m.}} = 0.598$, $CC_{1/2} = 0.445$, completeness 91.8% (Malý *et al.*, 2020).

PAIREF now supports both of the most frequently used refinement programs. The refinement approach of *phenix.refine* differs from that of *REFMAC5* in several aspects (Shabalin *et al.*, 2018), such as bulk-solvent modeling (Weichenberger *et al.*, 2015), second-derivatives approximation and separate refinement of coordinates and ADPs. Thus, certain variations in the paired refinement results could be expected. The use of *PAIREF* is not intended as a tool to decide on the choice of the refinement program, but rather as a step in structure refinement. Both refinement engines specifically treat special cases such as twinning (Campeotto *et al.*, 2018), extremes of resolution (Headd *et al.*, 2012; Kovalevskiy *et al.*, 2018), complex NCS and mixed anisotropic/isotropic ADP refinement *etc.* Hence, the support for multiple refinement programs in *PAIREF* can be useful when particular data qualities need to be addressed differently (Švecová *et al.*, 2021).

The complete cross-validation procedure is recommended for thorough determination of the proper resolution cutoff. The information value of the *PAIREF* analysis could be

Figure 1
PAIREF graphical user interface set for the execution of run 1.

Figure 2

Report plots given by *PAIREF* from three different runs of paired refinement using *phenix.refine* with the thermolysin data. The differences in the overall R values were obtained as follows: for each incremental step of resolution increase $X \rightarrow Y$, the R values were calculated at the lower resolution X . (a) Run 1: differences in the overall R values; resolution shells with a width of 0.10 Å were added stepwise. R_{free} decreases up to 1.60 Å resolution, indicating model improvement. (b) Run 2: graph of R_{gap} calculated using data up to 1.8 Å resolution depending on the high-resolution cutoff; resolution shells with a width of 0.01 Å were added stepwise. Minimal R_{gap} is observed at 1.52 Å resolution. (c) Run 3: differences in the overall R values averaged over all 20 sets of test reflections. The standard error of the mean is shown in orange. (d) Run 3: differences in the overall R values relating to all 20 sets of test reflections for the incremental step of resolution from 1.6 to 1.5 Å. Despite the increasing R_{free} value while using the original set (test flag equals 0) and four other sets, R_{free} decreases for 14 sets. After averaging over all 20 sets, a decreasing trend is observed for this resolution shell.

further increased by the implementation of a statistic which is independent of the selection of test reflections: R_{complete} (Luebben & Gruene, 2015). A detailed description of the *PAIREF* program and its algorithm, output and possibilities is provided in the primary reference (Malý *et al.*, 2020) and at the web page <https://pairef.fjfi.cvut.cz>.

Acknowledgements

We would like to thank Jan Stránský (Institute of Biotechnology, Czech Academy of Sciences) for helpful insights in software development and Christoph Parthier (Martin Luther University, Halle-Wittenberg) for program testing.

Funding information

The following funding is acknowledged: Ministry of Education, Youth and Sports CR (grant No. CZ.02.1.01/0.0/0.0/16_019/0000778; grant No. CZ.02.1.01/0.0/0.0/15_003/0000447; grant No. CZ.1.05/1.1.00/02.0109); Czech Science Foundation (grant No. 18-10687S); Czech Academy of Sciences (grant No. 86652036); Grant Agency of the Czech Technical University in Prague (grant No. SGS19/189/OHK4/3T/14).

References

- Afonine, P. V., Grosse-Kunstleve, R. W., Echols, N., Headd, J. J., Moriarty, N. W., Mustyakimov, M., Terwilliger, T. C., Urzhumtsev, A., Zwart, P. H. & Adams, P. D. (2012). *Acta Cryst.* **D68**, 352–367.

- Behnen, J., Köster, H., Neudert, G., Craan, T., Heine, A. & Klebe, G. (2012). *ChemMedChem*, **7**, 248–261.
- Brünger, A. T. (1993). *Acta Cryst. D***49**, 24–36.
- Campeotto, I., Lebedev, A., Schreurs, A. M. M., Kroon-Batenburg, L. M. J., Lowe, E., Phillips, S. E. V., Murshudov, G. N. & Pearson, A. R. (2018). *Sci. Rep.* **8**, 14876.
- Diederichs, K. & Karplus, P. A. (2013). *Acta Cryst. D***69**, 1215–1222.
- Evans, P. R. & Murshudov, G. N. (2013). *Acta Cryst. D***69**, 1204–1214.
- Headd, J. J., Echols, N., Afonine, P. V., Grosse-Kunstleve, R. W., Chen, V. B., Moriarty, N. W., Richardson, D. C., Richardson, J. S. & Adams, P. D. (2012). *Acta Cryst. D***68**, 381–390.
- Jiang, J.-S. & Brünger, A. T. (1994). *J. Mol. Biol.* **243**, 100–115.
- Karplus, P. A. & Diederichs, K. (2012). *Science*, **336**, 1030–1033.
- Karplus, P. A. & Diederichs, K. (2015). *Curr. Opin. Struct. Biol.* **34**, 60–68.
- Kovalevskiy, O., Nicholls, R. A., Long, F., Carlon, A. & Murshudov, G. N. (2018). *Acta Cryst. D***74**, 215–227.
- Liebschner, D., Afonine, P. V., Baker, M. L., Bunkóczi, G., Chen, V. B., Croll, T. I., Hintze, B., Hung, L.-W., Jain, S., McCoy, A. J., Moriarty, N. W., Oeffner, R. D., Poon, B. K., Prisant, M. G., Read, R. J., Richardson, J. S., Richardson, D. C., Sammito, M. D., Sobolev, O. V., Stockwell, D. H., Terwilliger, T. C., Urzhumtsev, A. G., Videau, L. L., Williams, C. J. & Adams, P. D. (2019). *Acta Cryst. D***75**, 861–877.
- Luebben, J. & Gruene, T. (2015). *Proc. Natl Acad. Sci. USA*, **112**, 8999–9003.
- Malý, M., Diederichs, K., Dohnálek, J. & Kolenko, P. (2020). *IUCrJ*, **7**, 681–692.
- Murshudov, G. N., Skubák, P., Lebedev, A. A., Pannu, N. S., Steiner, R. A., Nicholls, R. A., Winn, M. D., Long, F. & Vagin, A. A. (2011). *Acta Cryst. D***67**, 355–367.
- Shabalin, I. G., Porebski, P. J. & Minor, W. (2018). *Crystallogr. Rev.* **24**, 236–262.
- Švecová, L., Østergaard, L. H., Skálová, T., Schnorr, K. M., Koval', T., Kolenko, P., Stránský, J., Sedlák, D., Dušková, J., Trundová, M., Hašek, J. & Dohnálek, J. (2021). *Acta Cryst. D***77**, 755–775.
- Weichenberger, C. X., Afonine, P. V., Kantardjiev, K. & Rupp, B. (2015). *Acta Cryst. D***71**, 1023–1038.
- Winter, G. (2010). *J. Appl. Cryst.* **43**, 186–190.
- Winter, G., Waterman, D. G., Parkhurst, J. M., Brewster, A. S., Gildea, R. J., Gerstel, M., Fuentes-Montero, L., Vollmar, M., Michels-Clark, T., Young, I. D., Sauter, N. K. & Evans, G. (2018). *Acta Cryst. D***74**, 85–97.