ASYMMETRIC SYNTHESSES VIA HETEROCYCLIC INTERMEDIATES—XXII¹

ENANTIOSELECTIVE SYNTHESIS OF α -ALKENYL GLYCINE METHYL ESTERS AND α -ALKENYL GLYCINES (β,γ -UNSATURATED AMINO ACIDS)

ULRICH SCHÖLLKOPF,* JOACHIM NOZULAK and ULRICH GROTH
Institut für Organische Chemie der Universität, Tammannstraße 2, D-3400 Göttingen,
Federal Republic of Germany

(Received in USA 25 April 1983)

Abstract—Enantioselective syntheses of α -alkenyl glycines of type **10** and of type **23** are described that provide these uncommon amino acids with predictable configuration and with ee-values of >95%. Both approaches are based on the bislactim ether method developed by Schöllkopf *et al.* As for **10**: The lithiated bis-lactim ether **6** of cyclo(L-val-gly) is reacted with 2-[(dimethyl t-butyl)silyl]alkanals **2** to give the addition products **7** with de > 95%. These on acid hydrolysis afford L-valinate **8** and the methyl (2*R*)-2-amino-4-(dimethyl t-butyl)silyl-3-hydroxyalkanoates **9** which are convertible into the (*R*)- α -alkenyl glycines of type **10**. The scope of this synthesis is limited by the fact that the compounds **9** are thermolabile when disubstituted at C-4. As for **23**: The lithiated bis-lactim ether **6** is reacted with thioketones **14** to give the addition products **15** with de > 95%. The *S*-methyl compounds **16** undergo elimination to give regioselectively the olefins **18** when treated with Raney-Ni. Alternatively, the olefins **18** are obtained from the sulfonium salts **24** by dimethyl sulfide elimination, although this route is less regiospecific. The compounds **18** are cleaved by dilute hydrochloric acid, liberating L-valinate **8** and (*R*)- α -alkenyl glycine methyl esters **21**, which on further hydrolysis yield (*R*)- α -alkenyl glycines **23**. This synthesis is limited only by the availability of thioketones **14**.

Optically active, non-proteinogenic amino acids deserve attention because of their documented or potential biological activity. Some are valuable pharmaceuticals, such as L-Dopa, (*S*)- α -Methyldopa, D-Penicillamine, or D-Cycloserine. Others are components of pharmaceuticals, for instance D-phenylglycine or D-(*p*-hydroxy-phenylglycine) in the semisynthetic penicillins Ampicillin or Amoxycillin.

In biochemistry, they are valuable tools to investigate the mechanism of enzyme reactions.^{2,3} In fact, enzyme inhibition studies with non-proteinogenic amino acids have furnished valuable information about the mode of action of certain enzymes.^{2,3}

Obviously, there is a demand for optically active—if possible optically pure-uncommon amino acids both for pure and applied organic or bioorganic chemistry. Since asymmetric synthesis⁴ is—at least in principle—the shortest and most economic way to optically active compounds, it is a challenge for the synthetic organic chemist, to develop asymmetric syntheses of amino acids.⁵

Over the past four years our group has tried to elaborate asymmetric syntheses of uncommon amino acids. Our approach is based on heterocyclic chemistry and on the following concept.⁶ (1) From a racemic lower amino acid and a chiral auxiliary an heterocycle is built up, that is CH-acidic adjacent to the potential amino group and that contains two sites susceptible to hydrolysis. (2) An electrophile is introduced diastereoselectively via the anion of the heterocycle. (3) Subsequently the heterocycle is cleaved by

hydrolysis to liberate the chiral auxiliary and the new optically active amino acid.

In this strategy the heterocycle merely serves as a vehicle to construct finally an acyclic molecule with the proper structure and proper configuration. It makes use of the obvious fact, that an heterocyclic intermediate is necessarily more rigid than its open chain analog, hence a higher degree of asymmetric induction can be expected.

This communication describes the asymmetric synthesis of α -alkenyl glycines of type **10** and of type **23** (or their methyl esters **21**, respectively). These non-proteinogenic amino acids deserve attention, because they are potential "suicide inhibitors"⁷ of certain pyridoxal phosphate depending enzymes. The asymmetric syntheses described here are based on the fact, that the lithiated bis-lactim ether **6** of cyclo(L-val-gly) reacts both with carbonyl compounds^{8,9} and with thioketones with exceptionally high asymmetric induction (de = diastereomeric excess).

α -Alkenyl glycines of type 10 from 6 and 2-[(dimethyl t-butyl)silyl]alkanals 2

2-[(Dimethyl t-butyl)silyl]alkanals **2** are masked alkenyl groups and allow the electrophilic introduction of alkenyl groups into carbanions **1**, according to the following scheme.¹⁰

The silylated aldehydes **2** add to the lithiated bis-lactim ether **6**—obtained from the bislactim ether **5** of cyclo(L-val-gly) and butyllithium—to give (after protonation) the aldol-type addition products **7a–c**. Like other aldehydes or ketones,^{8,9} the incoming aldehyde **2** enters *trans* to the isopropyl group at the

2	a	b	c
R ¹	H	H	Me
R ²	H	Me	Me

(Si): *tert*-butyl(dimethyl)silyl

chiral inducing center C-6, i.e. the (3*R*) configuration is induced. The asymmetric induction is >95%; only (3*R*)-isomers are detectable in the ¹H- or ¹³C-NMR-spectrum. The enantioface selection at the CO group of 2 is less pronounced (see de at C-1'). Very likely the (3*R*,1'*S*)-diastereomers are formed predominantly.⁸

7	2	R ¹	R ²	de(%) C-3	de(%) C-1'
a	H	H	H	>95	58
b	H	H	Me	>95	0
c	Me	H	Me	>95	25

On acid hydrolysis—two equivalents of 0.25 N HCl at room temp—the products 7 are cleaved at the two imino ether groups liberating L-valine methyl ester 8 and methyl (2*R*)-2-amino-4-(dimethyl *t*-butyl)silyl-3-hydroxy-alkanoates 9. The esters 8 and 9 are separable by chromatography. However, the rather thermolabile 9c suffers retroaldol cleavage and uncontrolled silanol elimination during hydrolysis and chromatography. Probably, due to ground state strain, all 4,4-dialkylated substituted esters 9 will be unstable.

The esters 9a and b were converted into the (*R*)-α-alkenyl glycines 10a and b by refluxing with

5 N HCl. Alternative methods tried for silanol elimination 9→10 gave less good results. (*R*)-(-)-vinyl glycine 10a was obtained with [α]_D²⁰ = -81.1° (c = 0.2; H₂O). As the rotation indicates¹¹ 10a has the (*R*)-configuration and is of rather high optical purity, although—if the reported rotation of [α]_D²⁰ = -93.8° (c = 1.5; H₂O)¹¹ for optically pure (*R*)-10a is correct—some racemisation has taken place in going from 7a to 10a. *R*-(-)-Propen-1-yl glycine 10b¹² was obtained as a 1:1-*E/Z*-mixture with [α]_D²⁰ = -153.8° (c = 1.2; H₂O).

The 2-(dimethyl *t*-butyl)silyl aldehydes 2a and b were prepared according to lit.¹⁰ The aldehyde 2c was obtained by the following route. 2-[(dimethyl *t*-butyl)silyl]propyliden-cyclohexylimine 11 was treated with *t*-butyllithium to give the Li derivative 12, which on addition of methyl iodide furnished 13. This on hydrolysis gave cyclohexylamine and the aldehyde 2c.

Synthesis of α-alkenyl glycine methyl esters of type 21 and α-alkenyl glycines of type 23

Addition of thioketones 14 to the lithiated bis-lactim ester 6

As studied so far, thioketones 14 react with the lithiated bis-lactim ether 6 to give in good chemical yields the addition products 15. Due to our experience in the field we can safely assume, that—like other electrophiles such as ketones and aldehydes^{8,9} or alkyl halides⁶—also thioketones 14 enter in *trans* position to the iso-propyl group at C-6 of 6, i.e. that the (3*S*)-configuration is induced at C-3 in 15. Moreover, the coupling constant *J*_{3-H/6-H} in the ¹H-NMR-spectrum of 16 is *ca* 1–3 Hz, typical for a *trans* relation between H-3 and H-6 assuming a boat shape of the diaza-hexadienyl heterocycle.¹³ The *de*-values of 15 exceed 95%. For instance, the ¹H-NMR spectrum of 16b, obtained from 15b by *S*-

Table 1. Compounds 16

16	R ¹	R ²	Yield %	Config. C-3	de (%) C-3
a	Et	Me	76	S	>95
b	—(CH ₂) ₄ —		40	S	97
c	nPr	Et	36	S	>95
d	Me	Me	52	S	>95

a) (3*S*,1*S*):(3*S*,1*R*) ≈ 1:1

methylation, was carefully scrutinized and a *ca* 60:1 diastereomer ratio was found (tantamount to *de ca* 97%). As for the 16*a,c* and *d* (Table 1) only (3*S*)-diastereomers were detectable in the ¹H- and ¹³C-NMR spectrum.†

Usually, metallation of 5 to give 6 and addition of the thioketone 14 were performed at -70° (in THF). In one experiment with pentane-3-thione both reactions were carried out at room temp in hexane, whereby 16*a* was obtained with *de ca* 69%. Hence, the reaction 6 + 14 → 15 should be performed at low temps.

Hofmann olefins 18 by elimination of methylthiol from the compounds 16. Attempts to replace the methylthio group in 16 by hydrogen by the usual Raney-Ni treatment¹⁴ gave a surprising result. Nearly quantitatively, methylthiol elimination took place with virtually complete regioselectivity to give the Hofmann olefins 18. Olefin formation during a Raney-Ni treatment of thioethers has previously been observed occasionally, but only as a minor side reaction.¹⁴ The Ni was prepared from a commercial alloy¹⁵ according to a standard procedure.¹⁶ An excess of Raney-Ni was used and the mixture refluxed

in ethanol for *ca* 3 hr. As for 18*a*, 16*a* with SBZ instead of SMe was employed. With 16*d* the regio isomer 19 was formed as the main product besides the two *E/Z*-isomers of 18*d*.

The *E/Z*-assignments are based on the ¹H-NMR spectra of the isomers. In one of the diastereomers the 3-H suffers an upfield shift of $\Delta\delta \approx 0.5$ ppm. Obviously, in this isomer the 3-H is located within the shielding cone of the double bond. Models indicate, that this is more likely for the *Z*- than for the *E*-isomer.

Noteworthy, according to preliminary results, 3-methyl-thio amino acid esters of type 20 also undergo methylthiol elimination to give β,γ -unsaturated amino acid esters of type 21 when treated with Raney-Ni.

On hydrogenation (Pd/C, ethanol, 1 at H₂, room temp) 18*a* afforded (3*R*,6*S*)-2,5-dimethoxy-3-(1'-ethylprop-1'-yl)-6-isopropyl-3,6-dihydropyrazine (yield: 90%), the precursor of (*R*)-2-amino-3-ethyl-pentanoic acid.

Hydrolysis of 18; (*R*)- α -alkenyl glycine methyl esters 21 and α -alkenyl glycines 23

Hydrolysis of the bis-lactim ethers 18 proceeds under very mild conditions—two equivalents of 0.1 N HCl, room temp—to give methyl L-Valinate 8 and the (*R*)- α -alkenyl glycine methyl esters 21. L-Valinate 8 could be separated (as forerun) by fractional bulb-to-bulb distillation. Alternatively, it should be possible to remove the valinate 8 by chromatography. The (*R*)-amino acid esters 21 obtained were essentially enantiomerically pure; only the (*R*)-enantiomers were detectable in the ¹H-NMR spectrum [Eu(hfc)₃ as chiral shift reagent]. The 19/18*d* mixture gave a mixture of 22 and 21*d*.

On further hydrolysis—2 N HCl 90°—the esters 21*a* and *b* afforded (*R*)- α -2-amino-3-ethyl-pentenoic acid 23*a* (*E/Z*-mixture) and (*R*)-(cyclohex-1'-en-1'-yl)glycine 23*b* respectively.

†After preparation of the manuscript, a capillary-GC analysis of 16*a* revealed a diastereomer ration of *ca* 98:2, tantamount to *ca* 96% *de*.

A soln of 0.66 g (11 mmol) glacial AcOH in 5 ml THF was added. After warming to room temp the solvent was evaporated *in vacuo* and the residue shaken with 30 ml ether and 20 ml water. After separation, the water layer was extracted twice with 10 ml ether. The ether layers were dried with MgSO_4 . Usual workup. The crude compounds **7** were bulb-to-bulb distilled.

(3*R*,6*S*,1'*R*,*S*) - 2,5 - Dimethoxy - 3 - [1' - hydroxy - 2' - (dimethyl *t* - butyl)silyl - eth - 1' - yl] - 6 - isopropyl - 3,6 - dihydropyrazine (**7a**)

Compound **5** (1.75 g; 9.5 mmol) and **2a** (1.50 g; 9.5 mmol). Workup after 3 hr, yield: 2.46 g (76%) **7a**; b.p. 120–130°/0.2 torr; (3*R*,6*S*,1'*S*): (3*R*,6*S*,1'*R*) *ca* 3.8; no (3*S*)-diastereomers detectable in the ^{13}C -NMR spectrum. De at C-3 > 95%.

IR(film): ν = 3500–3200 (OH), 1690 cm^{-1} (N=C). ^1H -NMR (CDCl_3): δ = –0.04 and –0.01 (2s; 6H, $\text{Si}(\text{CH}_3)_2$), 0.67 and 0.99 (2d, J = 7 Hz; 6H, $\text{CH}(\text{CH}_3)_2$), 0.80 (s; 9H, $\text{C}(\text{CH}_3)_3$), 3*R*,6*S*,1'*R*), 0.84 (s; 9H, $\text{C}(\text{CH}_3)_3$), 3*R*,6*S*,1'*S*), 0.91 (d, J = 7 Hz; 2H, SiCH_2), 2.21 (dsp, J = 3 and 7 Hz; $\text{CH}(\text{CH}_3)_2$), 2.00–2.50 br.; 1H, OH), 3.63 and 3.66 (2s; 6H, OCH_3), 3.86 and 3.93 (2dd, J = 3 Hz, 3J = 3 Hz; 2H, 3- and 6-H), 4.11 (br. dt, J = 3 and 7 Hz; 1H, CHOH). ^{13}C -NMR (CDCl_3): (3*R*,6*S*,1'*S*): δ = –5.89 and –4.95 ($\text{Si}(\text{CH}_3)_2$), 16.56 and 17.66 ($\text{CH}_2\text{Si}(\text{CH}_3)_2$), 16.78 and 19.04 ($\text{CH}(\text{CH}_3)_2$), 26.50 ($\text{C}(\text{CH}_3)_3$), 31.98 ($\text{CH}(\text{CH}_3)_2$), 52.50 and 52.55 (OCH_3), 60.99 and 61.52 (3- and 6-CH), 70.81 (C–OH), 162.29 and 165.41 (C=N). (3*R*,6*S*,1'*R*): δ = –6.07 and –4.65 ($\text{Si}(\text{CH}_3)_2$), 15.11 and 17.66 ($\text{CH}_2\text{Si}(\text{CH}_3)_2$), 16.74 and 19.04 ($\text{CH}(\text{CH}_3)_2$), 26.50 ($\text{C}(\text{CH}_3)_3$), 32.24 ($\text{CH}(\text{CH}_3)_2$), 52.29 and 52.63 (OCH_3), 61.21 and 61.25 (3- and 6-CH), 69.52 (C–OH), 162.02 and 165.04 (C=N). (Found: C, 59.74; H, 9.85. Calc. for $\text{C}_{17}\text{H}_{34}\text{N}_2\text{O}_3\text{Si}$ (342.6): C, 59.61; H, 10.00%.)

(3*R*,6*S*,1'*R*,*S*,2'*R*,*S*) - 2,5 - Dimethoxy - 3 - [1' - hydroxy - 2' - (dimethyl *t* - butyl)silyl - prop - 1' - yl] - 6 - isopropyl - 3,6 - dihydropyrazine (**7b**)

Compound **5** (0.37 g; 2 mmol) and **2b** (0.34 g; 2 mmol). Workup after 4 hr, yield: 0.57 g (80%) **7b**; b.p. 110–120°/0.1 torr; (3*R*,6*S*,1'*S*,2'*S*): (3*R*,6*S*,1'*S*,2'*R*): (3*R*,6*S*,1'*R*,2'*S*): (3*R*,6*S*,1'*R*,2'*R*) = 2.6:2.6:1:1; no (3*S*)-diastereomers detectable in the ^{13}C -NMR spectrum.

IR(film): ν = 3400–3200 (OH), 1685 cm^{-1} (N=C). ^1H -NMR (CDCl_3): δ = –0.06, –0.04, –0.03 and 0.01 (4s; 24H, $\text{Si}(\text{CH}_3)_2$), 0.64, 0.66, 0.98 and 1.00 (4d, J = 7 Hz; 36H, $\text{CH}(\text{CH}_3)_2$ and CH_3CHSi), 0.81, 0.82, 0.85 and 0.87 (4s; 36H, $\text{C}(\text{CH}_3)_3$), 1.16–1.50 (m; 4H, CH_2CHSi), 2.00–2.30 (m; 4H, $\text{CH}(\text{CH}_3)_2$), 3.63, 3.65 and 3.66 (3s; 24H, OCH_3), 3.74–4.12 (m; 16H, 3- and 6-CH, CHOH). ^{13}C -NMR (CDCl_3): (3*R*,6*S*,1'*S*,2'*R*,*S*): δ = 27.12 and 27.63 ($\text{C}(\text{CH}_3)_3$), 31.77 and 31.94 ($\text{CH}(\text{CH}_3)_2$), 73.96 and 74.70 (C–OH), 162.93, 163.06, 165.18 and 166.33 (C=N); (3*R*,6*S*,1'*R*,2'*R*,*S*): δ = 26.34 and 27.02 ($\text{C}(\text{CH}_3)_3$), 31.08 and 31.63 ($\text{CH}(\text{CH}_3)_2$), 70.83 and 73.05 (C–OH), 162.43, 163.46, 163.84 and 165.71 (C=N). (Found: C, 61.02; H, 10.18. Calc. for $\text{C}_{18}\text{H}_{36}\text{N}_2\text{O}_3\text{Si}$ (356.6): C, 60.63; H, 10.18%.)

(3*R*,6*S*,1'*R*,*S*) - 2,5 - Dimethoxy - 3 - [1' - hydroxy - 2' - methyl - 2' - (dimethyl *t* - butyl)silyl - prop - 1' - yl] - 6 - isopropyl - 3,6 - dihydropyrazine (**7c**)

Compound **5** (0.18 g; 1 mmol) and **2c** (0.19 g; 1 mmol), yield: 0.26 g (70%) **7c**; b.p. 140–150°/0.1 torr; (3*R*,6*S*,1'*S*): (3*R*,6*S*,1'*R*) *ca* 1.7:1; no 3*S*-diastereomers detectable in the ^{13}C -NMR spectrum.

IR(film): ν = 3500–3200 (OH), 1685 cm^{-1} (N=C). (3*R*,6*S*,1'*S*): ^1H -NMR (CDCl_3): δ = –0.06 (s; 6H, $\text{Si}(\text{CH}_3)_2$), 0.67 and 0.97 (2d, J = 7 Hz; 6H, $\text{CH}(\text{CH}_3)_2$), 0.93 (s; 9H, $\text{C}(\text{CH}_3)_3$), 1.03 and 1.05 (2s; 6H, $(\text{CH}_2)_2\text{CSi}$), 1.54 (br.s; 1H, OH), 2.06–2.34 (m; 1H, $\text{CH}(\text{CH}_3)_2$), 3.62 and 3.66 (2s; 6H, OCH_3), 3.84–3.97 and 4.02–4.16 (2m; 2H, 3- and 6-CH), 4.13 (d, J = 3 Hz; 1H, CHOH). ^{13}C -NMR (CDCl_3): δ = –6.12 and –5.61 ($\text{Si}(\text{CH}_3)_2$), 16.97 and 19.00 ($\text{CH}(\text{CH}_3)_2$), 19.00 (CH_3), 20.27 and 22.19 (CH_3), 28.53

($\text{C}(\text{CH}_3)_3$), 29.25 (C–Si), 32.07 ($\text{CH}(\text{CH}_3)_2$), 52.50 and 52.67 (OCH_3), 60.53 and 60.72 (3- and 6-CH), 74.99 (C–OH), 162.95 and 165.85 (C=N). (3*R*,6*S*,1'*R*): ^1H -NMR (CDCl_3): δ = –0.02 (s; 6H, $\text{Si}(\text{CH}_3)_2$), 0.61 and 1.00 (2d, J = 7 Hz; 6H, $\text{CH}(\text{CH}_3)_2$), 0.92 (s; 9H, $\text{C}(\text{CH}_3)_3$), 3.64 and 3.65 (2s; 6H, OCH_3), 4.19 (d, J = 3 Hz; 1H, CHOH). ^{13}C -NMR (CDCl_3): δ = –6.22 and –5.51 ($\text{Si}(\text{CH}_3)_2$), 16.53 and 19.48 ($\text{CH}(\text{CH}_3)_2$), 19.00 ($\text{C}(\text{CH}_3)_3$), 19.10 and 21.77 (CH_3), 28.53 ($\text{C}(\text{CH}_3)_3$), 30.23 (C–Si), 31.46 ($\text{CH}(\text{CH}_3)_2$), 52.05 and 52.82 (OCH_3), 59.91 and 60.95 (3- and 6-CH), 76.15 (C–OH), 162.30 and 163.19 (C=N). (Found: C, 61.39; H, 10.14. Calc. for $\text{C}_{19}\text{H}_{38}\text{N}_2\text{O}_3\text{Si}$ (370.6): C, 61.58; H, 10.34%.)

Compounds 9; hydrolysis of compounds **7** (General Procedure)

The suspension of 2 mmol **7** in 16 ml (4 mmol) 0.25 N HCl was stirred for 12 hr at room temp. The mixture was extracted with ether which was discharged. The solvent was concentrated *in vacuo* (bath temp 40–60°) to 1–2 ml, 10 ml ether were added and then with shaking conc ammonia till pH 8–10. The ether layer was separated, the water layer saturated with NaCl and three times extracted with 5–10 ml ether. The combined ether layer was dried with MgSO_4 . Usual workup. The residual mixture of **8** and **9** was chromatographed on silica gel (low pressure) with ether–acetone 4:1. **8** was eluted before **9**.

Methyl(2*R*,3*S*,*R*) - 2 - amino - 3 - hydroxy - 4 - (dimethyl *t* - butyl)silyl - butanoate (**9a**)

Compound **7a** (0.68 g; 2 mmol), yield, 0.26 g (53%) **9a**; R_f = 0.66 (R_f for **8** = 0.15); **9a** can be distilled without decomposition; b.p. 110–120°/0.1 torr (bulb-to-bulb); (2*R*,3*S*): (2*R*,3*R*) *ca* 2.4 (^1H -NMR spectroscopically with $\text{Eu}(\text{fod})_3$).

IR(film): ν = 3550–3150 (NH_2 and OH), 1740 (C=O), 1640–1680 cm^{-1} (NH_2). ^1H -NMR (CDCl_3): δ = 0.03 and 0.06 (2s; 6H, $\text{Si}(\text{CH}_3)_2$), 0.78–0.91 (m; 2H, SiCH_2), 0.86 (s; 9H, $\text{C}(\text{CH}_3)_3$), 2.31–2.55 (br.s; 3H, NH_2 and OH), 3.23–3.35 (m; 1H, 2-H), 3.68 (s; 3H, OCH_3), 3.80–4.03 (m; 1H, 3-H). (Found: C, 53.41; H, 10.11. Calc. for $\text{C}_{11}\text{H}_{22}\text{NO}_3\text{Si}$ (247.4): C, 53.40; H, 10.19%.)

Methyl(2*R*,3*S*,*R*,4*R*,*S*)2 - amino - 3 - hydroxy - 4 - (dimethyl *t* - butyl)silyl pentanoate (**9b**)

Compound **7b** (0.57 g; 1.6 mmol) and 0.25 N HCl (12.8 ml; 3.2 mmol), yield: 0.24 g (58%) **9b**; R_f = 0.51 (R_f for **8** = 0.15); **9b** can be distilled without decomposition; b.p. 140–150°/0.1 torr (bulb-to-bulb distillation); (2*R*,3*S*,4*R*): (2*R*,3*S*,4*S*): (2*R*,3*R*,4*R*): (2*R*,3*R*,4*S*) *ca* 2.1:2.1:1:1 (^1H -NMR spectroscopically determined, *t*-Bu and OMe-signals).

IR(film): ν = 3500–3100 (NH_2 and OH), 1740 (C=O), 1630–1670 cm^{-1} (NH_2). ^1H -NMR (CDCl_3): δ = 0.01, 0.03, 0.05 and 0.06 (4s; 12H, $\text{Si}(\text{CH}_3)_2$), 0.87 and 0.88 (2s; 18H, $\text{C}(\text{CH}_3)_3$), 2*R*,3*R*,4*R*,*S*), 0.90 and 0.92 (2s; 18H, $\text{C}(\text{CH}_3)_3$), 2*R*,3*S*,4*R*,*S*), 0.98 (d, J = 7 Hz; 6H, CH_2CH), 1.06–1.39 (m; 2H, CH_2CH), 1.96–2.14 (br.s; 6H, NH_2 and OH), 3.22–3.91 (m; 4H, 2- and 3-H), 3.70 and 3.71 (2s; 6H, OCH_3), 2*R*,3*R*,4*R*,*S*), 3.73 (s; 6H, OCH_3), 2*R*,3*S*,4*R*,*S*). (Found: C, 55.46; H, 10.42. Calc. for $\text{C}_{12}\text{H}_{27}\text{NO}_3\text{Si}$ (261.4): C, 55.13; H, 10.41%.)

(*R*)- α -Alkenyl glycines **10** from **9** (General Procedure)

The soln of 1 mmol **9** in 10 ml 5 N HCl was refluxed for 1 hr. The solvent was evaporated *in vacuo*, the residue dried for 30 min at 30–40°/0.1 torr. 5–10 ml EtOH were added, the mixture was refluxed, 2–3 ml propene oxide were added and reflux continued for 15 min. After cooling to 0°, the precipitated **10** was isolated by suction, washed on the funnel with ether and dried *in vacuo* over P_2O_{10} .

(*R*)-(–)-Vinyl glycine (**10a**)

Compound **9a** (0.20 g), yield: *ca* 50 mg (62%) **10a**;

$[\alpha]_D^{20} = -81.1^\circ$ ($c = 0.2$, H_2O) [$lit^{11}[\alpha]_D^{20} = -93.8^\circ$ ($c = 1.5$, H_2O)].

1H -NMR(D_2O): $\delta = 4.27$ (br.d, $J = 7$ Hz, $^1J < 1$ Hz; 1H, α -H), 5.46 (br.dd, $J = 2$ Hz, $J_{trans} = 17$ Hz, $^1J < 1$ Hz; 1H, *trans*-HC=CH-CH), 5.47 (br.dd, $^2J = 2$ Hz, $J_{cis} = 10$ Hz, $^1J < 1$ Hz; 1H, *cis*-HC=CH-CH), 5.99 (ddd, $J = 7$ Hz, $J_{cis} = 10$ Hz, $J_{trans} = 17$ Hz; 1H, $H_2C=CH-CH$). Found: C, 47.64; H, 7.17. Calc for $C_4H_7NO_2$ (101.1): C, 47.52; H, 6.98%.

(R)-(-)-E/Z-2-Amino-3-pentenoic acid (α -propenyl glycine) (**10b**)

Compound **9b** (0.13 g; 0.5 mmol), yield, *ca* 40 mg (70%) **10b**; $[\alpha]_D^{20} = -153.8^\circ$ ($c = 1.2$, H_2O); $[\alpha]_D^{20} = -152.3^\circ$ ($c = 0.4$, 1 N HCl). *E:Z* ca 1:1.

1H -NMR(D_2O): $\delta = 1.73$ and 1.76 (2dd, $J = 7$ Hz, $^1J = 1.5$ Hz; je 3H, CH_3 von *E*- and *Z*-**10b**), 4.15 (br.d, $J = 8$ Hz, $^1J < 1$ Hz; 1H, α -H), 5.20–6.16 (m; 2H, $CH=CH$). (Found: C, 52.60; H, 7.73. Calc for $C_5H_9NO_2$ (115.1): C, 52.16; H, 7.88%.)

(R)- α -Alkenyl glycine methyl esters **21** and (R)- α -alkenyl glycines **23** from the bis-lactim ether **5**

Compounds **16** (General Procedure). A 1.55 N soln of BuLi (2.7 ml) in hexane (4.2 mmol BuLi) were added (syringe!) at -70° to a stirred soln of 0.74 g (4 mmol) bis-lactim ether **5**²⁰ in 10 ml dry THF. After 10 min (formation of **6**) the soln of 4.2 mmol thioketone (freshly prepared and distilled, see below) in 5 ml THF was added at -70° and stirring continued for 12 hr at -70° . A soln of 0.6 g (4.2 mmol) MeI [in one case 0.72 g (4.2 mmol) benzyl bromide] in 5 ml THF was added. After warming, stirring was continued at room temp for 40 hr. The solvent was evaporated *in vacuo*, the residue shaken with 15 ml ether and 20 ml water, the layers separated and the water layer extracted twice with 10 ml ether. The combined ether soln was dried with $MgSO_4$ with usual workup. The crude compounds **16** were bulb-to-bulb distilled *in vacuo*.

(3S,6S)-2,5-Dimethoxy-6-isopropyl-3-(1'-methylthio-1'-ethylprop-1'-yl)-3,6-dihydropyrazine (**16a**)

Pentane-2-thione (0.43 g; 4.2 mmol), yield: 0.91 g (76%) **16a**; b.p. 160–170°/10 torr. Only one diastereomer detectable in the 1H - and ^{13}C -NMR spectrum. *De* > 95%.

IR(film): $\nu = 1695\text{ cm}^{-1}$ (C=N). 1H -NMR($CDCl_3$): $\delta = 0.67$ and 1.10 (2d, $J = 7$ Hz; 6H, $CH(CH_3)_2$), 0.96 (t, $J = 7$ Hz; 3H, CH_2CH_3), 0.97 (t, $J = 7$ Hz; 3H, CH_2CH_3), 1.70 (q; $J = 7$ Hz; 2H, CH_2), 1.74 (q; $J = 7$ Hz; 2H, CH_2), 1.86 (s; 3H, S- CH_3), 2.39 (dsp, $J = 3$ and 7 Hz; 1H, $CH(CH_3)_2$), 3.66 and 3.71 (2s; 6H, 2- and 5-OCH₃), 4.07 (dd, $J = 3$ Hz, $^1J = 3$ Hz; 1H, 6-H), 4.13 (d, $^1J = 3$ Hz; 1H, 3-H). ^{13}C -NMR($CDCl_3$): $\delta = 8.05$, 8.37 and 11.83, 16.43 ($CH(CH_3)_2$ and CH_2-CH_3), 19.41 (S- CH_3), 26.06 and 26.22 (CH_2), 30.35 ($CH(CH_3)_2$), 52.02 and 52.51 (OCH₃), 57.94 (C-S), 60.63 and 61.27 (3- and 6-CH), 162.65 and 165.23 (C=N). High resolution MS: (Found: 300.1871. Calc for $C_{15}H_{28}N_2O_2S$; 300.1865).

(3S,6S)-3-(1'-Benzylthio-1'-ethylprop-1'-yl)-2,5-dimethoxy-6-isopropyl-3,6-dihydropyrazine (**16a**, SBz instead of SMe)

Pentane-2-thione (0.43 g; 4.2 mmol). After 12 hr stirring at -70° addition of 0.72 g (4.2 mmol) benzyl bromide; reflux for 12 hr, yield: 1.07 g (71%) **16a** (SBz instead of SMe); b.p. 120–130°/0.01 torr. Only one diastereomer detectable in the 1H -NMR spectrum. *De* > 95%.

IR(film): $\nu = 1690\text{ cm}^{-1}$ (C=N). 1H -NMR($CDCl_3$): $\delta = 0.67$ and 1.14 (2d, $J = 6.5$ Hz; 6H, $CH(CH_3)_2$), 0.98 (t, $J = 7$ Hz; 3H, CH_2-CH_3), 1.02 (t, $J = 7$ Hz; 3H, CH_2-CH_3), 1.74 (q, $J = 7$ Hz; 2H CH_2-CH_3), 1.77 (q, $J = 7$ Hz; 2H CH_2-CH_3), 2.39 (dsp, $J = 1$ und 6 Hz; 1H, $CH(CH_3)_2$), 3.54 (s; 2H, CH_2-Ph), 3.66 and 3.71 (2s; 6H, 2- and 5-OCH₃). (Found: C, 67.77; H, 8.59. Calc for $C_{21}H_{32}N_2O_2S$ (376.6): C, 66.98; H, 8.57%.)

(3S,6S)-2,5-Dimethoxy-6-isopropyl-3-(1'-methylthio-cyclohex-1'-yl)-3,6-dihydropyrazine (**16b**)

Cyclohexanethione (0.48 g; 4.2 mmol), yield: 0.5 g (40%) (**16b**); b.p. 170°/8 torr.

Careful examination of the 1H -NMR spectrum revealed a diastereomer ratio of *ca* 60:1 (*de ca* 97%) (SMe signals at $\delta = 1.88$ and 1.89).

IR(film): $\nu = 1685\text{ cm}^{-1}$ (C=N). 1H -NMR($CDCl_3$): (3S,6S)-diastereomer: $\delta = 0.64$ and 1.09 (2d, $J = 6.5$ Hz; 6H, $CH(CH_3)_2$), 1.4–2.0 (m; 10H, C_6H_{10}), 1.88 (s; 3H, S- CH_3), 2.35 (dsp, $J = 1.5$ and 6.5 Hz; 1H, $CH(CH_3)_2$), 3.65 and 3.68 (2s; 6H, 2- and 5-OCH₃), 4.03 and 4.05 (3-H and 6-H, $J = 1.5$ Hz, $^1J \leq 1$ Hz). (3R,6S)-diastereomer: 1.89 (s; 3H, S- CH_3). High resolution MS: (Found: 312.1872. Calc for $C_{16}H_{28}N_2O_2S$; 312.1865).

(3S,6S)-2,5-Dimethoxy-6-isopropyl-3-(1'-methylthio-1'-propylbut-1'-yl)-3,6-dihydropyrazine (**16c**)

Heptane-4-thione (0.55 g; 4.2 mmol), yield, 0.47 g (36%) **16c**; b.p. 170–180°/10 torr. Only one diastereomer detectable in the 1H -NMR spectrum. *De* > 95%.

IR(film): $\nu = 1690\text{ cm}^{-1}$ (C=N). 1H -NMR($CDCl_3$): $\delta = 0.68$ and 1.12 (2d, $J = 6.5$ Hz; 6H, $CH(CH_3)_2$), 0.94 and 0.97 (2t, $J = 4$ Hz; 6H, $[(CH_2)_2-CH_3]_2$), 1.23–1.76 (m; 8H, $[(CH_2)_2-CH_3]_2$), 1.88 (s; 3H, S- CH_3), 2.37 (dsp, $J = 3$ and 6.5 Hz; 1H, $CH(CH_3)_2$), 3.70 and 3.72 (2s; 6H, 2- and 5-OCH₃), 4.09 (dd, $J = 3$ Hz, $^1J = 3$ Hz; 1H, 6-H), 4.14 (d, $J = 3$ Hz; 1H, 3-H). High resolution MS: Found: 328.2185. Calc for $C_{17}H_{32}N_2O_2S$ (328.5): 328.2177.

(3S,6S,1'S,R)-2,5-Dimethoxy-6-isopropyl-3-(1'-methylthio-1'-methyl-prop-1'-yl)-3,6-dihydropyrazine (**16d**)

Butane-2-thione (0.37 g; 4.2 mmol), yield: 0.60 g (52%) **16d**; b.p. 140°/8 torr. (3S,6S,1'S): (3S,6S,1'R) = 1.05:1 (determined by 1H -NMR, signals at $\delta = 1.33$ and 1.37 or 1.94 and 1.97). (3R,6S)-diastereomers were not detectable in the 1H -NMR spectrum, even not with Eu(fod)₃.

IR(film): $\nu = 1690\text{ cm}^{-1}$ (C=N). 1H -NMR($CDCl_3$): (3S,6S,1'S)-diastereomer: $\delta = 0.68$ and 1.13 (2d, $j = 7$ Hz; 6H, $CH(CH_3)_2$), 1.05 (t, $J = 7$ Hz; 3H, CH_2-CH_3), 1.33 (s; 3H, 1'- $C-CH_3$), 1.78 (q, $J = 7$ Hz; CH_2-CH_3), 1.97 (s; S- CH_3), 2.38 (dsp, $J = 3$ and 7 Hz; $CH(CH_3)_2$), 3.71 and 3.73 (2s; 2- and 5-OCH₃), 4.01 (dd, $J = 3$ Hz, $^1J = 3$ Hz; 1H, 6-H), 4.12 (d, $J = 3$ Hz; 1H, 3-H); (3S,6S,1'R)-diastereomer: 1.37 (s; 3H, 1'- $C-CH_3$), 1.75 (q, $J = 7$ Hz; CH_2-CH_3), 1.94 s; CH_3). High resolution MS: Found: 286.1715. Calc for $C_{14}H_{26}N_2O_2S$; 286.1709.

Compounds **18** and **19**; methylthiol elimination with Raney-Ni (General Procedure)

A suspension of Raney-Ni prepared from 6 g Raney-Ni alloy¹⁵ according to *lit*¹⁶ in 20 ml 75% EtOH was added to the soln of 3 mmol **16** in 30 ml 75% EtOH and the mixture refluxed for 3 hr. The catalyst was filtered off when still hot and washed three times with hot EtOH. The solvent was evaporated *in vacuo* and the residual compounds **18** bulb-to-bulb distilled.

E/Z-(3R,6S)-2,5-Dimethoxy-3-(1'-ethylprop-1'-en-1'-yl)-6-isopropyl-3,6-dihydropyrazine (**18a**)

Compound **16a** (SBz instead of SMe) (1.13 g; 3 mmol), yield, 0.67 g (88%) *E/Z*-**18a** (*E:Z* = 1.3:1); b.p. 130°/1 torr.

IR(film): $\nu = 1620$ (C=C), 1695 cm^{-1} (C=N). 1H -NMR($CDCl_3$): *Z*-**18a**: $\delta = 0.72$ and 1.09 (2d; 6H, $CH(CH_3)_2$), 0.96 (t, $J = 7$ Hz; 3H, CH_2-CH_3), 1.68 (d, $J = 7$ Hz; 3H, C=C- CH_3), 1.95 (q, $J = 7$ Hz; 2H, C=C- CH_2), 2.32 (dsp, $J = 3$ and 7 Hz; 1H, $CH(CH_3)_2$), 3.66 and 3.67 (2s; 6H, 2- and 5-OCH₃), 3.96 (dd, $J = 3$ Hz, $^1J = 3$ Hz; 1H, 6-H), 4.48 (d, $J = 3$ Hz; 1H, 3-H), 5.47 (q, $J = 7$ Hz; 1H, C=C-H). *E*-**18a**: $\delta = 0.76$ and 1.09 (2d; 6H, $CH(CH_3)_2$), 0.98 (t, $J = 7$ Hz; 3H, CH_2-CH_3), 1.78 (d, $J = 7$ Hz; 3H, C=C- CH_3),

1.95 (q, $J = 7$ Hz; 2H, $C=C-CH_3$), 2.32 (dsp, $J = 3$ and 7 Hz; 1H, $CH(CH_3)_2$), 3.70 and 3.71 (2s; 6H, 2- and 5- OCH_3), 4.04 (dd, $J = 3$ Hz, $^3J = 4$ Hz; 1H, 6-H), 5.02 (d, $J = 4$ Hz; 1H, 3-H), 5.47 (q, $J = 7$ Hz; 1H, $C=C-H$). High resolution MS: Found: 252.1838. Calc for $C_{14}H_{24}N_2O_2$: 252.1832.

(3R,6S)-3-(1'-Cyclohex-1'-en-1'-yl)-2,5-dimethoxy-6-isopropyl-3,6-dihydropyrazine (18b)

Compound 16b (0.94 g; 3 mmol), yield, 0.74 g (93%) 18b; b.p. 160°/10 torr.

IR(film): $\nu = 1620$ ($C=C$), 1685 cm^{-1} ($C=N$). 1H -NMR ($CDCl_3$): $\delta = 0.72$ and 1.09 (2d, $J = 7$ Hz; 6H, $CH(CH_3)_2$), 1.40–2.20 (m; 8H, $-(CH_2)_4$), 2.32 (dsp, $J = 3$ and 7 Hz; 1H, $CH(CH_3)_2$), 3.68 and 3.70 (2s; 6H, 2- and 5- OCH_3), 3.93 (dd, $J = 3$ Hz, $^3J = 3$ Hz; 1H, 6-H), 4.41 (d, $J = 3$ Hz; 1H, 3-H), 5.68 (m; 1H, $C=C-H$). High resolution MS: Found: 264.1838. Calc for $C_{15}H_{25}N_2O_2$: 264.1832.

E/Z-(3R,6S)-2,5-Dimethoxy-6-isopropyl-3-(1'-propylbut-1'-en-1'-yl)-3,6-dihydropyrazine (18c)

Compound 16c (0.99 g; 3 mmol), yield, 0.71 g (84%) 18c; b.p. 180°/8 torr.

IR(film): $\nu = 1620$ ($C=C$), 1690 cm^{-1} ($C=N$). 1H -NMR ($CDCl_3$): Z-18c: $\delta = 0.72$ and 1.09 (2d, $J = 7$ Hz; 6H, $CH(CH_3)_2$), 0.85 and 1.02 (2t, $J = 7$ Hz; 6H, CH_2-CH_3 and $(CH_2)_2-CH_3$), 1.30 (m; 2H, $-CH_2-CH_2-CH_3$), 1.86 and 2.20 (2m; 4H, $-CH_2-CH_2-CH_3$ and $-CH_2-CH_3$), 2.30 (dsp, $J = 3$ and 7 Hz; 1H, $CH(CH_3)_2$), 3.69 and 3.73 (2s; 6H, 2- and 5- OCH_3), 4.08 (dd, $J = 3$ Hz, $^3J = 3.5$ Hz; 1H, 6-H), 4.43 (d, $J = 3.5$ Hz; 1H, 3-H), 5.36 (t, $J = 7.5$ Hz; 1H, $C=C-H$). E-18c: $\delta = 0.71$ and 1.07 (2d, $J = 7$ Hz; 6H, $CH(CH_3)_2$), 0.92 and 0.97 (2t, $J = 7$ Hz; 6H, CH_2-CH_3 and $(CH_2)_2-CH_3$), 2.45 (m; 2H, $-CH_2-CH_2-CH_3$), 1.88 and 2.23 (2m; 4H, $-CH_2-CH_2-CH_3$ and $-CH_2-CH_3$), 2.30 (dsp, $J = 3$ and 7 Hz; 1H, $CH(CH_3)_2$), 3.66 and 3.69 (2s; 6H, 2- and 5- OCH_3), 3.98 (dd, $J = 3$ Hz, $^3J = 3.5$ Hz; 1H, 6-H), 4.93 (d, $J = 3.5$ Hz; 1H, 3-H), 5.36 (t, $J = 7.5$ Hz; 1H, $C=C-H$). High resolution MS: Found: 280.2151. Calc for $C_{16}H_{28}N_2O_2$: 280.2144.

(3R,6S)-2,5-Dimethoxy-3-(1'-ethyl-ethenyl)-6-isopropyl-3,6-dihydropyrazine (19) and Z/E-(3R,6S)-2,5-dimethoxy-6-isopropyl-3-(1'-methyl-prop-1'-en-1'-yl)-3,6-dihydropyrazine (18d)

Compound 16d (0.86 g; 3 mmol), yield, 0.61 g (86%) 74:16:10-mixture of 19:Z-18d:E-18d; b.p. 130–140°/8 torr.

IR(film): $\delta = 1635$ ($C=C$), 1695 cm^{-1} ($C=N$). 1H -NMR($CDCl_3$): 19: $\delta = 0.66$ and 0.71 (2d, $J = 7$ Hz; 6H, $CH(CH_3)_2$), 1.04 (t, $J = 7$ Hz; 3H, CH_2-CH_3), 1.68 (q, $J = 7$ Hz; 2H, CH_2-CH_3), 2.30 (dsp, $J = 3$ and 7 Hz; 1H, $CH(CH_3)_2$), 3.70 (2s; 6H, 2- and 5- OCH_3), 3.70–4.20 (m; 4H, 3-H, 6-H, $=CH_2$). Z-18d: $\delta = 0.72$ and 0.95 (2d, $J = 7$ Hz; 6H, $CH(CH_3)_2$), 1.52 (d, $^3J = 1$ Hz; 3H, $HC-C-CH_3$), 1.77 (d, $J = 7$ Hz; 3H, $=CH(CH_3)$), 2.02 (dsp, $J = 3$ and 7 Hz; 1H, $CH(CH_3)_2$), 3.68 (2s; 6H, 2- and 5- OCH_3), 4.42 (d, $J = 3$ Hz; 1H, 3-H), 5.54 (q, $J = 7$ Hz, $^3J = 1$ Hz; 1H, $C=C-H$). E-18d: $\delta = 0.74$ and 0.90 (2d, $J = 7$ Hz; 6H, $CH(CH_3)_2$), 1.48 (d, $^3J = 1$ Hz; 3H, $HC-C-CH_3$), 1.79 (d, $J = 7$ Hz; 3H, $=CHCH_3$), 2.02 (dsp, $J = 3$ and 7 Hz; 1H, $CH(CH_3)_2$), 3.68 (2s; 6H, 2- and 5- OCH_3), 4.97 (d, $J = 3$ Hz; 1H, 3-H), 5.54 (q, $J = 7$ Hz, $^3J = 1$ Hz; 1H, $C=C-H$). (Found: C, 65.48; H, 9.53. Calc for $C_{13}H_{22}N_2O_2$ (238.3): C, 65.52; H, 9.30%).

Compounds 18a–c and 25a–c from the sulfonium salts 24 (General Procedure)

MeI (1.42 g; 10 mmol) was added to a soln of 3 mmol 16 in 15 ml dry acetonitrile. The mixture was refluxed for 12–14 hr. The solvent was evaporated *in vacuo* and the residue shaken with 20 ml water and 10 ml ether. The layers were separated and the water layer extracted twice with 10 ml ether. The combined ether layer was dried with $MgSO_4$ with usual workup. The crude products 18/25 were

purified by bulb-to-bulb distillation. The product ratios were analyzed by 1H -NMR spectroscopy.

Compounds 18a and (6S)-25a: 0.9 g (3 mmol) 16a, yield, 0.55 g (73%) 88:12-mixture of Z/E-18a and 25a. Z:E = 2.5; b.p. 130°/1 torr.

Spectra of Z/E-18a see above.

Compound 25a: 1H -NMR($CDCl_3$): $\delta = 0.84$ (d, $J = 7$ Hz; 3H, $CH_3-CH-CH_3$), 3.75 and 3.76 (2s; 6H, 2- and 5- OCH_3).

Compound 18b and (6S)-3-Cyclohexylidene-2,5-dimethoxy-6-isopropyl-3,6-dihydropyrazine (25b)

Compound 16b (0.94 g; 3 mmol), yield 0.53 g (67%) 90:10-mixture of 18b and 25b; b.p. 160°/10 torr.

IR(film): $\nu = 1620$ ($C=C$), 1685 cm^{-1} ($C=N$). 1H -NMR($CDCl_3$) for 25b (for 18b see above): $\delta = 0.93$ and 1.22 (2d, $CH(CH_3)_2$), 3.74 and 3.76 (2s, 2- and 5- OMe). High resolution MS: Found: 264.1838. Calc for $C_{15}H_{24}N_2O_2$: 264.1832.

Compound 18c and (6S)-2,5-Dimethoxy-6-isopropyl-3-(1'-propylbutylidene)-3,6-dihydropyrazine (25c)

Compound 16d (0.99 g; 3 mmol), yield, 0.67 g (80%) 87:13-mixture of E/Z-18c and 25c. Z:E = 2.

IR(film): $\nu = 1620$ ($C=C$), 1690 cm^{-1} ($C=N$). 1H -NMR($CDCl_3$) for 25c (for 18c see above): $\delta = 0.70$ (d, $-CH-CH_3$), 3.64 and 3.65 (2s, 2- and 5- OMe). High resolution MS: Found: 280.2151. Calc for $C_{16}H_{28}N_2O_2$: 280.2144.

(R)- α -Alkenyl glycine methyl esters 21 and α -alkenyl glycines 23

Hydrolysis of 18 or 18/25-mixtures (General Procedure). 2 mmol 18 (or the mixture of 18 and 25, see above) were suspended in 40 ml 0.1 N HCl and vigorously stirred for 40–60 hr at room temp. The aqueous layer was extracted twice with 30 ml ether which was discarded (in case of the 18/25-mixture it contained 26). The solvent was evaporated *in vacuo* (bath temp 40–60°); ca 15 ml water and 20 ml ether were added to the residue and with vigorous shaking conc. ammonia till pH 8. The ether layer was separated and the water layer extracted three times with 15 ml ether. The combined ether layer was dried with $MgSO_4$ with usual workup. The crude mixture of 21 and valinate 8 was bulb-to-bulb distilled and 8 removed as forerun.

E/Z-Methyl(R)-2-amino-3-ethyl-3-pentenoate (21a)

Compound 18a (0.5 g; 2 mmol). Workup after 40 hr, yield: 0.23 g (73%) E/Z-21a (E:Z = 1.3:1); b.p. 130°/5 torr; $[\alpha]_D^{20} = -150.6^\circ$ ($c = 1.0$, $CHCl_3$).

IR(film): $\nu = 1740$ ($C=O$), 3200–3500 cm^{-1} (NH_2). 1H -NMR($CDCl_3$): E-21a: $\delta = 1.04$ (t, $J = 7$ Hz; 3H, CH_2-CH_3), 1.75 (d, $J = 6$ Hz; 3H, $=CHCH_3$), 1.93 (s; 2H, NH_2), 2.16 (q, $J = 7$ Hz; 2H, CH_2-CH_3), 3.72 (s; 3H, OCH_3), 4.49 (s; 1H, $\alpha-H$), 5.49 (q, $J = 6$ Hz; 1H, $=CH$). Z-21a: $\delta = 1.02$ (t, $J = 7$ Hz; 3H, CH_2-CH_3), 1.67 (d, $J = 6$ Hz; 3H, $=CHCH_3$), 1.93 (s; 2H, NH_2), 2.16 (q, $J = 7$ Hz; 2H, CH_2-CH_3), 3.72 (s; 3H, OCH_3), 3.99 (s; 1H, $\alpha-H$), 5.49 (q, $J = 6$ Hz, $=CH$). (Found: C, 60.98; H, 9.92. Calc for $C_8H_{15}NO_2$ (157.2): C, 61.12; H, 9.62%).

Methyl(R)-(cyclohex-1'-ene-1-yl)glycinate (21b)

Compound 18b (0.53 g; 2 mmol). Workup after 40 hr, yield: 0.26 g (76%) 21b; b.p. 140–150°/10 torr. Enantiomerically pure by 1H -NMR. $[\alpha]_D^{20} = -89.4^\circ$ ($c = 1.0$, $CHCl_3$).

IR(film): $\nu = 1740$ ($C=O$), 3200–3500 cm^{-1} (NH_2). 1H -NMR($CDCl_3$): $\delta = 1.50$ –2.20 (m; 8H, $-(CH_2)_4-$), 1.78 (s; 2H, NH_2), 3.72 (s; 3H, OCH_3), 3.93 (s; 1H, $\alpha-H$), 5.70 (m; 1H, $=CH$). (Found: C, 64.00; H, 8.95. Calc for $C_9H_{15}NO_2$ (169.2): C, 63.88; H, 8.93%).

Methyl(R)-2-amino-3-ethyl-3-butenate (22) and Z/E-methyl(R)-2-amino-3-methyl-3-pentenoate (21d)

A mixture of 19 and E/Z-18d (0.48 g; 2 mmol). Workup

after 40 hr, yield, 0.18 g (64%) 74:16:10-mixture of **22** and **Z/E-21d**; b.p. 110°/8 torr. Contaminated with ca 20% of **8**.

IR(film): $\nu = 1730$ (C=O), 3200–3400 cm^{-1} (NH₂). ¹H-NMR(CDCl₃): **22**: $\delta = 0.84$ (t, J = 7 Hz; 3H, CH₂-CH₃), 1.65 (s; 2H, NH₂), 1.75 (q, J = 7 Hz; 2H, CH₂-CH₃), 3.65 (s; 3H, OCH₃), 3.70–4.00 (m; 3H, =CH₂, α -H). **Z-21d**: $\delta = 1.60$ (s; 3H, CH₃), 1.65 (s; 2H, NH₂), 1.70 (d, J = 6 Hz; 3H, =CHCH₃), 3.65 (s; 3H, OCH₃), 3.84 (s; 1H, α -H), 5.39 (q, J = 6 Hz; 1H, =CH). **E-21d**: $\delta = 1.60$ (s; 3H, CH₃), 1.65 (s; 2H, NH₂), 1.70 (d, J = 6 Hz; 3H, =CHCH₃), 3.65 (s; 3H, OCH₃), 4.35 (s; 1H, α -H), 5.39 (q, J = 6 Hz; 1H, =CH).

α -Alkenyl glycines **23** (General Procedure)

Compound **21** (1 mmol) was refluxed for 3 hr with 10 ml 2 N HCl. The soln was evaporated *in vacuo* to dryness, 5 ml dry EtOH and 2 ml propylene oxide added and the mixture refluxed for 20 min. After cooling to room temp the precipitated **23** was isolated by suction, washed with some cold acetone and dried several days at room temp at 10 torr.

Z/E-(R)-2-Amino-3-ethyl-3-pentenoic acid (**23a**)

Compound *E/Z*-**21a** (0.16 g; 1 mmol) (obtained from the sulfonium salt **25**), yield, 0.10 g (72%) *Z/E*-**23a** (*Z*:*E* = 2.5); m.p. 164° (dec) [α]_D²⁰ = -113.1° (*c* = 1.0; 1 N HCl).

IR(KBr): $\nu = 1620$ (CO₂⁻), 3050 cm^{-1} (NH₃⁺, br.). ¹H-NMR(D₆-DMSO/D₂O): **Z-23a**: $\delta = 0.95$ (t, J = 7.5 Hz; 3H, CH₂CH₃), 1.64 (d, J = 6.5 Hz; 3H, =CHCH₃), 2.08 (q, J = 7.5 Hz; 2H, CH₂-CH₃), 3.85 (s; 1H, α -H), 5.54 (q, J = 6.5 Hz; 1H, =CH). **E-23a**: $\delta = 0.98$ (t, J = 7.5 Hz; 3H, CH₂-CH₃), 1.67 (d, J = 6.5 Hz; 3H, =CHCH₃), 2.08 (q, J = 7.5 Hz; 2H, CH₂-CH₃), 5.54 (q, J = 6.5 Hz; 1H, =CH). (Found: C, 56.42; H, 9.19. Calc for C₁₇H₁₃NO₃ (143.2): C, 58.72; H, 9.15%).

(R)-(Cyclohex-1'-en-1'-yl)glycine (**23b**)

Compound **21b** (0.17 g; 1 mmol), yield, 0.12 g (76%) **23b**; m.p. 154° (dec) [α]_D²⁰ = -96.8° (*c* = 0.5; 1 N HCl).

IR(KBr): $\nu = 1600$ (CO₂⁻), 3050 cm^{-1} (NH₃⁺, br.). ¹H-NMR(D₂O): $\delta = 1.25$ –2.15 (m; 8H, -(CH₂)₄-), 4.03 (s; 1H, α -H), 5.83 (m; 1H, =CH). (Found: C, 61.57; H, 8.39. Calc for C₈H₁₃NO₂ (155.2): C, 61.91; H, 8.44%).

Compound **21a** from the bis-lactim ether **27**

(3*S*)-2,5-Dimethoxy-3-(3,4-dimethoxybenzyl)-3-methyl-3,6-dihydropyrazine (**27**) was obtained according to lit.¹⁸

(3*S*,6*S*)-2,5-Dimethoxy-3-(3,4-dimethoxybenzyl)-3-methyl-6-(1'-methylthio-1'-ethylprop-1'-yl)-3,6-dihydropyrazine (**30b**)

A soln of 4.2 mmol BuLi (2.7 ml of a 1.55 N soln in hexane) was added at -70° to the soln of 1.23 g (4 mmol) **27**. After stirring for 20 min, the soln of 0.43 g (4.2 mmol) pentane-3-thione in 5 ml dry THF was added. After 12 hr stirring at -70° the soln of 0.60 g (4 mmol) MeI was added. After stirring at room temp for 40 hr workup as described above for **18**, yield: 0.91 g (54%) **30b**; b.p. 180°/0.005 torr (bulb-to-bulb distillation).

IR(film): $\nu = 1696$ cm^{-1} (C=N). ¹H-NMR(CDCl₃): $\delta = 0.87$ and 0.92 (2t, J = 7 Hz; 6H, (CH₂-CH₃)₂), 1.60 (s; 3H, 3-CH₃), 1.62 and 1.74 (2q, J = 7 Hz; 4H, (CH₂-CH₃)₂), 1.82 (s; 3H, S-CH₃), 2.85 and 2.98 (AB, C₆H₃-CH₂; J = 13 Hz), 3.59 (s; 1H, 6-H), 3.65 and 3.66 (2s; 6H, 2- and 5-OCH₃), 3.80 and 3.83 (2s; 6H, C₆H₃-(OCH₃)₂), 6.61 (m; 3H, C₆H₃). High resolution MS: Found: 422.2239. Calc for C₂₂H₂₄N₂O₄S: 422.2231.

Z/E-(3*S*,6*R*)-2,5-Dimethoxy-3-(3,4-dimethoxybenzyl)-6-(1'-ethylprop-1'-en-1'-yl)-3-methyl-3,6-dihydropyrazine (**31**) and (3*S*)-2,5-dimethoxy-3-(3,4-dimethoxybenzyl)-6-(1'-ethyl-propyliden)-3-methyl-3,6-dihydropyrazine (**32**)

Elimination via the sulfonium salt as described above.

Compound **30b** (1.2 g; 3 mmol); reflux with MeI in acetonitrile for 14 hr, yield: 0.97 g (86%) *Z/E*-**31** and **32** (76:24); *Z*:*E* = 4; b.p. 130–140°/0.04 torr (bulb-to-bulb distillation).

IR(film): $\nu = 1620$ (C=C), 1680 cm^{-1} (C=N). **Z-31**: ¹H-NMR(CDCl₃): $\delta = 0.89$ (t, J = 7 Hz; 3H, CH₂-CH₃), 1.52 (s; 3H, 3-CH₃), 1.61 (d, J = 6 Hz; 3H, C=C-CH₃), 1.86 (q, J = 7 Hz; 2H, C=C-CH₂), 2.83 and 2.97 (AB, C₆H₃-CH₂; J = 13 Hz), 3.65 (2s; 6H, 2- and 5-OCH₃), 3.79 and 3.80 (2s; 6H, C₆H₃-(OCH₃)₂), 4.05 (s; 1H, 6-H), 5.33 (q, J = 6 Hz; 1H, C=C-H), 6.59 (m; 3H, C₆H₃). **E-31**: ¹H-NMR(CDCl₃): $\delta = 0.87$ (t, J = 7 Hz; 3H, CH₂-CH₃), 1.49 (s; 3H, 3-CH₃), 1.59 (d, J = 6 Hz; 3H, C=C-CH₃), 1.86 (q, J = 7 Hz; 2H, C=C-CH₂), 2.83 and 2.97 (AB, C₆H₃-CH₂; J = 13 Hz), 3.65 (2s; 6H, 2- and 5-OCH₃), 3.77 and 3.78 (2s; 6H, C₆H₃-(OCH₃)₂), 4.05 (s; 1H, 6-H), 5.27 (q, J = 6 Hz; 1H, C=C-H), 6.59 (m; 3H, C₆H₃). Compound **32**: ¹H-NMR(CDCl₃): $\delta = 0.79$ and 0.94 (2t, J = 7 Hz; 6H, -(CH₂CH₃)₂), 1.46 (s; 3H, 3-CH₃), 3.82 (2s; 6H, C₆H₃-(OCH₃)₂). High resolution MS: Found: 374.2206. Calc for C₂₁H₂₀N₂O₄: 374.2198.

Compound **21a**. Hydrolysis of 0.75 g (2 mmol) **31/32**-mixture according to hydrolysis **18**–**21** (see above) for 60 hr, yield: 0.16 g (52%) *Z/E*-**21a** (*Z*:*E* = 4); [α]_D²⁰ = -104° (*c* = 1.0, CHCl₃); physical data and spectra see above.

Thioketones **14**

The thioketones **14** were basically prepared according to lit.²¹, the procedure modified according to lit.²²: Through the soln of the dimethyl ketals in AcOH, that contained some hydrochinone and a catalytic amount of conc H₂SO₄, a rather strong stream of H₂S was passed. For further information about the preparation of thioketones, see lit.²³

Acknowledgements—We gratefully acknowledge the support of this research by the Deutsche Forschungsgemeinschaft, the Fonds der Chemischen Industrie, the BASF AG, the Degussa AG and by Boehringer, Mannheim.

REFERENCES

- For part XXI, see: U. Schöllkopf, U. Busse, R. Kilger and P. Lehr, *Synthesis* in press (1984).
- G. Nass, K. Poralla and H. Zähler, *Naturwissenschaften* **58**, 603 (1971).
- W. Trowitzsch and H. Sahn, *Z. Naturforschung, Teil C* **32**, 78 (1977).
- J. D. Morrison and H. S. Mosher, *Asymmetric Organic Reactions*. Prentice-Hall, Englewood Cliffs, New Jersey (1971).
- Recent reviews on asymmetric syntheses, including amino acids. J. W. ApSimon and R. P. Seguin, *Tetrahedron* **35**, 2797 (1979); K. Weinges and B. Stemmler, *Recent Develop. Chem. Nat. Carbon Compd.* **7**, 91 (1976); D. Valentine, Jr. and J. W. Scott, *Synthesis* 329 (1978); H. B. Kagan, *Pure and Appl. Chem.* **43**, 401 (1975); H. B. Kagan and J. C. Fiaud, *Topics in Stereochemistry* **10**, 175 (1978).
- Reviews: U. Schöllkopf, *Topics in Current Chemistry* **109**, (1983); *Tetrahedron* **39**, 2085 (1983).
- R. R. Rando, *Acc. Chem. Res.* **8**, 281 (1975); R. H. Abeles and A. L. Maycock, *Ibid.* **9**, 313 (1976); R. H. Abeles, *Pure and Appl. Chem.* **53**, 149 (1981); C. Walsh, *Tetrahedron* **38**, 871 (1982).
- U. Schöllkopf, U. Groth, M.-R. Gull and J. Nozulak, *Liebigs Ann. Chem.* 1133 (1983); *cf. lit.* 6.
- U. Schöllkopf, J. Nozulak and U. Groth, *Synthesis* 868 (1982).
- P. F. Hudrlik and A. K. Kulkarni, *J. Am. Chem. Soc.* **103**, 6251 (1981).
- P. Friis, P. Helboe and P. O. Larsen, *Acta Chem. Scand. B* **28**, 317 (1974); for a synthesis of L-vinyl glycine from L-methionine *cf.* A. Afzali-Ardakani and H. Rapoport, *J. Org. Chem.* **45**, 4817 (1980); for a synthesis of D,L-vinyl

- glycine *cf* J. E. Baldwin, S. B. Haber, C. Hoskins and L. I. Kruse, *J. Org. Chem.* **42**, 1239 (1977).
- ¹²For syntheses of D,L-, (Z)- and (E)-2-Amino-3-pentenoic acid *cf* M. Johnston, R. Raines, M. Chang, N. Esaki, K. Soda and C. Walsh, *Biochemistry* **20**, 4325 (1981).
- ¹³An X-ray analysis of the addition product of **6** and benzaldehyde revealed, that the dihydropyrazine is a flat boat; E. Egert, J. Nozulak and U. Schöllkopf, unpublished. With regard to the coupling constants in boat-shaped heterocycles *cf* H. Günther, *NMR-Spektroskopie*. 1. Aufl., S. 113 ff. Georg Thieme Verlag, Stuttgart (1973).
- ¹⁴G. R. Pettit and E. E. van Tamelen, *Organic Reactions* **12**, 356 (1962); H. Hauptmann and W. F. Walter, *Chem. Rev.* **62**, 347 (1962).
- ¹⁵Obtained from Fluka AG, CH-9470 Buchs, Switzerland.
- ¹⁶Organikum, Deutscher Verlag der Wissenschaften, Berlin (1976).
- ¹⁷The racemic mixtures of **21b** and **23b** have been described; *cf* K. Nunami, M. Suzuki and N. Yoneda, *J. Chem. Soc. Perkin I*, 2224 (1979); R. D. Allan, *Austral. J. Chem.* **32**, 2507 (1979).
- ¹⁸U. Schöllkopf, W. Hartwig, K.-H. Pospischil and H. Kehne, *Synthesis* 966 (1981).
- ¹⁹We thank Dr. Dick and Dr. Lettenbauer of Boehringer, Mannheim, for a generous gift of (S)-O,O-dimethyl- α -methyl dopa.
- ²⁰U. Schöllkopf, U. Groth and C. Deng, *Angew. Chem. Int. Ed. Engl.* **20**, 798 (1981).
- ²¹R. Mayer and H. Berthold, *Chem. Ber.* **96**, 3096 (1963).
- ²²K. Madawinata, Dissertation Univ. Göttingen (1977).
- ²³S. Bleisch and R. Mayer, *Chem. Ber.* **99**, 1771 (1966); R. Mayer, J. Morgenstern and J. Fabian, *Angew. Chem.* **76**, 157 (1964); *Ibid.* Int. Ed. Engl. **3**, 277 (1964); B. S. Pedersen, S. Scheibye, N. H. Nilsson and S. O. Lawesson, *Bull. Soc. Chim. Belg.* **87**, 223 (1978); E. Schaumann, U. Wriede and G. Rüther, *Angew. Chem.* **95**, 52 (1983); *Ibid.* Int. Ed. Engl. **22**, 55 (1983).