Total Synthesis of the Angucyclinone Antibiotic (+)-Rubiginone B₂¹

Aris Kalogerakis, Ulrich Groth*

Fachbereich Chemie, Universität Konstanz, Fach M-720, Universitätsstraße 10, 78457 Konstanz, Germany

Fax +49(7531)884155; E-mail: ulrich.groth@uni-konstanz.de

Received 30 June 2003

Abstract: A new chiral synthesis of (+)-rubiginone B₂ is reported. The intramolecular cobalt-mediated [2+2+2]-cycloaddition of a triyne precursor, synthesized from (+)-citronellal, afforded a chiral anthracene, which led after a two-step oxidation to the angucyclinone antibiotic.

Key words: angucyclinone antibiotics, benz[*a*]anthraquinones, natural products, catalysis, cobalt, cycloaddition

The rubiginones (Figure 1),² isolated from the strain of *Streptomyces griseorubiginosus*, exhibit potential vincristine-induced cytotoxicity against multi-drug-resistant tumor cells. They feature the typical benz[*a*]anthraquinone structure of the angucyclinone antibiotics, a growing class of secondary metabolites with remarkable biological properties.³ While several racemic syntheses of these angucyclinones have been achieved,⁴ only a few asymmetric total syntheses have been reported.⁵ The only known total synthesis of (+)-rubiginone B₂ has been reported by Carreño and co-workers based on the Diels–Alder reaction between an enantiopure substituted naphthoquinone and a racemic vinylcyclohexene through the kinetic resolution of the diene. The natural product was synthesized with 80% ee and 9% overall yield for the longest sequence.⁶

Figure 1

We have recently reported a new method¹ for the construction of the angucyclinone core via an intramolecular cobalt-mediated [2+2+2]-cycloaddition of a triyne.⁷ Herein, we describe a new asymmetric access to the angucyclinone antibiotic (+)-rubiginone B₂ **12**. Starting from commercially available *R*-(+)-citronellal ($[\alpha]_D^{20} +13.0$,

neat, 92% ee)⁸ we have synthesized the chiral diyne **6** (Scheme 1). By employing our previously reported method¹ we used **6** for the synthesis of the triyne **9**, which after cyclization and two-step oxidation led to (+)-rubiginone B₂ **12** (Scheme 2).

Our synthesis began with a Corey–Fuchs olefination⁹ of (+)-citronellal **1**. The resulting dibromoalkene **2** was used without further purification and converted to the protected alkyne **3** by treatment with *n*-BuLi and then TMS-Cl¹⁰ (76% yield, 2 steps). Ozonolysis of **3** (Me₂S workup) afforded **4** in quantitative yield, which was transformed without purification via another Corey–Fuchs olefination into the dibromoalkene **5**. Treatment of **5** with *n*-BuLi and hydrolysis of the resulting lithiated acetylene gave the diyne **6** (74% yield, 3 steps).

Scheme 1 Reaction Conditions: (a) 2 equiv CBr₄, 4 equiv PPh₃, CH₂Cl₂, 0 °C, 30 min, then **1**, 25 °C, 2 h. (b) i. 2 equiv *n*-BuLi, THF, –80 °C, 1 h, ii. 2 equiv TMS-Cl, –80 °C to r.t., 18 h (76%, 2 steps). (c) i. O₃, CH₂Cl₂, –80 °C, 15 min, ii. Me₂S, AcOH, –80 °C to r.t., 18 h. (d) 2 equiv CBr₄, 4 equiv PPh₃, CH₂Cl₂, 0 °C, 30 min, then **4**, 25 °C, 2 h. (e) i. 2 equiv *n*-BuLi, THF, –80 °C, 1 h, ii. H₂O, –80 °C to r.t., 2 h (74%, 3 steps).

After addition of the lithiated diyne **6** to the benzaldehyde **7**¹ and deprotection of the triple bonds of the resulting triyne **8** with NH₄F,¹¹ the hydroxy group of the obtained triyne was protected with TBDMSOTf.¹² Cyclization of **9** with 10% CpCo(CO)₂¹³ in refluxing toluene under irradiation (tungsten-lamp) gave the anthracene **10** in 74% yield. By using 5% CpCo(CO)₂ **10** was obtained in 61% yield. The use of the expensive CpCo(ethene)₂¹⁴ for the cyclization (see ref.¹) of the triyne **9** was not necessary. Oxidation with [Ag(Py)₂]MnO₄¹⁵ led to the anthraquinone **11** (62% yield), which was converted via photooxidation^{4c} to (+)-rubiginone B₂ **12** (67% yield).¹⁶ The spectroscopic data of synthetic (+)-rubiginone B₂ **12** coincided with those of the

natural compound.² The structure of **12** was proven by two-dimensional NMR experiments (ROESY-, H,H-COSY-spectra, see the H,H-correlations in Figure 2). Additionally, the observed optical rotation value $[\alpha]_{\text{D}}^{25} +71.64$ (*c* 0.275, CHCl₃, 92% ee) agreed satisfactory with the reported value $[\alpha]_{\text{D}}^{25} +78.00$ (*c* 0.5, CHCl₃) for the natural product.

Scheme 2 Reaction Conditions: (a) *n*-BuLi, **6**, THF, -80°C , 1 h, then **7**, -80°C to -30°C , 4 h (93%), (b) i. NH₄F, Bu₄NHSO₄, CH₂Cl₂, r.t., 48 h (96%), ii. TBDMSTf, 2,6-lutidine, CH₂Cl₂, 25°C , 2 h (95%) (c) 10% CpCo(CO)₂, toluene, reflux, h ν , 4 h (74%). (d) 8 equiv [Ag(Py)₂]MnO₄, CH₂Cl₂, 25°C , 8 h (62%). (e) h ν , air, CHCl₃, 25°C , 18 h (67%).

Figure 2 H,H-correlations in ROESY- and H,H-COSY-Spectra

In summary, a new effective chiral synthesis of (+)-rubiginone B₂ has been achieved starting from commercially available *R*-(+)-citronellal in 11 steps and 15% overall yield. Applications to the asymmetric synthesis of other angucyclinone antibiotics using the reported methodology¹ are currently under investigation.

Acknowledgment

The authors are grateful to the Fonds der Chemischen Industrie and the EU-Comission, Directorate XII, for financial support. We thank Dr. H. Frauendorf, analytical laboratory of the institute of organic chemistry at the university of Göttingen, for measuring the HRMS-analysis of compound **10**.

References

- (1) Transition metal catalyzed reactions in organic synthesis, part 4. For part 3, see: Kalogerakis, A.; Groth, U. *Org. Lett.* **2003**, *6*, 843.
- (2) Oka, M.; Kamei, H.; Hamagishi, Y.; Omita, K.; Miyaki, T.; Konish, M.; Oki, T. *J. Antibiot.* **1990**, *43*, 967.
- (3) (a) Rohr, J.; Thiericke, R. *Nat. Prod. Rep.* **1992**, 103. (b) Krohn, K.; Rohr, J. *Top. Curr. Chem.* **1997**, *188*, 127.
- (4) (a) Uemura, M.; Take, K.; Minami, T.; Hayashi, Y. *Tetrahedron* **1985**, *41*, 5771. (b) Kraus, G. A.; Wu, Y. *Tetrahedron Lett.* **1991**, *32*, 3803. (c) Krohn, K.; Khanbabaee, K.; Jones, P. G.; Chrapowski, A. *Liebigs Ann. Chem.* **1994**, 471. (d) Krohn, K.; Khanbabaee, K. *Liebigs Ann. Chem.* **1994**, 1109. (e) Krohn, K.; Mcheel, J.; Zukowski, M. *Tetrahedron* **2000**, *56*, 4753. (f) Krohn, K.; Böker, N.; Flörke, U.; Freund, C. *J. Org. Chem.* **1997**, *62*, 2350. (g) Katsura, K.; Snieckus, V. *Can. J. Chem.* **1987**, *65*, 124. (h) Guingant, A.; Barreto, M. *Tetrahedron Lett.* **1987**, *28*, 3107. (i) Patil, M. G.; Borate, H. B.; Ponde, D. E.; Bhawal, B. M.; Deshpande, V. H. *Tetrahedron Lett.* **1999**, *40*, 4437. (j) Mal, D.; Roy, H. N. *J. Chem. Soc., Perkin Trans 1* **1999**, 3167. (k) Rozek, T.; Bowie, J. H.; Pyke, S. M.; Skelton, B. W.; White, A. H. *J. Chem. Soc., Perkin Trans 1* **2001**, 1826.
- (5) (a) Larsen, D. S.; O'Shea, M. D.; Brooker, S. *Chem. Commun.* **1996**, 203. (b) Caygill, G. B.; Larsen, D. S.; Brooker, S. *J. Org. Chem.* **2001**, *66*, 7427. (c) Carreño, M. C.; Ribagorda, M.; Somoza, A.; Urbano, A. *Angew. Chem., Int. Ed.* **2002**, *41*, 2755. (d) Boyd, V. A.; Sulikowski, G. A. *J. Am. Chem. Soc.* **1995**, *117*, 8472.
- (6) (a) Carreño, M. C.; Urbano, A.; Vitta, C. D. *Chem. Commun.* **1999**, 817. (b) Carreño, M. C.; Urbano, A.; Vitta, C. D. *J. Org. Chem.* **1998**, *63*, 8320. (c) Hecker, S. J.; Heathcock, C. H. *J. Org. Chem.* **1985**, *50*, 5159.
- (7) (a) Vollhardt, K. P. C. *Acc. Chem. Res.* **1977**, *10*, 1. (b) Vollhardt, K. P. C. *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 539. (c) Lecker, S. H.; Nguyen, N. H.; Vollhardt, K. P. C. *J. Am. Chem. Soc.* **1986**, *108*, 856. (d) Dosa, P. I.; Whitener, G. D.; Vollhardt, K. P. C.; Bond, A. D.; Teat, S. J. *Org. Lett.* **2002**, *4*, 2075.
- (8) (a) Purchased from Fluka. (b) Tietze, L. F.; Kiedrovski, G. V.; Fahlbusch, K.-G.; Voss, E. *Org. Synth.* **1993**, *8*, 353. (c) Chapuis, C.; Barthe, M.; Laumer, J.-Y. *Helv. Chim. Acta* **2001**, *84*, 230.
- (9) Corey, E. J.; Fuchs, P. L. *Tetrahedron Lett.* **1972**, *36*, 3769.
- (10) Hosokawa, S.; Isobe, M. *J. Org. Chem.* **1999**, *64*, 37.
- (11) Herold, P. *Helv. Chim. Acta* **1988**, *71*, 354.
- (12) Corey, E. J.; Cho, H.; Ruecker, C.; Hua, D. H. *Tetrahedron Lett.* **1981**, *36*, 3455.
- (13) **Experimental Procedure:** A solution of CpCo(CO)₂ (25.00 mg, 0.142 mmol, 10 mol%) in 2 mL toluene was added via canula to a solution of triyne **9** (0.58 g, 1.42 mmol) in 120 mL toluene and the mixture was heated to reflux and irradiated with a tungsten-lamp (osram vitalux 300 W) for 4 h. The organic phase was then concentrated in vacuo. Chromatography on silica gel (Et₂O–petroleum ether, 1:200) provided anthracene **10** (0.29 g, 1.051 mmol, 74% yield) as a white solid.

- Mp 136–138 °C. $[\alpha]_{\text{D}}^{22} +86.43$ (*c* 0.14, CHCl₃). $R_f = 0.31$ (Et₂O–petroleum ether, 1:100). IR (CCl₄): 2935–2810 [C–H (OCH₃)] cm^{−1}. ¹H NMR (600 MHz, CDCl₃): $\delta = 1.16$ (d, ³*J* = 6.6 Hz, 3 H, 3-CH₃), 1.58 (m, 1 H, H-2), 2.00 (m, 1 H, H-3), 2.14 (m, 1 H, H-2), 2.60 (dd, ²*J* = 16.5 Hz, ³*J* = 10.5 Hz, 1 H, H-4), 2.95 (dd, ²*J* = 16.5 Hz, ³*J* = 3.9 Hz, 1 H, H-4), 3.14 (m, 1 H, H-1), 3.40 (d, ²*J* = 16.5 Hz, 1 H, H-1), 4.09 (s, 3 H, OCH₃), 6.73 (d, *J*_o = 8.0 Hz, 1 H, H-9), 7.18 (d, *J*_o = 8.6 Hz, 1 H, H-5), 7.37 (t, *J*_o = 8.0 Hz, 1 H, H-10), 7.62 (d, *J*_o = 8.0 Hz, 1 H, H-11), 7.82 (d, *J*_o = 8.6 Hz, 1 H, H-6), 8.44 (s, 1 H, H-12), 8.78 (s, 1 H, H-7). ¹³C NMR (100 MHz, CDCl₃): $\delta = 21.79$ (3-CH₃), 25.83 (C-1), 28.88 (C-3), 31.35 (C-2), 39.14 (C-4), 55.45 (OCH₃), 101.34 (C-9), 120.75 (C-12), 120.83 (C-11), 121.35 (C-7), 125.06 (C-10), 126.58 (C-6), 128.05 (C-5), 124.09, 130.34, 131.42, 132.64, 133.57, 155.38 (C_{quart}-arom.). EI-MS (70 eV): *m/z* = 276 (100%, M⁺), 261 (15%, M⁺ – CH₃), 246 (7%, 261 – CH₃), 233 (77%, 261 – C₂H₄). HRMS: calcd 276.1514 for C₂₀H₂₀O, found 276.1508.
- (14) (a) Jonas, K.; Deffense, E.; Habermann, D. *Angew. Chem., Int. Ed. Engl.* **1983**, 22, 716. (b) Cammack, J. K.; Jalisatgi, S.; Matzger, A. J.; Négron, A.; Vollhardt, K. P. C. *J. Org. Chem.* **1996**, 61, 4798.
- (15) (a) Tius, M. A.; Galeno, J. G.; Gu, X.; Zaid, J. H. *J. Am. Chem. Soc.* **1991**, 113, 5775. (b) Firouzabadi, H.; Vessel, B.; Naderi, M. *Tetrahedron Lett.* **1982**, 23, 1847.
- (16) **Experimental Procedure:** Anthraquinone **11** (60.00 mg, 0.195 mmol) was dissolved in 5 mL CHCl₃ and irradiated with a tungsten-lamp (osram vitalux 300 W) over 18 h. The organic layer was then removed in vacuo. Chromatography on silica gel (petroleum ether–Et₂O, 1:2) provided (+)-rubiginone B₂ **12** (42.00 mg, 0.131 mmol, 67% yield) as a yellow solid. Mp > 262 °C (dec.). $[\alpha]_{\text{D}}^{25} +71.64$ (*c* 0.275, CHCl₃, recrystallized from petroleum ether–CHCl₃, 2:1). $R_f = 0.14$ (Et₂O–petroleum ether, 1:4). IR (CCl₄): 1673, 1677, 1708 (C=O) cm^{−1}. ¹H NMR (400 MHz, CDCl₃): $\delta = 1.20$ (d, ³*J* = 6.4 Hz, 3 H, 3-CH₃), 2.45 (m, 1 H, H-3), 2.55 (dd, ²*J* = 15.6 Hz, ³*J* = 11.0 Hz, 1 H, H-2), 2.67 (dd, ²*J* = 16.4 Hz, ³*J* = 11.0 Hz, 1 H, H-2), 2.98 (m, 2 H, H-1 and H-4), 4.04 (s, 3 H, OCH₃), 7.24 (d, *J*_o = 8.2 Hz, 1 H, H-9), 7.45 (d, *J*_o = 8.0 Hz, 1 H, H-5), 7.65 (t, *J*_o = 8.2 Hz, *J*_o = 7.8 Hz, 1 H, H-10), 7.71 (d, *J*_o = 7.8 Hz, 1 H, H-11), 8.20 (d, *J*_o = 8.0 Hz, 1 H, H-6). ¹³C NMR (150 MHz, CDCl₃): $\delta = 21.43$ (3-CH₃), 30.82 (C-3), 38.33 (C-4), 47.55 (C-2), 56.50 (OCH₃), 117.15 (C-9), 119.68 (C-11), 129.60 (C-6), 132.99 (C-5), 135.34 (C-10), 120.56, 134.98, 135.07, 137.67, 149.13, 159.81 (C_{quart}-arom.), 181.59, 184.51, 198.89 (C=O). EI-MS (70 eV): *m/z* = 320 (100%, M⁺), 305 (15%, M⁺ – CH₃), 292 (35%, M⁺ – CO), 291 (15%, M⁺ – HCO), 261 (27%, 292 – OCH₃), 233 (24%, 261 – CO).