

Short communication

When the ringing in the ears gets unbearable: Illness representations, self-instructions and adjustment to tinnitus

Manja Vollmann ^{a,*}, Natallia Kalkouskaya ^a, Berthold Langguth ^b, Margreet Scharloo ^c

^a University of Konstanz, Germany

^b Department of Psychiatry and Psychotherapy, University of Regensburg, Germany

^c Leiden University Medical Center, The Netherlands

A B S T R A C T

Objective: Chronic tinnitus can severely impair a person's quality of life. The degree of impairment, however, is not closely related to tinnitus loudness. Applying the common sense model (CSM) of self regulation of health and illness, this study investigated to what extent psychological factors, i.e. illness representations and positive/negative self instructions, are associated with the degree of tinnitus related complaints.

Methods: In this cross sectional study, 118 patients diagnosed with chronic tinnitus filled in questionnaires assessing illness representations (IPQ R), positive and negative self instructions (TRSS), and tinnitus related complaints (TQ).

Results: The regression analysis yielded a number of significant associations between illness representations and tinnitus related complaints, particularly for the IPQ R dimensions identity, consequences, coherence, and emotional representations. With regard to self instructions and tinnitus related complaints, significant effects were found only for negative self instructions. Moreover, multiple mediation analyses revealed that the effects of consequences and emotional representations on tinnitus related complaints were (partially) due to the use of negative self instructions.

Conclusion: Psychological factors are strongly related to the extent of tinnitus related complaints. The findings provide an indication of which aspects should be targeted in psychological and psychotherapeutic tinnitus treatment.

Keywords:

Tinnitus
Illness representations
Self-instructions
Adjustment
Coping
Common Sense Model (CSM) of self-regulation

Introduction

Tinnitus, defined as the perception of an internal sound in the absence of an external stimulus [1], is a widespread phenomenon in the general population [2]. Chronic tinnitus can severely impair a person's quality of life as it causes a number of psychological and somatic complaints [3–6]. For about 10–20% of affected people, tinnitus becomes an impairing condition. The degree of impairment, however, does not appear to be closely related to objective aspects of the tinnitus, such as its loudness [7,8]. This suggests that adjustment to tinnitus is likely to be influenced by psychological factors.

The Common Sense Model (CSM) of self regulation of health and illness [9] addresses how psychological factors, i.e. illness representations and coping responses, may influence adjustment to medical conditions. The CSM proposes that individuals' mental representations of an illness determine their coping responses which, in turn, have an impact on their illness adjustment. There is growing cross

sectional and prospective evidence for the usefulness of the CSM in predicting illness outcomes in a wide range of illnesses [10–12].

Previous studies have examined the impact of illness representations and coping responses on tinnitus adjustment, but only separately. The results from these studies are in line with the assumptions of the CSM. It has been found that negative illness representations, i.e. a strong illness identity, perceptions of serious consequences, and perceptions of low control, were associated with lower levels of adjustment to tinnitus [6,13–15]. With regard to coping, researchers proposed that especially cognitive processes play an important role in tinnitus adjustment [16]. Studies showed that coping strategies such as dwelling on tinnitus, catastrophizing about consequences, and masking the tinnitus were related to more psychological distress, whereas strategies such as acceptance, positive self talk, and engaging in activities were related to less psychological distress [17–20]. To our knowledge, there are no studies that have examined the postulated mediating effect of coping responses on the relationship between illness representations and adjustment in patients with chronic tinnitus.

Therefore, the aim of the present study was to extend previous research on tinnitus adjustment by jointly investigating the impact of illness representations and coping responses, i.e. positive and

negative self instructions, on tinnitus related complaints based on the assumptions of the CSM.

Method

Participants and procedure

Participants were recruited from the Tinnitus Center of the University of Regensburg, Germany. A total of 200 patients diagnosed with tinnitus were invited to participate. They were provided with information about the study and received a booklet containing the questionnaires. A total of 118 patients returned the completed booklet (response rate: 59%). The study was approved by the Ethical Committee of the University of Regensburg and informed consent was obtained from all participants.

Measures

Illness Perceptions Questionnaire (IPQ R)

Participants' cognitive and emotional representations of tinnitus were assessed by a modified German version of the Illness Perceptions Questionnaire Revised [21,6].

The first part measures *illness identity* that refers to symptoms perceived to be part of the illness. For 15 symptoms, participants indicated whether they have experienced the symptom in the course of their tinnitus, and if yes, whether they believe the symptom to be specifically related to their tinnitus.

The second part contains 7 subscales assessing *chronic timeline* (5 items), *cyclical timeline* (4 items), *consequences* (5 items), *personal control* (4 items), *treatment control* (4 items), *coherence* (5 items), and *emotional representations* (5 items). Responses were given on a 5 point rating scale. Cronbach's alpha ranged between .65 and .91, except for cyclical timeline ($\alpha = .51$) which was therefore excluded from further analyses.

The third part measures *causal attributions*. Participants rated 21 potential causes of their tinnitus. A principal component analysis with VARIMAX rotation yielded two factors that can be labeled as *internal causes* (7 items, $\alpha = .86$) and *external causes* (5 items, $\alpha = .62$).

Tinnitus Related Self Statements Scale (TRSS)

The Tinnitus Related Self Statement Scale [22] was used to assess situation specific cognitions during the occurrence of tinnitus. The TRSS consists of two subscales capturing *negative self instructions* (9 items, $\alpha = .86$) and *positive self instructions* (9 items, $\alpha = .80$).

Tinnitus Questionnaire (TQ)

The German version of the Tinnitus Questionnaire [23] was used to assess tinnitus related complaints. The TQ contains 5 subscales relating to psychological distress, intrusiveness, auditory perception difficulties, sleep disturbance, and somatic complaints. A total score that includes 42 items was calculated ($\alpha = 0.95$).

Statistics

Data were analyzed following the procedure described by Evans and Norman [11]. A regression analysis quantified the extent to which illness representations and self instructions explained variance in tinnitus related complaints. Illness representations were entered in the first step and self instructions were entered in the second step. Subsequent multiple mediation analyses [24] were conducted when the effect of an illness representation dimension in step 1 was substantially reduced (by at least 0.5 points) after the addition of self instructions in step 2 in order to test whether self instructions mediated the effect of the illness representation dimension on tinnitus related complaints.

Results

Sample characteristics

The sample consisted of 27 women and 91 men with a mean age of 55.64 years ($SD = 11.31$). The majority of participants have received some kind of tinnitus treatment in the past (94.8%). Most participants were employed (57%) or retired (36%), while only a few received illness benefits (2%) or were unemployed/homemakers (3%). The mean sum score for the overall subjective severity of tinnitus (TQ-score $M = 40.49$, $SD = 17.87$) was below the cut-off point for decompensated tinnitus, although 28 (23.7%) participants reported severe distress (TQ-score: 47–59) and 16 (13.6%) participants reported very severe distress (TQ-score: 60–84).

Explained variance in tinnitus-related complaints and mediating effects of self-instructions

The illness representations entered in the first step of the regression explained 57% of the variance in tinnitus-related complaints (see Table 1). The dimensions identity, consequences, coherence, and emotional representations showed significant associations with the outcome. The addition of the self-instructions in the second step of the regression produced a significant increment by 5% in the amount of explained variance. Negative self-instructions were significantly associated with the outcome.

As the effects of consequences and emotional representations were substantially reduced when the self-instructions were added to the regression, two multiple mediation analyses were conducted. Both analyses revealed a significant mediation effect of negative self-instructions (see Table 2), indicating that the use of negative self-instructions partially mediated the effect of consequences and fully mediated the effect of emotional representations on tinnitus-related complaints.

Discussion

The present findings confirm the assumption that tinnitus adjustment is strongly influenced by psychological factors. Thus, the way people think and feel about their tinnitus and the way they cope with their tinnitus are substantially related to their adjustment to tinnitus. Among the illness representations, the dimensions identity, consequences, coherence, and emotional representations are of special importance which is in line with previous research on tinnitus sufferers [9] and other patients [10]. Attributing many symptoms to tinnitus, the perception of serious consequences, a low sense of understanding of the tinnitus, and strong emotional representations are directly and, in part, indirectly via negative self instructions associated with elevated levels of tinnitus related complaints. With regard to coping, negative self instructions have an unfavorable effect on tinnitus adjustment which concurs with previous studies examining the effectiveness of coping strategies in tinnitus patients [17,18]. Positive self instructions were not independently associated with better adjustment. Additionally, negative self instructions mediated at

Table 1
Results of the regression analysis

Independent variables	β step 1	β step 2
Step 1: adj. $R^2 = .58$, $F(7, 103) = 22.97^{***}$		
Identity	.23**	.20**
Chronic timeline	.09	.07
Consequences ^a	.33***	.27**
Coherence	.18*	.15*
Emotional representations ^a	.28**	.14
Internal causes	.01	.04
External causes	.04	.03
Step 2: $\Delta R^2 = .05$, $F(2, 101) = 7.05^{**}$; adj. $R^2 = .63$, $F(9, 101) = 21.54^{***}$		
Negative self-instructions		.35***
Positive self-instructions		.06

Note. Only those illness representation dimensions that correlated significantly with tinnitus-related complaints were entered into the regression analysis in order to keep the number of independent variables to a minimum.

^a Mediation was subsequently tested (see Table 2).

*** $p < .001$.

** $p < .01$.

* $p < .05$.

Table 2
Results of the mediation analyses

Effect	β	BCa 95% CI	
		Lower	Upper
Independent variable: consequences			
IV – Mediator 1	.22*		
IV – Mediator 2	.25*		
Mediator 1 – DV (direct effect)	.36***		
Mediator 2 – DV (direct effect)	.06		
IV – DV (total effect)	.33***		
IV – DV (direct effect)	.27**		
Mediation mediator 1	.08 ^a	.018	.189
Mediation mediator 2	.02 ^a	.077	.011
Independent variable: emotional representations			
IV – Mediator 1	.44***		
IV – Mediator 2	.19		
Mediator 1 – DV (direct effect)	.36***		
Mediator 2 – DV (direct effect)	.06		
IV – DV (total effect)	.28**		
IV – DV (direct effect)	.14		
Mediation effect mediator 1	.16 ^a	.057	.274
Mediation effect mediator 2	.01 ^a	.064	.010

Note. Mediation analyses were calculated with consequences/emotional representations as independent variable (IV), negative and positive self-instructions as mediators 1 and 2, tinnitus-related complaints as dependent variable (DV), and the remaining dimensions of illness representations as covariates.

BCa 95% CI = bias corrected and accelerated 95% confidence interval; 5000 bootstrap samples.

^a The mediation effects were estimated by bootstrap analyses [25].

*** $p < .001$.

** $p < .01$.

* $p < .05$.

least partially the unfavorable impact of consequences and emotional representations on tinnitus adjustment. Thus, strong emotional representations and the perception of serious consequences promote the use of negative self instructions which, in turn, hinder adjustment to tinnitus.

These findings are of practical interest as they provide an indication of which aspects should mainly be targeted in psychological and psychotherapeutic interventions for tinnitus, namely, negative illness representations and catastrophizing. Patients with chronic tinnitus may benefit from information provision on and discussions about the symptoms, timeline, possible treatments, and likely consequences of tinnitus as this might foster their sense of understanding of tinnitus. It might also lead to an increase of patients' perceptions of control which, in turn, might minimize the emotional impact of tinnitus and reduce the perceived severity of consequences. Interventions developed to modify illness representations in other patient groups could generate the desired changes [26,27]. Additionally, patients should be dissuaded from negative self instructions and instructed on how to deal with the tinnitus and how to accommodate to its consequences. Previous research identified acceptance of the tinnitus and engaging in activities as effective ways of coping with tinnitus [19,20]. The suggestions derived from our findings are in line with cognitive behavioral therapy (CBT) which has been found to have positive effects on qualitative aspects of tinnitus and the management of tinnitus. CBT for tinnitus usually combines the provision of information, the modification of negative thinking, and the enhancement of effective coping [28,29]. The present findings could help to specify the areas of negative cognitions and indicate that especially negative illness perceptions and catastrophizing should be targeted in psychological tinnitus treatment. Furthermore, our findings could also be of use in the context of tinnitus counseling as the manner in which information about tinnitus and its treatment is communicated to patients influences their illness representations and their use of coping strategies.

Some limitations need to be acknowledged. Due to the cross sectional design, the direction of causality in the associations between the constructs could not be determined. Longitudinal studies are

needed to clarify this issue. Additionally, longitudinal studies could investigate the dynamic processes specified in the CSM [30]. From the various possible ways of coping with tinnitus, this study focused on two cognitive coping responses. Further research would profit from including more diverse coping strategies to get a broader insight in the interplay between illness representations, coping responses, and tinnitus related complaints.

Role of funding source

Funding was provided by the Committee for Research Funding (AFF) of the University of Konstanz, Germany to Natallia Kalkouskaya and Manja Vollmann. The AFF had no further role in study design; in collection, analysis and interpretation of the data; in the writing of the report; and in the decision to submit the paper for publication.

Conflict of interest

The authors have no competing interests to report.

References

- Lockwood AH, Salvi RJ, Burkard RF. Tinnitus. *N Engl J Med* 2002;347:904–10.
- Sanchez L. The epidemiology of tinnitus. *Audiol Med* 2004;2:8–17.
- Andersson G, Vretblad P, Larsen HC, Lyttkens L. Longitudinal follow-up of tinnitus complaints. *Arch Otolaryngol Head Neck Surg* 2001;127:175–9.
- Sanchez L, Stephens SDG. Perceived problems of tinnitus clinic clients at long-term follow up. *J Audiol Med* 2000;9:94–103.
- Langguth B, Landgrebe M, Kleinjung T, Sand GP, Hajak G. Tinnitus and depression. *World J Biol Psychiatry* 2011;12:489–500.
- Reynolds P, Gardner D, Lee R. Tinnitus and psychological morbidity: a cross-sectional study to investigate psychological morbidity in tinnitus patients and its relationship with severity of symptoms and illness perceptions. *Clin Otolaryngol Allied Sci* 2004;29:628–34.
- Andersson G, Baguley DM, McKenna L, McFerran DJ. Tinnitus: a multidisciplinary approach. London: Whurr; 2005.
- Ward LM, Baumann M. Measuring tinnitus loudness using constrained psychophysical scaling. *Am J Audiol* 2009;18:119–28.
- Leventhal H, Meyer D, Nerenz D. The common sense model of illness danger. In: Rachman S, editor. *Medical psychology*. New York: Pergamon; 1980. p. 7–30.
- Hagger MS, Orbell S. A meta-analytic review of the common-sense model of illness representations. *Psychol Health* 2003;18:141–84.
- Evans D, Norman P. Illness representations, coping and psychological adjustment to Parkinson's disease. *Psychol Health* 2009;24:1181–96.
- Förster C, Taubert S. Subjektive Krankheitstheorien und Selbstregulation von Tumorpatienten. *Z Med Psychol* 2006;15:117–27.
- Budd RJ, Pugh R. The relationship between locus of control, tinnitus severity, and emotional distress in a group of tinnitus sufferers. *J Psychosom Res* 1995;39:1015–8.
- Sirois FM, Davis CG, Morgan MS. "Learning to live with what you can't rise above": control beliefs, symptom control, and adjustment to tinnitus. *Health Psychol* 2006;25:119–23.
- von Osterhausen K, Kessler B, D'Amelio R, Delb W. Adaptivität und Maladaptivität der Krankheitsverarbeitung bei Patienten mit subjektivem chronischen Tinnitus. *Verhaltenstherapie Verhaltensmedizin* 2001;22:39–54.
- Andersson G, McKenna L. The role of cognition in tinnitus. *Acta Otolaryngol* 2006;126:39–43.
- Budd RJ, Pugh R. The relationship between coping style, tinnitus severity and emotional distress in a group of tinnitus sufferers. *Br J Health Psychol* 1996;1:219–29.
- Budd RJ, Pugh R. Tinnitus coping style and its relationship to tinnitus severity and emotional distress. *J Psychosom Res* 1996;41:327–35.
- Davis CG, Morgan MS. Finding meaning, perceiving growth, and acceptance of tinnitus. *Rehabil Psychol* 2008;53:128–38.
- Westin V, Hayes SC, Andersson G. Is it the sound or your relationship to it? The role of acceptance in predicting tinnitus impact. *Behav Res Ther* 2008;46:1259–65.
- Gaaf J, Latanzia-Bunschoten S, Sprott H, Ehler U, IPQ. Illness Perception Questionnaire (Deutsche version). In: Bengel J, Wirtz M, Zwingmann C, editors. *Diagnostische Verfahren in der Rehabilitation*. Göttingen: Hogrefe; 2008. p. 109–11.
- Weise C, Heinecke K, Rief W. Biofeedback-based behavioral treatment for chronic tinnitus: results of a randomized controlled trial. *J Consult Clin Psychol* 2008;76:1046–57.
- Goebel G, Hiller W. Tinnitus-Fragebogen. Göttingen: Hogrefe; 1998.
- Preacher KJ, Hayes AF. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav Res Methods* 2008;40:879–91.
- Shrout PE, Bolger N. Mediation in experimental and nonexperimental studies: new procedures and recommendations. *Psychol Methods* 2002;7:422–45.

- [26] Broadbent E, Ellis CJ, Thomas J, Gamble G, Petrie KJ. Further development of an illness perception intervention for myocardial infarction patients: a randomized controlled trial. *J Psychosom Res* 2009;67:17–23.
- [27] Glattacker M, Heyduck K, Meffert C. Illness beliefs, treatment beliefs and information needs as starting points for patient information — evaluation of an intervention for patients with chronic back pain. *Patient Educ Couns* 2012;86:378–89.
- [28] Martinez-Devesa P, Waddell A, Perera R, Theodoulou M. Cognitive behavioural therapy for tinnitus. *Cochrane Database Syst Rev* 2007(1) [Art. No.: CD005233].
- [29] Zachriat C, Kröner-Herwig B. Treating chronic tinnitus: comparison of cognitive-behavioural and habituation-based treatments. *Cogn Behav Ther* 2004;33:187–98.
- [30] Leventhal H, Benyamini Y, Brownlee S, Diefenbach M, Leventhal EA, Patrick-Miller L, et al. Illness representations: theoretical foundations. In: Petrie KJ, Weinman JA, editors. *Perceptions of health and illness: current research and applications*. Harwood Academic Publishers: Amsterdam, The Netherlands; 1997. p. 19–45.