

## **ansa-METALLOCENE DERIVATIVES**

### **VI \*. SYNTHESIS AND MOLECULAR STRUCTURE OF A STABLE TETRAMETHYLETHYLENE-BRIDGED CHROMOCENE CARBONYL COMPLEX, $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4)_2\text{Cr}(\text{CO})$**

HEINZ SCHWEMLEIN, LASZLO ZSOLNAI, GOTTFRIED HUTTNER and HANS H. BRINTZINGER \*\*

*Fakultät für Chemie, Universität Konstanz, Postfach 5560, 7750 Konstanz (F.R.G.)*

#### **Summary**

The tetramethylethylene-bridged chromocene carbonyl complex  $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4)_2\text{Cr}(\text{CO})$  is obtained by reaction of the Grignard compound  $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4\text{MgCl})_2 \cdot 4\text{THF}$  with  $\text{CrCl}_2 \cdot \text{THF}$  under CO. The diamagnetic carbonyl complex is the first thermally stable chromocene ligand adduct; its structure is that of a normal wedge-shaped metallocene derivative whose ring ligands span a centroid–chromium–centroid angle of  $143^\circ$ ; the molecule closely approximates to  $C_2$  symmetry with respect to the  $\text{Cr}-\text{C}\equiv\text{O}$  axis.

#### **Introduction**

While molybdocene and tungstenocene form a great variety of ligand derivatives, coordinative adducts of chromocene are rare, and none of those reported are thermally stable. Wilkinson, Cotton and Birmingham observed the formation of  $\text{NH}_3$  adducts of  $(\text{C}_5\text{H}_5)_2\text{Cr}$  in liquid ammonia [2]; at ambient temperatures  $(\text{C}_5\text{H}_5)_2\text{Cr}$  is regenerated from these complexes [3]. Later, we found that  $(\text{C}_5\text{H}_5)_2\text{Cr}$  is quantitatively converted to the diamagnetic adduct  $(\text{C}_5\text{H}_5)_2\text{Cr}(\text{CO})$  by reaction with CO in pentane at  $-78^\circ\text{C}$  [4]; at room temperature, however, the CO dissociation pressure rises above 100 mbar. Finally, Chisholm et al. reported that  $\text{F}_3\text{CC}\equiv\text{CCF}_3$  to form a diamagnetic  $(\text{C}_5\text{H}_5)_2\text{Cr}$  adduct at temperatures below  $-30^\circ\text{C}$ ; at room temperature, this complex is converted into an incompletely characterized paramagnetic material [5]. We now report the synthesis and molecular structure of a

\* For Part V see ref. 1.

\*\* To whom correspondence should be addressed.

surprisingly stable carbonyl derivative of an *ansa*-chromocene containing an interannular tetramethylethylene bridge.

## Results and discussion

When the Grignard compound  $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4\text{MgCl})_2 \cdot 4\text{THF}$ , obtained by reductive coupling of 6,6-dimethyl fulvene with  $\text{Mg}/\text{CCl}_4$  [1] was treated with an equimolar quantity of  $\text{CrCl}_2 \cdot \text{THF}$  in THF in an attempt to obtain the tetramethylethylene-bridged chromocene  $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4)_2\text{Cr}$ , only an insoluble, very air-sensitive, red product, probably of polymeric nature was obtained. However, when the reaction is carried out under an atmosphere of CO, a dark red solution is formed, from which, by evaporation of solvent, extraction with pentane and subsequent recrystallization, dark red needles of  $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4)_2\text{Cr}(\text{CO})$  are obtained in ca. 50% yield. This product can be further purified by sublimation in vacuo, which is, however, accompanied by loss of material due to partial decomposition to an unsublimable residue.

The product is characterized by an IR spectrum with a strong  $\nu(\text{CO})$  absorption at  $1905\text{ cm}^{-1}$  (pentane solution). In the reversibly formed unbridged CO adduct  $(\text{C}_5\text{H}_5)_2\text{Cr}(\text{CO})$  the  $\nu(\text{CO})$  absorption appears at  $1900\text{ cm}^{-1}$  [4]. This close spectral coincidence indicates a very similar degree of back donation in the two CO adducts; the increased stability of the bridged derivative would thus have to be ascribed primarily to steric reasons, i.e. to the fixation of a deformed ligand framework by the interannular bridge.

In its  $^1\text{H}$  NMR spectrum, the ring protons of  $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4)_2\text{Cr}(\text{CO})$  give rise to the AA'XX' pair of triplets normally observed in *ansa*-metallocenes. The chemical shifts for these resonances, at 3.60 and 4.38 ppm, are rather similar to those, viz. 3.89 and 4.63 ppm, observed for tetramethylethylene-bridged ferrocene in  $\text{CDCl}_3$  [6]; the  $^1\text{H}$  NMR spectrum confirms the diamagnetism of the *ansa*-chromocene carbonyl adduct.

A crystal of approximate dimensions  $0.2 \times 0.2 \times 0.9\text{ mm}$ , sealed in a glass capillary under argon, was used for X-ray measurements. Space group, cell parameters and X-ray diffraction intensities were determined on a Syntex-P3 four-circle diffractometer at 233 K ( $\text{Mo-K}\alpha$ ,  $\lambda\ 71.069\text{ pm}$ ), graphite monochromator,  $\omega$ -scan with  $1.4 \leq \dot{\omega} \leq 29.3^\circ\text{ min}^{-1}$  and  $1 < 2\theta < 44^\circ$ . The crystals are monoclinic, space group  $P2_1/c$ ;  $a\ 1036(1)$ ,  $b\ 725.2(7)$ ,  $c\ 1859(4)\text{ pm}$ ;  $\beta\ 97.05(0.15)^\circ$ ; 4 crystallographically equivalent molecules per unit cell;  $V\ 1387 \times 10^6\text{ pm}^3$ ;  $d_{\text{calc}}\ 1.25\text{ g/m}^3$ ; absorption coefficient  $\mu\ 8.2\text{ cm}^{-1}$ .

Of 1658 reflexes measured, 1320 with  $I \geq 2\sigma$  were used for solving and refining the structure. The structure was solved by direct methods (SHELXTL program [7]) and refined with a partially anisotropic model (H atom positions were taken from the Fourier difference map, but not refined). The refinement converged at  $R_1 = (\sum |F_o| - |F_c|) / \sum |F_o| = 0.076$  and  $R_2 = [\sum w(|F_o| - |F_c|)^2]^{1/2} / [\sum w|F_o|^2]^{1/2} = 0.087$ . Structural data are listed in Tab. 1; bond lengths and bond angles are listed in Table 2; the molecular structure obtained is represented in Fig. 1.

The Cr–C(ring) distances are identical within standard deviations; their mean value of  $216 \pm 1\text{ pm}$  is in close agreement with a value of  $216.9(0.4)\text{ pm}$  observed for  $(\text{C}_5\text{H}_5)_2\text{Cr}$  by gas phase electron diffraction [8]. The centroid–metal–centroid angle is  $143.3^\circ$ , substantially larger than the angles of  $128$  and  $130^\circ$  observed in ethylene-

TABLE 1  
STRUCTURAL PARAMETERS FOR  $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4)_2\text{Cr}(\text{CO})$

Fractional coordinates (with e.s.d.'s) and isotropic temperature factors <sup>a</sup>

Atom	<i>x/a</i>	<i>y/b</i>	<i>z/c</i>	<i>U</i>
Cr	0.7610(1)	0.3858(2)	0.5108(1)	
C(1)	0.8123(9)	0.135(1)	0.4579(5)	0.029(2)
C(2)	0.8580(8)	0.126(1)	0.5316(5)	0.029(2)
C(3)	0.7461(8)	0.137(1)	0.5714(5)	0.025(2)
C(4)	0.6345(8)	0.155(1)	0.5188(5)	0.026(2)
C(5)	0.6738(9)	0.152(1)	0.4496(5)	0.031(2)
C(6)	0.7366(8)	0.475(1)	0.6178(5)	0.025(2)
C(7)	0.8652(9)	0.524(1)	0.6032(5)	0.032(2)
C(8)	0.8557(9)	0.644(1)	0.5454(5)	0.034(2)
C(9)	0.7208(9)	0.672(1)	0.5228(5)	0.033(2)
C(10)	0.6467(9)	0.570(1)	0.5665(5)	0.032(5)
C(11)	0.7005(8)	0.339(1)	0.6751(5)	0.025(2)
C(12)	0.7496(8)	0.141(1)	0.6529(5)	0.029(2)
C(13)	0.554(1)	0.345(2)	0.6768(6)	0.045(3)
C(14)	0.763(1)	0.402(2)	0.7490(6)	0.047(3)
C(15)	0.891(1)	0.100(2)	0.6890(6)	0.045(3)
C(16)	0.663(1)	-0.017(2)	0.6766(6)	0.043(3)
C(17)	0.7821(9)	0.479(1)	0.4207(5)	0.032(2)
O	0.7982(8)	0.540(1)	0.3648(4)	
H(1)	0.852	0.146	0.418	
H(2)	0.947	0.103	0.560	
H(4)	0.548	0.165	0.524	
H(5)	0.611	0.149	0.403	
H(7)	0.928	0.487	0.633	
H(8)	0.930	0.705	0.519	
H(9)	0.676	0.715	0.484	
H(10)	0.558	0.558	0.561	
H(13A)	0.512	0.310	0.636	
H(13B)	0.511	0.258	0.709	
H(13C)	0.523	0.456	0.691	
H(14A)	0.842	0.396	0.753	
H(14B)	0.710	0.495	0.768	
H(14C)	0.755	0.336	0.797	
H(15A)	0.803	0.116	0.699	
H(15B)	0.908	0.012	0.666	
H(15C)	0.927	0.156	0.679	
H(16A)	0.586	-0.006	0.658	
H(16B)	0.724	-0.077	0.674	
H(16C)	0.673	-0.005	0.730	

Anisotropic temperature factors <sup>b</sup>

Atom	<i>U</i> <sub>11</sub>	<i>U</i> <sub>22</sub>	<i>U</i> <sub>33</sub>	<i>U</i> <sub>23</sub>	<i>U</i> <sub>13</sub>	<i>U</i> <sub>12</sub>
Cr	0.0256(8)	0.0156(8)	0.0256(8)	-0.0004(7)	0.0047(6)	-0.0006(7)
O(1)	0.076(6)	0.050(5)	0.032(4)	0.010(4)	0.014(4)	0.000(4)

<sup>a</sup> The designations HxA, HxB, HxC refer to the methyl hydrogen atoms bonded to Cx. Other hydrogens: Hx bonded to Cx. <sup>b</sup> The anisotropic factors *U*<sub>*ij*</sub> refer to:  $T = \exp(-2\pi^2[U_{11}h^2a^* + \dots + 2hka^*b^*U_{12} + \dots]); U_{ij}$  in  $10^4 \text{ pm}^2$ .

TABLE 2

INTERATOMIC DISTANCES (pm) AND BOND ANGLES ( $^{\circ}$ ) (Cp(1) = centroid of ring 1, plane 1 = mean plane of ring 1)

Cr-C(1)	217(1)	Cr-C(6)	214(1)
Cr-C(2)	216(1)	Cr-C(7)	217(1)
Cr-C(3)	214(1)	Cr-C(8)	218(1)
Cr-C(4)	216(1)	Cr-C(9)	213(1)
Cr-C(5)	218(1)	Cr-C(10)	215(1)
Cr-Cp(1)	178(1)	Cr-Cp(2)	178(1)
Cr-C(17)	185(1)	C(17)-O	116(1)
C-C(ring 1)	$144 \pm 3$	C-C(ring 2)	$143 \pm 3$
C(6)-C(11)	153(1)	C(3)-C(12)	151(1)
	C(11)-C(12)	160(1)	
Cr-C(17)-O	178.4(9)	Cp(1)-Cr-Cp(2)	143.3(5)
C(6)-C(11)-C(12)	106.1(7)	C(3)-C(12)-C(11)	107.9(7)
<i>Angles between ring planes (<math>^{\circ}</math>)</i>			
Plane 1-Plane 2	38.5(5)		




Fig. 1. Molecular structure of  $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4)_2\text{Cr}(\text{CO})$ . Mutually perpendicular projections on to plane containing the two ring centroids and the  $\text{Cr}-\text{C}\equiv\text{O}$  axis (left) and on to the bisector plane between the two ring mean planes (right). Atomic positions designated by numbers only refer to carbon atoms of the ligand framework.

bridged [9] and in unbridged titanocene dichloride [10], but quite close to a centroid-metal-centroid angle of  $145.8^{\circ}$  observed for  $(\text{C}_5\text{H}_5)_2\text{MoH}_2$  [11]. Although not fixed by crystallographic site symmetry, the structure depicted in Fig. 1 maintains, without major deviations, the  $\text{C}_2$  symmetry along the  $\text{Cr}-\text{C}\equiv\text{O}$  axis expected for a monoligated metallocene derivative with low-spin  $d^4$  electron configuration [12].

## Experimental

All operations were carried out with exclusion of air and moisture, using Schlenk and vacuum line techniques. Anhydrous  $\text{CrCl}_2 \cdot \text{THF}$  was obtained from Cr metal and  $\text{HCl}/\text{THF}$  [13]; the Grignard compound  $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4\text{MgCl})_2 \cdot 4\text{THF}$  was prepared as described in ref. 1.



In a 250 ml side-arm flask, 80 ml THF are condensed at  $-78^\circ\text{C}$  on to 0.72 g (3.7 mmol)  $\text{CrCl}_2 \cdot \text{THF}$  and 2.34 g (3.7 mmol)  $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4\text{MgCl})_2 \cdot 4\text{THF}$ . After introduction of an atmosphere of CO, the mixture is stirred and allowed to warm during 12 h to room temperature, at which it is stirred for another day. The solvent is then removed in vacuo. The residue is dried without further warming and taken up in ca. 100 ml of pentane, filtered off and washed twice with pentane. The combined pentane extracts are concentrated to 10 ml and kept at  $-40^\circ\text{C}$  to give 570 mg (1.94 mmol, 52% theoretical yield) of  $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4)_2\text{Cr}(\text{CO})$  as red-brown crystals. The material can be further purified by sublimation at  $40-45^\circ\text{C}$  on to a cold finger or at  $60-70^\circ\text{C}$  in a sublimation tube; it is very air-sensitive.

$^1\text{H}$  NMR in benzene- $d_6$  at  $25^\circ\text{C}$ : 0.91 ppm (s, 12H), 3.60 ppm (t, 4H,  $J$  2.14 Hz), 4.38 ppm (t, 4H,  $J$  2.14 Hz).

### Acknowledgements

This work was supported by Deutsche Forschungsgemeinschaft (grant Bri 542/82), by funds of the University of Konstanz and by Fonds der Chemischen Industrie.

### References

- 1 H. Schwemlein and H.H. Brintzinger, *J. Organomet. Chem.*, 254 (1983) 69.
- 2 G. Wilkinson, F.A. Cotton and J.M. Birmingham, *J. Inorg. Nucl. Chem.*, 2 (1956) 95.
- 3 G.W. Watt and L.J. Baye, *J. Inorg. Nucl. Chem.*, 26 (1964) 2099.
- 4 K.L. Tang Wong and H.H. Brintzinger, *J. Amer. Chem. Soc.*, 97 (1975) 5143.
- 5 M.H. Chisholm, T.D. Gallagher and W.W. Reichert, *Synth. React. Inorg. Metalorgan. Chem.*, 7 (1977) 279.
- 6 K.L. Rinehart, Jr., A.K. Frerichs, P.A. Kittle, L.F. Westman, D.H. Gustafson, R.L. Pruett and J.E. McMahon, *J. Amer. Chem. Soc.*, 82 (1960) 4111.
- 7 G.M. Sheldrick, Universität Göttingen 1981.
- 8 E. Gard, A. Haaland, D.P. Novak and R. Seip, *J. Organomet. Chem.*, 88 (1975) 181.
- 9 J.A. Smith, J. v.Seyerl, G. Huttner and H.H. Brintzinger, *J. Organomet. Chem.*, 173 (1979) 175.
- 10 A. Clearfield, D.K. Warner, C.H. Saldarriaga-Molina, R. Ropal and I. Bernal, *Can. J. Chem.*, 53 (1975) 1622.
- 11 A.J. Schultz, K.L. Stearley, J.M. Williams, R. Mink and G.D. Stucky, *Inorg. Chem.*, 16 (1977) 3303.
- 12 J.W. Lauher and R. Hoffmann, *J. Amer. Chem. Soc.*, 98 (1976) 1729.
- 13 L.F. Larkworthy and M.H.O. Nelson-Richardson, *Chem. Ind.* (1974) 164.  
F.H. Köhler and W. Prössdorf, *Z. Naturforsch. B*, 32 (1977) 1026.