

Research article

Open Access

## A link of $\text{Ca}^{2+}$ to cAMP oscillations in *Dictyostelium*: the calmodulin antagonist W-7 potentiates cAMP relay and transiently inhibits the acidic $\text{Ca}^{2+}$ -store

Dieter Malchow\*, Daniel F Lusche and Christina Schlatterer

Address: Department of Biology, University of Konstanz, D-78457 Konstanz, Germany

Email: Dieter Malchow\* - [Renate.Gimmi@uni-konstanz.de](mailto:Renate.Gimmi@uni-konstanz.de); Daniel F Lusche - [daniel.lusche@uni-konstanz.de](mailto:daniel.lusche@uni-konstanz.de); Christina Schlatterer - [Christina.Schlatterer@uni-konstanz.de](mailto:Christina.Schlatterer@uni-konstanz.de)

\* Corresponding author

Published: 17 May 2004

Received: 31 October 2003

*BMC Developmental Biology* 2004, 4:7

Accepted: 17 May 2004

This article is available from: <http://www.biomedcentral.com/1471-213X/4/7>

© 2004 Malchow et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

### Abstract

**Background:** During early differentiation of *Dictyostelium* the attractant cAMP is released periodically to induce aggregation of the cells. Here we pursue the question whether pulsatile cAMP signaling is coupled to a basic  $\text{Ca}^{2+}$ -oscillation.

**Results:** We found that the calmodulin antagonist W-7 transiently enhanced cAMP spikes. We show that W-7 acts on an acidic  $\text{Ca}^{2+}$ -store: it abolished ATP-dependent vesicular acidification, inhibited V-type  $\text{H}^+$ ATPase activity more potently than the weaker antagonist W-5 and caused vesicular  $\text{Ca}^{2+}$ -leakage. Concanamycin A, an inhibitor of the V-type  $\text{H}^+$ -pump, blocked the  $\text{Ca}^{2+}$ -leakage elicited by W-7 as well as cAMP-oscillations in the presence of W-7. Concanamycin A caused an increase of the cytosolic  $\text{Ca}^{2+}$ -concentration whereas W-7 did not. In case of the latter,  $\text{Ca}^{2+}$  was secreted by the cells. In accord with our hypothesis that the link from  $\text{Ca}^{2+}$  to cAMP synthesis is mediated by a  $\text{Ca}^{2+}$ -dependent phospholipase C we found that W-7 was not active in the phospholipase C knockout mutant.

**Conclusion:** We conclude that the potentiation of cAMP relay by W-7 is due to a transient inhibition of the acidic  $\text{Ca}^{2+}$ -store. The inhibition of the proton pump by W-7 causes a leakage of  $\text{Ca}^{2+}$  that indirectly stimulates adenylyl cyclase activity via phospholipase C.

### Background

The development of *Dictyostelium discoideum* to the multicellular stage and formation of fruiting bodies is largely determined by cAMP. During aggregation cAMP is required extracellularly to attract amoebae to the aggregation centre and intracellularly to activate PKA. PKA in turn is involved in an oscillatory loop of adenylyl cyclase activation as well as in gene regulation. Later in development, after tip formation, cAMP waves emitted from the tip reg-

ulate cell movement, whereas intracellular cAMP is necessary for differentiation of prespore and prestalk cells [for review see [1,2]].

There are three distinct genes that encode adenylyl cyclases [3-6]. The main enzyme for aggregation (ACA) displays little activity in growing cells and accumulates during differentiation with a maximum at aggregation. If ACA is genetically inactivated cells do not aggregate. The

second enzyme (ACRA) is required at culmination for sporulation and construction of the stalk but its mRNA accumulates to high levels after 4 h of starvation. Transcripts of the third class (ACG) are found during spore germination. Its activity is upregulated by high osmolarity and is essential for the maintenance of dormancy [6].

The aggregative enzyme (ACA) is activated following cAMP-binding to the G-protein coupled cell surface cAMP receptor (cAR1). The signal is transmitted via  $G_{\beta\gamma}$  and a cytosolic activator protein named CRAC that is thought to mediate the function of  $G_{\beta\gamma}$  in stimulating ACA [for review see [7]]. Among further components of this regulatory loop are RasC [8], a MAP-Kinase ERK2 [9] and RegA, a two-component signaling cAMP-phosphodiesterase [10,11].


cAMP is secreted periodically after about 4 h of differentiation with a period length in cell suspension of about 7 min. cAMP oscillations are accompanied by oscillations of cGMP,  $Ca^{2+}$ ,  $K^+$  and  $H^+$ . A basal extracellular  $Ca^{2+}$ -oscillation is transformed to spikes during cAMP relay and continues after cessation of the latter [12,13].

cAMP induces an influx of  $Ca^{2+}$  and a cytosolic  $Ca^{2+}$  transient [14-21]. Cytosolic  $Ca^{2+}$  is taken up into an  $IP_3$ -sensitive  $Ca^{2+}$ -store as well as into acidic  $Ca^{2+}$ -stores [22-26]. Activation of plasma membrane  $Ca^{2+}$ -ATPase results in  $Ca^{2+}$ -efflux which decreases during the course of differentiation [14,27,28] with a concomitant increase of sequestered  $Ca^{2+}$  [29].  $Ca^{2+}$ -influx is strongly reduced in the IplA minus strain that should lack an  $IP_3$  receptor-like  $Ca^{2+}$ -channel [30]. One type of calmodulin antagonists, calmidazolium, was found to stimulate  $Ca^{2+}$ -entry, to cause a cytosolic  $Ca^{2+}$ -increase and to delay light scattering oscillation and cAMP relay [31,32]. Calmidazolium, but not the other calmodulin antagonist W-7, blocked *Dictyostelium* calcineurin activity [[33], unpublished results].

In contrast to calmidazolium, W-7 inhibited  $Ca^{2+}$ -influx, enhanced light scattering as well as cAMP oscillation. Since W-7 was the first known agent stimulating oscillations we investigated its mode of action. The results show that W-7 transiently inhibited the acidic  $Ca^{2+}$ -store and that phospholipase C is required as a link of  $Ca^{2+}$  to cAMP oscillations.

## Results

*Dictyostelium* cell suspensions start to oscillate at about 4 h after induction of differentiation. This can be measured by the light scattering technique or with a  $Ca^{2+}$ -sensitive electrode. The light scattering oscillations coincide with cAMP relay and extracellular  $Ca^{2+}$ -oscillations [34-36]. Oscillations start out with low amplitude and gain maximal height after 5 to 8 cycles due to feedback stimulation



**Figure 1**


Early light scattering oscillations are enhanced by W-7. Free running oscillations of a cell suspension at  $2 \times 10^7$  cells per ml were recorded using the light scattering technique as described in Material and Methods 4 h after induction of differentiation. 30  $\mu$ M W-7 was added where indicated. One out of ten independent experiments is shown.

by extracellular cAMP. We found that the calmodulin (CaM) antagonist W-7 transiently enhanced light scattering oscillation amplitude when added early during oscillation (Fig. 1). W-7 addition also caused a phase advance indicating that W-7 interferes with the principal oscillator. Fig. 2 demonstrates that W-7 augmented cAMP relay and caused a phase advance in cAMP synthesis. The effect of W-7 was most pronounced at the onset of oscillations with a  $6.3 \pm 2.7$  fold ( $n = 3$ ) increase in cAMP relay. Later on, smaller increases were measured ( $1.3 \pm 0.1$  fold,  $n = 3$ ) because oscillation amplitude had reached their maximal values. In order to find the target of W-7, we first tested whether W-7 directly activates adenylcyclase in cell lysates. In four separate experiments no significant alteration of the catalytic enzyme activity by W-7 could be detected ( $0.96 \pm 0.25$ ,  $n = 10$ ). We then investigated the following processes that involve  $Ca^{2+}$ .

### cAMP-induced $Ca^{2+}$ -influx

Addition of cAMP to a cell suspension induces a rapid transient  $Ca^{2+}$ -influx [13-17]. In the presence of W-7 cAMP-induced  $Ca^{2+}$ -influx was reduced by 60–70% (Fig. 3). Maximal inhibition occurred at about 30  $\mu$ M W-7.

Inhibition of  $Ca^{2+}$ -influx is not sufficient as an explanation for the enhancement of oscillation amplitude by W-7. The IplA minus strain oscillated and responded to W-7




**Figure 2**  
Just starting cAMP oscillations are upregulated by W-7. cAMP concentrations were determined as described in Materials and Methods. 80  $\mu$ l samples were withdrawn from the cell suspension before and after addition of 30  $\mu$ M W-7. The peak cAMP value corresponds to 89 pmol/ $10^7$  cells. One out of three independent experiments is shown.


although receptor-mediated  $Ca^{2+}$ -influx is strongly reduced (30, Fig. 4). The amplitude increased  $1.8 \pm 0.3$  fold ( $n = 5$ ) and cAMP  $2.7 \pm 0.5$  fold ( $n = 3$ ) following addition of 30  $\mu$ M W-7.

**Vesicular  $Ca^{2+}$ -release**

Previously, we have shown that vesicular  $^{45}Ca^{2+}$ -uptake was inhibited by a CaM antibody and CaM antagonists including W-7 [25]. We now show that W-7 also induced substantial  $Ca^{2+}$ -release (Fig. 5). A dose response curve revealed that 30  $\mu$ M W-7 nearly saturated the response (Fig. 6). To examine whether CaM is involved we asked whether a peptide containing the CaM-binding domain of CaMKII (CaMBP) competes for W-7-induced  $Ca^{2+}$ -release. We found that CaMBP dose dependently reduced the response to W-7. Half maximal inhibition occurred at



**Figure 3**  
Dose response curve for inhibition of cAMP-induced  $Ca^{2+}$ -influx in the presence of W-7. Each data point represents a separate experiment.  $Ca^{2+}$ -influx was determined as described in Materials and Methods. The mean control influx was  $165 \pm 68$  pmol per  $10^7$  cells. The extracellular  $Ca^{2+}$ -concentration was about 4  $\mu$ M.




**Figure 4**  
Light scattering oscillation of the IplA minus strain and its response to W-7. The experiment was performed as described in the legend of figure 1.

about 20  $\mu$ M of the peptide (Fig. 7). CaMBP also induced  $Ca^{2+}$ -release albeit at a slower rate than W-7 (data not shown).

**The acidic  $Ca^{2+}$ -store**

*Dictyostelium* cells contain at least two  $Ca^{2+}$ -stores besides mitochondria and mass dense granules, the endoplasmic




**Figure 5**

Induction of vesicular  $\text{Ca}^{2+}$ -release by W-7. Vesicular  $\text{Ca}^{2+}$ -uptake was measured as described in Materials and Methods. Addition of 400  $\mu\text{M}$  ATP initiated  $\text{Ca}^{2+}$ -uptake. Following a  $\text{Ca}^{2+}$  calibration pulse W-7 was added. One out of ten independent experiments is shown.


reticulum (ER) and an acidic  $\text{Ca}^{2+}$ -store [22-26]. The acidic  $\text{Ca}^{2+}$ -store contains a V-type  $\text{H}^+$ ATPase that acidifies the vesicles. The proton gradient is used to transport  $\text{Ca}^{2+}$  into the store. In the presence of W-7 ATP-dependent acidification was abolished (Fig. 8). The half maximal concentration for W-7 was  $16 \pm 4 \mu\text{M}$  ( $n = 3$ ). The less potent CaM antagonist W-5 was effective at a correspondingly higher concentration. Half maximal inactivation occurred at  $50 \pm 14 \mu\text{M}$  ( $n = 3$ ). The graded potency of W-7 and W-5 to block ATP-dependent acidification supports the notion that the CaM antagonists act via CaM. To find out whether CaM regulates the  $\text{H}^+$ -pump we measured V-type  $\text{H}^+$ ATPase activity in the presence of W-7 or W-5. 90  $\mu\text{M}$  W-7 or W-5 reduced enzyme activity ( $3.7 \text{ nmol P}_i/\text{min} \cdot \text{mg}$ ) by  $34 \pm 3\%$  and by  $9.7 \pm 5\%$  ( $n = 3$ ), respectively. Measurement of  $\text{H}^+$ ATPase activity requires 30 min

of incubation. The incomplete inhibition of the ATPase is probably the reason for the transient effect of W-7 on light scattering oscillation.


If W-7 acts on the acidic store inhibition of the  $\text{H}^+$ ATPase should eliminate or potentially reduce the response to W-7. A potent inhibitor of the V-type  $\text{H}^+$ ATPase is concanamycin A (CMA). We found that preincubation with CMA reduced both, vesicular  $\text{Ca}^{2+}$ -uptake and W-7-induced  $\text{Ca}^{2+}$ -release (Fig. 9).  $\text{Ca}^{2+}$ -uptake was inhibited by  $64 \pm 9\%$  ( $n = 3$ ),  $\text{Ca}^{2+}$ -release by  $86 \pm 2\%$  ( $n = 3$ ). These results show that the acidic  $\text{Ca}^{2+}$ -store accounts for 65% of vesicular transport. Blockade of the proton pump abolished  $\text{Ca}^{2+}$ -release induced by W-7. The occurrence of  $\text{Ca}^{2+}$ -release can thus be explained by leakage of  $\text{Ca}^{2+}$  from the store due to transient inhibition of the proton pump.



**Figure 6**  
Dose response curve for W-7-induced vesicular Ca<sup>2+</sup>-release. The experiment was performed as described in the legend of figure 5. One out of two independent experiments is shown.



**Figure 7**  
Dose response curve for inhibition of W-7-induced vesicular Ca<sup>2+</sup>-release by CaMBP. The experiment was performed as described in the legend of figure 4. 100 μM W-7 was added after preincubation of CaMBP with the vesicles for 10 to 15 min to allow for Ca<sup>2+</sup>-release elicited by CaMBP. 100% Ca<sup>2+</sup>-release amounted to 6 nmol Ca<sup>2+</sup> per mg protein. One out of three independent experiments is shown.




**Figure 8**  
Inhibition of ATP-dependent vesicular acidification by CaM antagonists. Determination of ATP-dependent acidification was performed as described in Material and Methods. Control acidification amounted to 20 relative fluorescence units. One out of three independent experiments is shown.

**Effect of CMA on oscillation**

Since W-7 and CMA act on the H<sup>+</sup>-pump and W-7 enhances cAMP synthesis we wondered whether CMA influences cAMP relay. We measured light scattering and cAMP concentrations before and after application of CMA and W-7. We found that after CMA application, oscillation continued for one cycle and then ceased. Addition of W-7 caused a small increase in light scattering but cAMP relay was blocked (Fig. 10). In 3 independent experiments inhibition of cAMP relay amounted to 92 ± 5%.

**W-7 induces Ca<sup>2+</sup>-efflux in intact cells**

We have shown that W-7 caused vesicular Ca<sup>2+</sup>-leakage. Does this result in an increase of [Ca<sup>2+</sup>]<sub>i</sub>? In 210 cells analyzed 30 μM W-7 did not cause a cytosolic Ca<sup>2+</sup>-concentration change. By contrast, CMA raised [Ca<sup>2+</sup>]<sub>i</sub>, indicating that CMA acts more potently on the H<sup>+</sup>-pump than W-7 (Fig. 11). However, W-7 elicited Ca<sup>2+</sup>-efflux from cells. The efflux amounted to 1.6 ± 0.5 nmol/10<sup>7</sup> cells (n = 8). If we compare efflux with leakage from the acidic Ca<sup>2+</sup>-store which amounted to 13.0 ± 4, 8 nmol/mg protein (n = 8) and account for a six-fold enrichment of Ca<sup>2+</sup>-stores in the vesicle preparation with respect to cells (1 × 10<sup>7</sup> cells contain 1 mg protein) the 13 nmol reduce to 2.2 nmol/mg protein. Thus, the most part (1.6/2.2 = 72%) of W-7-induced Ca<sup>2+</sup>-leakage is secreted by the cells.

**Figure 9**

Vesicular  $\text{Ca}^{2+}$ -uptake and W-7-induced  $\text{Ca}^{2+}$ -release in the absence or presence of CMA. The experiment was performed as described in the legend of figure 5.  $30 \mu\text{M}$  CMA was added 2 min before ATP addition (right panel) the control received the solvent DMSO (left panel). One out of three independent experiments is shown.


#### Augmentation of cAMP relay by W-7 requires phospholipase C

In order to find out whether the  $\text{Ca}^{2+}$ -activated phospholipase C (PLC) of *Dictyostelium* represents a link to cAMP relay we analyzed the PLC null strain HD10, for light scattering and cAMP oscillations. In HD10, light scattering amplitudes contain a negative component as compared to Ax-2 (Figs. 1,2,9). W-7 neither elicited an increase in light scattering amplitude nor in cAMP relay (Fig. 12). In four experiments only  $76 \pm 22\%$  ( $n = 4$ ) of the previous control amplitude of cAMP synthesis was measured. The parental strain of HD10, Ax3, responded to W-7 with a  $2.8 \pm 0.6$  fold ( $n = 6$ ) increase in light scattering amplitude and did not show a negative component in light scattering oscillations (data not shown).

The occurrence of oscillations in the PLC null strain can be explained by an  $\text{IP}_5$ -phosphatase that hydrolyses  $\text{IP}_5$  to yield  $\text{IP}_3$ . The phosphatase is also activated by  $\text{Ca}^{2+}$  but requires higher concentrations [37-39].

#### Discussion

The  $\text{Ca}^{2+}$ -binding protein CaM regulates many proteins and enzymes in mammalian cells but in *Dictyostelium* only few targets in signal transduction are known: calcineurin, a CaM-dependent protein kinase, and the contractile vacuole [33,40,41]. Neither the cyclases, nor the phosphodiesterases, PLC or the plasma membrane  $\text{Ca}^{2+}$  ATPase were reported to be regulated by CaM [1,27]. Therefore, it came as a surprise that the CaM antagonist W-7 enhanced light scattering spikes and cAMP relay. The spike-shaped light scattering response, as an indicator for




**Figure 10**  
Light scattering and cAMP oscillation in the presence of CMA and W-7. The experiment was performed as described in the legend of figure 2. cAMP concentration were measured before the addition of CMA and after the addition of W-7. The results are expressed as xfold increase in cAMP concentration. One out of three independent experiments is shown.


cAMP relay, saturated at 30 μM W-7. The reported IC<sub>50</sub> value for W-7 as an antagonist for CaM is 25 μM [42,43] indicating that the effective concentration of W-7 matches with known CaM-targets.

In mammalian cells a variety of ion channels are regulated by CaM. Indeed, W-7 inhibited cAMP-induced Ca<sup>2+</sup>-influx (Fig. 3). However, in the IplA minus mutant, where receptor-mediated Ca<sup>2+</sup>-influx is almost absent, W-7 still augmented light scattering spikes and cAMP relay. This result excludes inhibition of Ca<sup>2+</sup>-influx as the main cause for the effect of W-7.

Another potent response to W-7 was vesicular Ca<sup>2+</sup>-leakage (Fig. 5). Leakage may be brought about because W-7



**Figure 11**  
Cytosolic Ca<sup>2+</sup>-concentration in the presence of CMA or W-7. Single cells were loaded with Fura-2-Dextran and the fluorescence ratio was measured as described in Materials and Methods. One out of three independent experiments is shown.



**Figure 12**  
Light scattering and cAMP-oscillation of the phospholipase C-null-strain, HD10. W-7 was added were indicated. Note that the light scattering change underlying cAMP relay is negative. The experiment was performed as described in the legend of Fig. 2.

inhibited the vesicular proton pump, thus abolishing the proton gradient (Fig. 8) and  $\text{Ca}^{2+}$ -sequestration. Moreover, blocking the proton pump by CMA eliminated not only the effect of W-7 on the acidic  $\text{Ca}^{2+}$ -store but also on cAMP relay (Fig. 10)

W-7 seems to antagonise CaM since W-5 was less potent than W-7 and a CaM-binding peptide dose dependently inhibited the  $\text{Ca}^{2+}$ -response of W-7 (Fig. 7). CaM was shown to be located to the contractile vacuole, an organelle thought to be identical with or at least part of the acidic  $\text{Ca}^{2+}$ -store [41,44].

Although both, CMA and W-7, inhibit the acidic  $\text{Ca}^{2+}$ -store and CMA blocks the effects of W-7, the compounds differ:

1. CMA is lipid soluble and targets the  $V_0$ -portion of the  $\text{H}^+$ ATPase. By contrast, W-7 is fairly water soluble and seems to inhibit the  $\text{H}^+$ ATPase indirectly via CaM.
2. The effect of CMA is long lasting (Figs. 9,10) whereas the effect of W-7 is transient (Fig. 1,5).
3. CMA elicits an increase of the cytosolic  $\text{Ca}^{2+}$ -concentration whereas W-7 does not (Fig. 11). A likely explanation for the different outcomes could be that W-7 acts on the acidic stores located at or near the plasma membrane. This would explain why  $\text{Ca}^{2+}$ -release can be measured extracellularly and not as a global, cytosolic  $\text{Ca}^{2+}$ -increase. The  $\text{Ca}^{2+}$ -released, close to the plasma membrane, however, could still activate PLC [37,38].

How could the  $\text{Ca}^{2+}$ -oscillation be regulated? Previously, we have shown that signal transduction elicited by extracellular cAMP is mediated by phospholipase $_2$  activity: the fatty acids liberated induce  $\text{Ca}^{2+}$ -release from the acidic  $\text{Ca}^{2+}$ -store [26] as shown in Fig. 13.  $\text{Ca}^{2+}$  may stimulate a positive feedback loop by activation of PLC. One of the products,  $\text{IP}_3$ , elicits  $\text{Ca}^{2+}$ -release from the ER that may accumulate in the acidic store thereby completing the oscillatory cycle. Alternatively, PLC activity is reinforced. DAG, the second product of PLC, represents a link to adenylyl cyclase activity. In principle, cAMP levels could be raised likewise by inhibition of phosphodiesterase activity, for instance via phosphorylation by a DAG-dependent PKC. However, the peak of cAMP relay occurs earlier after W-7 application (Fig. 2). This shift cannot be due simply to a reduction in degradation of cAMP. In mammalian cells CalDAG-GEFs are known to activate Ras [45]. In *Dictyostelium*, RasC activity is required for adenylyl cyclase activity (ACA) [8]. Direct regulation of ACA by  $\text{Ca}^{2+}$  which occurs in two types of mammalian adenylyl cyclases (I-AC and VIII-AC) seems unlikely in *Dictyostelium* since no EF hand nor CaM binding site has been reported. The same holds true for other regulators of ACA,

namely Pianissimo [46], RasGEF (Aimless) [7], CRAC and RipA [47].


We have shown that disturbing the function of the acidic  $\text{Ca}^{2+}$ -store leads to inhibition of light scattering oscillation and cAMP relay. We have obtained the same result by inhibition of the other  $\text{Ca}^{2+}$ -store with Xestospongine C (XeC), a drug which, in *Dictyostelium*, inhibits the  $\text{Ca}^{2+}$ -pump of the ER [48]. Both drugs led to an increase of the cytosolic  $\text{Ca}^{2+}$ -concentration.

A basic conclusion is that the oscillatory behaviour of *Dictyostelium* depends on both  $\text{Ca}^{2+}$ -stores. Extracellular  $\text{Ca}^{2+}$ -oscillations occur simultaneously with light scattering and cAMP oscillations [36]. The phase advance induced by the calmodulin antagonist W-7 suggests that W-7 interferes with an internal  $\text{Ca}^{2+}$ -oscillation to which light scattering changes and cAMP production are coupled.

When the cytosolic  $\text{Ca}^{2+}$ -concentration remains elevated above the basal level as found in case of inhibition of  $\text{Ca}^{2+}$ -uptake by XeC or CMA, a hitherto unknown target seems to mediate inhibition of ACA. A possible candidate could be CaM Kinase II which is known to inhibit mammalian adenylyl cyclase III *in vivo* [49].

We postulate that ACA can be regulated twice by  $\text{Ca}^{2+}$  during the oscillatory cycle, at least at its onset. A localised  $\text{Ca}^{2+}$ -increase stimulates ACA whereas an elevated cytosolic  $\text{Ca}^{2+}$ -concentration inhibits enzyme activity. A likely candidate mediating the activating  $\text{Ca}^{2+}$ -link to ACA activity is PLC as shown in Fig. 13. As soon as cAMP oscillations are fully developed, the extracellular loop, that is: cAMP released by the cells binds to the cAMP receptor and reinforces signal transduction, provides for stimulation of cAMP synthesis and enhances  $\text{Ca}^{2+}$ -oscillations. This is true for the wild type and for the PLC null strain. Under this condition, W-7 augmented spike-shaped light scattering oscillations and cAMP relay only to a small extent. However, the regulatory link persists to a later stage (sinusoidal light scattering oscillations) where cAMP relay is absent. W-7 even then restarted spike-shaped light scattering oscillations (unpublished).

As stated above, in the absence of PLC a  $\text{Ca}^{2+}$ -sensitive  $\text{IP}_5$ -phosphatase hydrolyses  $\text{IP}_5$  to yield  $\text{IP}_3$ . Thus  $\text{Ca}^{2+}$ -oscillations still occur (unpublished results) but is there a link to cAMP synthesis although DAG is not provided for by  $\text{IP}_5$ -phosphatase? In order to obtain cAMP oscillations the extracellular loop, that is restimulation of cAR1 by extracellular cAMP, might be sufficient. Alternatively, a hitherto unknown  $\text{Ca}^{2+}$ -sensitive mechanism exists that is independent of DAG. We know that PLC null cells display a reduced sensitivity to FA for  $\text{Ca}^{2+}$ -influx [26] indicating that the sensitivity to  $\text{Ca}^{2+}$  may have changed in the strain.



**Figure 13**

Hypothetical scheme for  $Ca^{2+}$ -oscillations in *Dictyostelium* and a link to adenylyl cyclase. Adenylyl cyclase was shown to be activated by  $G_{\beta\gamma}$ , rasC, and CRAC.  $PI_3K$  activity provides for  $PIP_3$  allowing cytosolic CRAC to bind to the plasma membrane. Aimless and RipA are thought to act in concert with rasC. The direct target of pianissimo is not yet known nor the detailed sequence of events leading to cAMP synthesis. ERK2 stimulation by cAR1 is independent of rasC [8]. ERK2 inhibits RegA, a cytosolic phosphodiesterase. cAR1 also activates phospholipase  $A_2$  that liberates fatty acids (FA) and elicits  $Ca^{2+}$ -influx [17,26].  $Ca^{2+}$  released by fatty acids from the acidic store activates a positive feedback loop via PLC activity.  $IP_3$  stimulates  $Ca^{2+}$ -release from the ER, thereby raising cytosolic  $Ca^{2+}$  and enhancing PLC activity.  $Ca^{2+}$  sinks are provided by  $Ca^{2+}$ -uptake into the ER and the acidic  $Ca^{2+}$ -store. Inhibition of either pump, the  $Ca^{2+}$ -pump by XeC or the  $H^+$ -pump by CMA leads to an increase of  $[Ca^{2+}]_i$  and to a blockade of  $Ca^{2+}$ -oscillation as well as cAMP relay [48]. ACA is suggested to be activated by a CalDAG stimulated exchange factor acting on RasC. cAMP released to the outside of the cell elicits feed back stimulation during oscillation in cell suspension. Extracellular phosphodiesterase (PdsA) provides for a low extracellular cAMP level allowing sensitive detection of the next wave.

This result is in accord with the lower  $Ca^{2+}$  affinity of the  $IP_5$ -phosphatase as compared to PLC. Therefore it is not unlikely that another pathway to cAMP production is open.

**Conclusions**

Unexpectedly, the CaM antagonist W-7 augmented pulsatile cAMP relay in *Dictyostelium*. We found that W-7 caused  $Ca^{2+}$ -leakage by transient inhibition of the V-type  $H^+$ -pump. A target of  $Ca^{2+}$  is phospholipase C. A PLC-null-

mutant failed to respond to W-7. Inhibition of the acidic  $\text{Ca}^{2+}$ -store by CMA blocked the effect of W-7 on  $\text{Ca}^{2+}$ -leakage and cAMP relay. We conclude that a basal  $\text{Ca}^{2+}$ -oscillation regulates cAMP synthesis (Fig. 13):  $\text{Ca}^{2+}$ -released by one of the stores activates PLC.  $\text{IP}_3$  reinforces  $\text{Ca}^{2+}$ -release and PLC-activity. The  $\text{Ca}^{2+}$ -signal is terminated by transport into the stores. DAG may stimulate adenylyl cyclase activity via RasC. cAMP in an extracellular loop triggers receptor-mediated fatty acid release that in turn causes  $\text{Ca}^{2+}$ -release from the acidic store thereby enhancing the cycle.

## Methods

### Chemicals

N-(6-aminohexyl)5-chloro-1-naphthalene sulfonamide (W-7), N-1-(6-aminohexyl)naphthalene sulfonamide (W-5) and concanamycin A were from Fluka, Buchs, Switzerland. Fluo-3, Fura-2 and Fura-2-Dextran were purchased by MobiTec, Göttingen, Germany, and calmodulin binding peptide of CaMKII by Calbiochem, Bad Soden, Germany.

### Cells and culture

The axenic strain Ax2 and the phospholipase C-null mutant, HD10, kindly provided by P.J. Van Haastert, University of Groningen, The Netherlands, were grown in shaking culture as described [50,51]. Differentiation was induced by washing the cells twice in ice-cold Sørensen phosphate buffer (17 mM  $(\text{KH}_2/\text{Na}_2\text{H})\text{PO}_4$ , pH 6.0). Cells were shaken on a rotary shaker at 23 °C, 150 rpm at  $2 \times 10^7$  cells/ml until use.

### Preparation of vesicles

20 ml  $2 \times 10^7$  cells/ml were shaken in Sørensen phosphate buffer containing 5 mM EGTA for 2–4 h, washed once in ice-cold 20 mM Hepes buffer, pH 7.2, resuspended at  $2 \times 10^8$  cells/ml, and lysed by passage through nuclepore filters. Immediately, 3% sucrose, 50 mM KCl, 1 mM  $\text{MgCl}_2$ , 10  $\mu\text{g}/\text{ml}$  leupeptin, 1  $\mu\text{g}/\mu\text{l}$  aprotinin and 1.25 mM dithiothreitol were added (final concentration). After centrifugation for 5 min at  $150 \times g$  in order to remove unbroken cells, the supernatant was further fractionated by centrifugation for 5 min at  $3800 \times g$ . The sediment ( $P_0$ ) was resuspended in 1 ml 10 mM Tris buffer, pH 7.8, 3% sucrose, 50 mM KCl, 1 mM  $\text{MgCl}_2$ , 20  $\mu\text{g}/\text{ml}$  leupeptin and 2  $\mu\text{g}/\text{ml}$  aprotinin yielding a protein concentration of  $2.6 \pm 0.5$  mg/ml ( $n = 19$ ).  $P_0$  contained part of the ER and the acidic vesicle fraction with about 70% of the  $\text{Ca}^{2+}$ -transport activity.

### $\text{Ca}^{2+}$ -transport

$\text{Ca}^{2+}$ -transport was measured essentially as described [26]. In brief, about 60  $\mu\text{l}$   $P_0$  was added to 10 mM Hepes, pH 7.2, 50 mM KCl, 3% sucrose, 6  $\mu\text{g}/\text{ml}$  antimycin A, 6  $\mu\text{g}/\text{ml}$  oligomycin A, 100  $\mu\text{M}$   $\text{NaN}_3$ , 2 mM  $\text{MgCl}_2$ , and about

0.5  $\mu\text{M}$  Fluo-3 in a total volume of 1 ml.  $\text{Ca}^{2+}$ -uptake was initiated by addition of 400  $\mu\text{M}$  ATP. Fluo-3 fluorescence was monitored at 505 nm excitation and 526 nm emission with a fluorimeter (Perkin Elmer 650–10S, Überlingen, Germany). Statistical data are expressed throughout as the mean  $\pm$  S.D.

### Determination of $[\text{Ca}^{2+}]_i$

Cells were loaded with the  $\text{Ca}^{2+}$ -indicator Fura-2-Dextran by electroporation [50]. After a wash in 5 mM Hepes, 5 mM KCl, pH 7.0 (H5 buffer), 10  $\mu\text{l}$  aliquots of cell suspension were placed on cover slips and incubated in a humid chamber. At 10 min before  $[\text{Ca}^{2+}]_i$  recordings buffer covering the cells was replaced by 90  $\mu\text{l}$  of fresh H5 buffer with or without  $\text{Ca}^{2+}$ , as indicated.  $[\text{Ca}^{2+}]_i$  imaging was performed on an Axiovert F100 TV (Zeiss, Jena, Germany). The cells were viewed with a  $100 \times$  Fluor objective (NA1.3). 334 and 380 nm excitation was performed with a mercury lamp and a rotating filter wheel. Images of the cells were recorded with an ICCD Camera (HL-A; Proxitronic, Bensheim, Germany). Image digitization and calibration were as described previously [34].

### Determination of $\text{Ca}^{2+}$ -efflux

70  $\mu\text{l}$  of 3 h starved cells ( $2 \times 10^7$  cells/ml) were added to 930  $\mu\text{l}$  Sørensen phosphate buffer, pH 6.0, containing 1.6  $\mu\text{M}$  of the fluorescent  $\text{Ca}^{2+}$ -indicator Fura-2. Cells were stirred at 23 °C and fluorescence emission was recorded at 510 nm and excitation wavelengths of 340 nm and 380 nm with a Perkin Elmer fluorimeter.  $\text{Ca}^{2+}$ -concentrations were calculated as described [31]. Following equilibration of the external  $\text{Ca}^{2+}$ -concentration to about 0.3  $\mu\text{M}$   $\text{Ca}^{2+}$  30  $\mu\text{M}$  W-7 was added.  $\text{Ca}^{2+}$ -efflux occurred within about 4 min and was calibrated by addition of known amounts of  $\text{Ca}^{2+}$ .

### ATP-dependent acidification

Acidification was measured by fluorescence quenching of the weak base acridine orange (480 nm excitation, 528 nm emission) according to Rooney and Gross [22]. 80 to 150  $\mu\text{g}$  protein were assayed in 1 ml buffer containing 10 mM Hepes buffer pH 7.2, 50 mM KCl, 3% sucrose, 2 mM  $\text{MgCl}_2$ , 2  $\mu\text{M}$  acridine orange, and 6  $\mu\text{g}/\text{ml}$  antimycin A, 6  $\mu\text{g}/\text{ml}$  oligomycin A, 100  $\mu\text{M}$   $\text{NaN}_3$  in order to inhibit mitochondrial activity. Since  $P_0$  contained vesicles with ATP-independent acidification the supernatant was used after centrifugation for 10 min at  $5800 \times g$ . Acidification was started with 1 mM ATP.

### Other measurements

$\text{Ca}^{2+}$ -influx was determined with a  $\text{Ca}^{2+}$ -sensitive electrode [17], and light scattering oscillations were recorded as described [32]. The amount of cAMP (intracellular and extracellular) present during oscillations in the presence or absence of drugs was measured using an enzyme

immunoassay (Biotrak, Amersham Pharmacia Biotech). Samples were prepared as described [32]. V-type H<sup>+</sup>ATPase activity was determined as described previously [26]. Adenylyl cyclase assays in cell lysates were performed essentially as described [52]. cAMP levels, determined by enzyme immunoassay, amounted to 11–85 pmol cAMP per min per mg protein. Protein concentrations were determined with the Coomassie protein assay reagent (Pierce) using bovine serum albumin as standard.

### Abbreviations

ACA, adenylyl cyclase A, CRAC, cytosolic activator of adenylyl cyclase; CaM, calmodulin; [Ca<sup>2+</sup>]<sub>i</sub>, intracellular Ca<sup>2+</sup>-concentration; CaMKII, calmodulin Kinase II, CMA, concanamycin A; IP<sub>3</sub>, inositol 1, 4, 5-trisphosphate; PLC, phospholipase C; RipA, Ras-interacting protein; XcC, Xestospongin C.

### Authors' contributions

DM conceived the study, participated in its design and coordination and performed Ca<sup>2+</sup>-transport experiments. DFL participated in the design of the study and carried out the cAMP relay studies. CS participated in the design of the study and performed cytosolic Ca<sup>2+</sup>-measurements. All authors read and approved the final manuscript.

### Acknowledgements

We thank E. Jaworski, U. Eichert and D. Horak for expert technical assistance, Dr. R. Schaloske for helpful discussions and the Deutsche Forschungsgemeinschaft for support.

### References

- Kessin RH: *Dictyostelium* Cambridge University Press; 2001.
- Saran S, Meima ME, Alvarrez-Curto E, Weening KE, Rozen DE, Schaap P: **cAMP signaling in Dictyostelium.** *J Muscle Res Cell Mot* 2002, **23**:793-802.
- Pitt GS, Milona N, Borleis J, Lin KC, Reed RR, Devreotes PN: **Structurally distinct and stage-specific adenylyl cyclase genes play different roles in Dictyostelium development.** *Cell* 1992, **69**:305-315.
- Kim H-J, Chang W-T, Meima ME, Gross JD, Schaap P: **A novel adenylyl cyclase detected in rapidly developing mutants of Dictyostelium.** *J Biol Chem* 1998, **273**:30859-30862.
- Söderbom F, Anjard C, Iranfar N, Fuller D, Loomis WF: **An adenylyl cyclase that functions during late development of Dictyostelium.** *Development* 1999, **126**:5463-5471.
- van Es S, Viridy KJ, Pitt GS, Meima ME, Sands TW, Devreotes PN, Cotter DA, Schaap P: **Adenylyl cyclase G, an osmosensor controlling germination of Dictyostelium spores.** *J Biol Chem* 1996, **271**:23623-23625.
- Parent CA, Devreotes PN: **Molecular genetics of signal transduction in Dictyostelium.** *Annu Rev Biochem* 1996, **65**:411-440.
- Lim CJ, Spiegelman G, Weeks G: **RasC is required for optimal activation of adenylyl cyclase and Akt/PKB during aggregation.** *EMBO J* 2001, **20**:4490-4499.
- Segall JE, Kuspa A, Shaulsky G, Ecke M, Maeda M, Gaskins C, Firtel RA, Loomis WF: **A MAP kinase necessary for receptor-mediated activation of adenylyl cyclase in Dictyostelium.** *J Cell Biol* 1995, **128**:405-413.
- Shaulsky G, Fuller D, Loomis WF: **A cAMP phosphodiesterase controls PKA-dependent differentiation.** *Development* 1998, **125**:691-699.
- Thomason PA, Traynor D, Cavet G, Chang WT, Harwood AJ, Kay RR: **An intersection of the cAMP/PKA and two-component signal transduction systems in Dictyostelium.** *EMBO J* 1998, **17**:2838-2845.
- Wurster B, Nanjundiah V, Malchow D: **Calcium oscillations in Dictyostelium discoideum.** In *Calcium as an intracellular Messenger in Eucaryotic Microbes* Edited by: O'Day DH. Washington: Am Soc Microbiol; 1990:228-242.
- Newell PC, Malchow D, Gross JD: **The role of calcium in aggregation and development of Dictyostelium.** *Experientia* 1995, **51**:1155-1165.
- Bumann J, Wurster B, Malchow D: **Attractant induced changes and oscillations of the extracellular Ca<sup>2+</sup> concentration in suspensions of differentiating Dictyostelium cells.** *J Cell Biol* 1984, **98**:173-178.
- Milne JL, Coukell MB: **A Ca<sup>2+</sup> transport system associated with the plasma membrane of Dictyostelium discoideum is activated by different chemoattractant receptors.** *J Cell Biol* 1991, **112**:103-110.
- Milne JLS, Wu LJ, Caterina MJ, Devreotes PN: **Seven helix cAMP receptors stimulate Ca<sup>2+</sup> entry in the absence of functional G proteins in Dictyostelium.** *J Biol Chem* 1995, **270**:5926-5931.
- Schaloske R, Malchow D: **Mechanism of cAMP-induced Ca<sup>2+</sup>-influx in Dictyostelium: role of phospholipase A<sub>2</sub>.** *Biochem J* 1997, **327**:233-238.
- Schlatterer C, Gollnick F, Schmidt E, Meyer R, Knoll G: **Challenge with high concentrations of cyclic AMP induces transient changes in the cytosolic free calcium concentration in Dictyostelium discoideum.** *J Cell Sci* 1994, **107**:2107-2115.
- Saran S, Nahao H, Tasaka M, Iida H, Tsuji FI, Nanjundiah V, Takeuchi I: **Intracellular free calcium level and its response to cAMP stimulation in developing Dictyostelium cells transformed with jellyfish apoaequorin.** *FEBS Lett* 1994, **337**:43-47.
- Sonnemann J, Knoll G, Schlatterer C: **cAMP-induced changes in the cytosolic free Ca<sup>2+</sup> concentration in Dictyostelium discoideum are light sensitive.** *Cell Calcium* 1997, **22**:65-74.
- Nebl T, Fisher PR: **Intracellular Ca<sup>2+</sup> signals in Dictyostelium chemotaxis are mediated exclusively by Ca<sup>2+</sup> influx.** *J Cell Sci* 1997, **110**:2845-2853.
- Rooney EK, Gross JD: **ATP-driven Ca<sup>2+</sup>/H<sup>+</sup>-antiport in acid vesicles from Dictyostelium.** *Proc Natl Acad Sci USA* 1992, **89**:8025-8029.
- Flaadt H, Jaworski E, Malchow D: **Evidence for two intracellular calcium pools in Dictyostelium – the cAMP-induced calcium influx is directed into a NBD-Cl<sup>-</sup> and 2, 5-di-(tert-butyl)-1, 4-hydroquinone-sensitive pool.** *J Cell Sci* 1993, **105**:1131-1135.
- Rooney EK, Gross JD, Satre M: **Characterisation of an intracellular Ca<sup>2+</sup> pump in Dictyostelium.** *Cell Calcium* 1994, **16**:509-522.
- Gröner M, Malchow D: **Calmodulin-antagonists inhibit vesicular Ca<sup>2+</sup>-uptake in Dictyostelium.** *Cell Calcium* 1996, **19**:105-111.
- Schaloske R, Sonnemann J, Malchow D, Schlatterer C: **Fatty acids induce release of Ca<sup>2+</sup> from acidosomal stores and activate capacitative Ca<sup>2+</sup>-entry in Dictyostelium discoideum.** *Biochem J* 1998, **332**:541-548.
- Böhme R, Bumann J, Aeckerle S, Malchow D: **A high affinity plasma membrane Ca<sup>2+</sup>-ATPase in Dictyostelium discoideum: its relation to cAMP-induced Ca<sup>2+</sup>-fluxes.** *Biochim Biophys Acta* 1987, **904**:125-130.
- Milne JL, Coukell MB: **Isolation and characterization of a plasma membrane calcium pump from Dictyostelium discoideum.** *Biochem J* 1988, **249**:223-230.
- Schlatterer C, Buravkov S, Zierold K, Knoll G: **Calcium-sequestering organelles of Dictyostelium discoideum: changes in element content during early development as measured by electron probe X-ray microanalysis.** *Cell Calcium* 1994, **16**:101-111.
- Traynor D, Milne JL, Insall RH, Kay RR: **Ca<sup>2+</sup> signalling is not required for chemotaxis in Dictyostelium.** *EMBO J* 2000, **19**:4846-4854.
- Schlatterer C, Schaloske R: **Calmidazolium leads to an increase in the cytosolic Ca<sup>2+</sup>-concentration in Dictyostelium discoideum by induction of Ca<sup>2+</sup>-release from intracellular stores and influx of extracellular Ca<sup>2+</sup>.** *Biochem J* 1996, **313**:661-667.
- Malchow D, Schaloske R, Schlatterer C: **An increase in cytosolic Ca<sup>2+</sup>-delays cAMP oscillations in Dictyostelium cells.** *Biochem J* 1996, **319**:323-327.

33. Dammann H, Hellstern S, Husain Q, Mutzel R: **Primary structure, expression and developmental regulation of a Dictyostelium calcineurin A homologue.** *Eur J Biochem* 1996, **238**:391-399.
34. Gerisch G, Hess B: **cAMP-controlled oscillations in suspended Dictyostelium cells: their relation to morphogenetic cell interactions.** *Proc Natl Acad Sci USA* 1974, **71**:2118-2122.
35. Gerisch G, Wick U: **Intracellular oscillations and release of cAMP from Dictyostelium cells.** *Biochem Biophys Res Commun* 1975, **65**:364-370.
36. Bumann J, Malchow D, Wurster B: **Oscillations of Ca<sup>2+</sup>-concentration during the cell differentiation of Dictyostelium discoideum. Their relation to oscillations in cAMP and other components.** *Differentiation* 1986, **31**:85-91.
37. Lundberg GA, Newell PC: **Membrane-associated phosphoinositidase C activity in Dictyostelium discoideum.** *FEBS Lett* 1990, **270**:181-183.
38. Cubitt AB, Firtel RA: **Characterization of phospholipase activity in Dictyostelium discoideum – identification of a Ca<sup>2+</sup>-dependent polyphosphoinositidase-specific phospholipase-C.** *Biochem J* 1992, **283**:371-378.
39. Van Dijken P, Bergsma JCT, Van Haastert PJM: **Phospholipase-C-independent inositol 1, 4, 5-triphosphate formation in Dictyostelium cells.** *Eur J Biochem* 1997, **244**:113-119.
40. Dunbar AJ, Wheldrake JF: **A calcium and calmodulin-dependent protein kinase present in differentiating Dictyostelium discoideum.** *FEMS Microbiol Lett* 1994, **115**:113-118.
41. Zhu Q, Clarke M: **Association of calmodulin and an unconventional myosin with the contractile vacuole complex of Dictyostelium discoideum.** *J Cell Biol* 1992, **118**:347-358.
42. Hill TD, Campos-Gonzalez R, Kindmark H, Boyton AL: **Inhibition of inositoltriphosphate-stimulated calcium mobilization by calmodulin antagonists in rat liver epithelial cells.** *J Biol Chem* 1988, **263**:16479-16484.
43. Peters C, Mayer A: **Ca<sup>2+</sup>-calmodulin signals the completion of docking and triggers a late step of vacuole fusion.** *Nature* 1998, **396**:575-580.
44. Moniakis J, Coukell MB, Janiec A: **Involvement of the Ca<sup>2+</sup>-ATPase PAT1 and the contractile vacuole in calcium regulation in Dictyostelium discoideum.** *J Cell Sci* 1999, **112**:405-414.
45. Clyde-Smith J, Silins G, Gartside M, Grimmond S, Etheridge M, Apolloni A, Hayward N: **Characterization of RasGRP2, a plasma membrane-targeted, dual specificity Ras factor.** *J Biol Chem* 2000, **275**:32260-32267.
46. Chen MY, Long Y, Devreotes PN: **A novel cytosolic regulator, pianissimo, is required for chemoattractant receptor and G-protein-mediated activation of the 12 transmembrane domain adenylyl cyclase in Dictyostelium.** *Genes Dev* 1997, **11**:3218-3231.
47. Lee S, Parent CA, Insall R, Firtel RA: **A novel Ras-interacting protein required for chemotaxis and cyclic adenosine monophosphate signal relay in Dictyostelium.** *Mol Biol Cell* 1999, **10**:2829-2845.
48. Schaloske R, Schlatterer C, Malchow D: **A Xestospongine C-sensitive Ca<sup>2+</sup>-store is required for cAMP-induced Ca<sup>2+</sup>-influx and cAMP oscillations in Dictyostelium.** *J Biol Chem* 2000, **275**:8404-8408.
49. Wayman GA, Impey S, Storm DR: **Ca<sup>2+</sup>-inhibition of type III adenylyl cyclase in vivo.** *J Biol Chem* 1995, **270**:21480-21486.
50. Sonnemann J, Aichele A, Schlatterer C: **Dissection of the cAMP-induced cytosolic calcium response in Dictyostelium discoideum: the role of cAMP receptor subtypes and G-protein subunits.** *FEBS Lett* 1998, **436**:271-276.
51. Meima ME, Schaap P: **Fingerprinting of adenylyl cyclase activities during Dictyostelium development indicates a dominant role for adenylyl cyclase B in terminal differentiation.** *Dev Biol* 1999, **212**:182-190.
52. Drayer AL, Van der Kaay J, Mayr GW, Van Haastert PJM: **Role of Phospholipase C in Dictyostelium – formation of inositol 1, 4, 5-triphosphate and normal development in cells lacking phospholipase C activity.** *EMBO J* 1994, **13**:1601-1609.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

