

# Independent Association Between Lower Level of Social Support and Higher Coagulation Activity Before and After Acute Psychosocial Stress

PETRA H. WIRTZ, PhD, LAURA S. REDWINE, PhD, ULRIKE EHLERT, PhD, AND ROLAND VON KANEL, MD

**Objective:** To investigate the relationship between social support and coagulation parameter reactivity to mental stress in men and to determine if norepinephrine is involved. Lower social support is associated with higher basal coagulation activity and greater norepinephrine stress reactivity, which in turn, is linked with hypercoagulability. However, it is not known if low social support interacts with stress to further increase coagulation reactivity or if norepinephrine affects this association. These findings may be important for determining if low social support influences thrombosis and possible acute coronary events in response to acute stress. We investigated the relationship between social support and coagulation parameter reactivity to mental stress in men and determined if norepinephrine is involved. **Methods:** We measured perceived social support in 63 medication-free nonsmoking men (age (mean  $\pm$  standard error of the mean) = 36.7  $\pm$  1.7 years) who underwent an acute standardized psychosocial stress task combining public speaking and mental arithmetic in front of an audience. We measured plasma D-dimer, fibrinogen, clotting Factor VII activity (FVII:C), and plasma norepinephrine at rest as well as immediately after stress and 20 minutes after stress. **Results:** Independent of body mass index, mean arterial pressure, and age, lower social support was associated with higher D-dimer and fibrinogen levels at baseline ( $p < .012$ ) and with greater increases in fibrinogen ( $\beta = -0.36, p = .001; \Delta R^2 = .12$ ), and D-dimer ( $\beta = -0.21, p = .017; \Delta R^2 = .04$ ), but not in FVII:C ( $p = .83$ ) from baseline to 20 minutes after stress. General linear models revealed significant main effects of social support and stress on fibrinogen, D-dimer, and norepinephrine ( $p < .035$ ). Controlling for norepinephrine did not change the significance of the reported associations between social support and the coagulation measures D-dimer and fibrinogen. **Conclusions:** Our results suggest that lower social support is associated with greater coagulation activity before and after acute stress, which was unrelated to norepinephrine reactivity. **Key words:** social support, coagulation, clotting factor VII:C, D-dimer, fibrinogen, psychological stress.

**CVD** = cardiovascular disease; **ACS** = acute coronary syndromes; **BMI** = body mass index; **MAP** = mean arterial pressure; **BP** = blood pressure; **SD** = standard deviation; **AUC** = area under the curve; **TSST** = Trier Social Stress Test; **FVII:C** = clotting Factor VII activity; **FVIII:C** = clotting Factor VIII activity.

## INTRODUCTION

Social support is a context-specific advocative interpersonal process centered on the reciprocal exchange of information on one or more of the following three classes: information leading the person to believe that he or she (1) is cared for and loved (2), esteemed and valued (3), and belongs to a network of communication and mutual obligation (1,2). In the literature, measures of perceived social support are more commonly used than measures of received social support (3). Perceived social support is conceptualized as the subjective appraisal of the degree of match between the amount and type of support needed and the amount and type of support available, or the perception that support would be available if needed (3).

Accumulating evidence suggests that social support is associated with health outcomes (2). For example, poor social support prospectively increases the risk of coronary artery disease (CAD) (3–6) with a relative risk two- to three-fold and a robust effect observed even after controlling for conventional (e.g., age, body

mass index (BMI), and high blood pressure (BP)), social, and behavioral cardiovascular risk factors (4–7).

Previous research addressed the biological pathways through which low social support increases cardiovascular risk or, alternatively, high social support enhances health. Based on the effects of social support on physiological processes implicated in cardiovascular disease, the so-called “social support-reactivity hypothesis” has been proposed. This view posits that social support maintains cardiovascular health by reducing psychobiological reactivity to stressors, thereby acting as a “stress buffering” variable (8,9). Evidence suggests that, in healthy individuals, social support attenuates psychological and physiological stress responses as indexed by cardiovascular, autonomic, and hypothalamo-pituitary-adrenal (HPA) responses (8,10–13). In hypertensive and normotensive middle-aged men, greater levels of perceived social support were associated with reduced catecholamines reactivity to acute psychosocial stress (14). Notably, the accumulation of physiological hyperreactivity to stress throughout a life span that may occur in individuals with low social support has been proposed to enhance CAD risk (15–18).

Coagulation and fibrinolysis are implicated as stress-reactive physiological systems important in the development of CAD and acute coronary syndromes (ACS) (19). Patients whose ACS had been triggered by intense emotions showed greater platelet activation in response to psychological stress compared with patients without emotion-triggering ACS (20). In healthy individuals, acute mental stress activates both the coagulation and the fibrinolysis components of hemostasis to result in net hypercoagulability (19,21). Acute psychological stress-induced increases in several hemostatic parameters, such as clotting Factor VII activity (FVII:C), clotting Factor VIII activity (FVIII:C) and FXII:C, thrombin-antithrombin complex, fibrinogen, von Willebrand Factor, and D-dimer levels, have been observed (19).

In contrast to stress effects, higher social support may reduce coagulation activity. Greater social support as assessed

From the Department of Clinical Psychology and Psychotherapy (P.H.W., U.E.), University of Zurich, Switzerland; Department of Medicine (L.S.R.), Cardiology Branch, University of California, San Diego, California; and the Department of General Internal Medicine (R.v.K.), Bern University Hospital, Inselspital, and University of Bern, Bern, Switzerland.

Address correspondence and reprint requests to Petra H. Wirtz, Department of Clinical Psychology and Psychotherapy, University of Zurich, Binzmühlestrasse 14, Box 26, CH-8050 Zurich, Switzerland. E-mail: p.wirtz@psychologie.uzh.ch

Supported by research Grants 56233203 and 56233204 from the University of Zurich (P.H.W.) and by a research grant from the University of Bern (R.v.K.).

via social integration, social isolation, and social network is associated with lower amount of fibrinogen (22–27) even when controlling for cardiovascular risk factors (22,24,26,27). However, the role of social support as a buffer of acute stress for hemostasis is not straightforward. In a small sample of 27 healthy middle-aged men, we recently did not find significant associations between social support as assessed by the German Social Support Questionnaire (F-SozU) and stress-induced changes in coagulation parameters (28). However, in that study, due to the small sample size, we did not control for age, BMI, and BP. In contrast, a study by Steptoe and co-workers assessed associations between social support measures and fibrinogen stress reactivity (27) and found higher fibrinogen levels in socially isolated participants compared with nonisolated participants before and after acute mild psychological stress (27).

Catecholamines play a role in stress-induced hemostatic changes because catecholamine infusion elicited hypercoagulability in healthy persons (29). In addition, we recently found associations between epinephrine and FVIII:C and between norepinephrine and fibrinogen under resting conditions, independent of age, BMI, and mean arterial pressure (MAP) in middle-aged hypertensive and normotensive men (30). Moreover, in the same study, stress-induced changes in D-dimer were predicted by norepinephrine stress change (30). In another study, we found that the stress-induced increase in norepinephrine significantly correlated with thrombin formation (31). However, it is unknown whether lower social support is associated with greater catecholamine stress reactivity, which in turn could induce hypercoagulability. More specifically, the role of norepinephrine in stress-induced D-dimer changes in those with low social support has not previously been examined.

The aim of this study was to extend previous research investigating whether perceived social support is associated with several coagulation parameter levels at rest and throughout a potent acute psychosocial stressor in a sizeable group of medication-free nonsmoking men. We hypothesized that lower social support is associated with higher baseline levels and greater increases in response to stress in the coagulation measures fibrinogen and FVII:C, and the hypercoagulability marker D-dimer. Moreover, we examined whether such associations were moderated or mediated by norepinephrine levels.

## METHODS

### Study Participants

The study is part of a project studying the biological response in general and coagulation activation in particular to acute psychosocial stress in men as previously described (32,33). Whereas our previous report addressed the acute stress-induced increases of the coagulation parameters D-dimer and fibrinogen in relationship to age (33), in the present study, we specifically examined the relationship of social support to acute stress-induced coagulation changes. In the present study, we also tested whether the relationship between social support and coagulation activation to stress would be modulated by norepinephrine.

The Ethics Committee of the State of Zurich, Switzerland, formally approved the research protocol. Of the original 66 subjects, three persons had to be excluded because of incomplete blood samples for coagulation param-

eters. The final study sample consisted of 63 subjects who provided their written informed consent. The study was conducted between April 2004 and August 2005. We intentionally recruited nonsmoking men between the ages of 20 and 65 years who were in excellent physical and mental health confirmed by an extensive health questionnaire (34) and telephone interview. Specific exclusion criteria were obtained by the subjects' self-report and were as follows: regular strenuous exercise; alcohol and illicit drug abuse; any heart disease, varicosis or thrombotic diseases; elevated blood sugar and diabetes; elevated cholesterol, liver, and renal diseases; chronic obstructive pulmonary disease; allergies and atopic diathesis, rheumatic diseases; and current infectious diseases. In addition, participants were included only if they reported taking no prescribed and/or over-the-counter medication, either regularly or occasionally, and if their BP was in the normotensive or moderately hypertensive range (systolic BP <160 mm Hg and diastolic BP <100 mm Hg). When the personal or medication history was not conclusive, the subject's primary care physician was contacted for verification.

### Assessment of Social Support

The first part of the Berlin Social Support Scale consists of 17 items assessing perceived social support (PSS), support seeking (SS), and need for support (35). Previous literature on social support and cardiovascular disease as well as our own previous findings of greater catecholamine stress reactivity with increasing PSS led us to use the 8-item PSS subscale composed of the two subscales "emotional support" and "instrumental support" in the present study (3,14). Using a 4-point rating scale ranging from 1 (completely wrong) to 4 (completely right), we asked the participants whether they agree with certain statements on their perception of social support (e.g., "there are people who help me if I need help"; "there are people cheering me up when I am sad"). The PSS renders scores between 1 (minimum score) and 4 (maximum score). Higher scores mean higher PSS. Cronbach's  $\alpha$  ( $n = 437$ ) is 0.83 for the PSS subscale (35).

### Stress Protocol

Subjects were tested between 2 PM and 4 PM. They abstained from physical exercise, alcohol, and caffeinated beverages for at least 24 hours before testing. We used the Trier Social Stress Test (TSST) combining a 5-minute preparation phase followed by a 5-minute mock job interview, and 5-minute mental arithmetic task in front of an audience (36). The TSST evokes reliable physiological responses across different biological systems, including coagulation factors also investigated in the present study (37). During recovery, the subjects remained seated in a quiet room for 40 minutes.

Blood for coagulation and norepinephrine measures was obtained immediately before stress, immediately after stress, and 20 minutes after stress. BP was measured immediately before and 40 minutes after stress by sphygmomanometry (Omron 773, Omron Healthcare Europe B.V., Hoofddorp, Netherlands) and MAP was calculated by the formula ( $2/3 \times$  mean diastolic BP) + (1/3 mean systolic BP).

### Biochemical Analyses

Venous blood was drawn through an indwelling forearm catheter into polypropylene tubes containing 3.8% sodium citrate and centrifuged at 2000 g for 20 minutes at 4°C. The obtained plasma sample was immediately aliquoted in polypropylene tubes and frozen at -80°C. All analyses of coagulation factors used the BCS Coagulation Analyzer (Dade Behring, Liederbach, Germany). Determination of FVII:C used standard coagulometric methods, factor-deficient standard human plasma, and reagents (Dade Behring); plasma fibrinogen was determined using a modified Clauss method (Multifibren U, Dade Behring). Plasma D-dimer was measured by means of an enzyme-linked immunosorbent assay (Asserachrom Stago, Asnières, France). Inter- and intra-assay coefficients of variation were <10% for all coagulation measures.

Blood samples for measurement of plasma norepinephrine were drawn into ethylenediaminetetraacetic acid-coated monovettes (Sarstedt, Numbrecht, Germany), and immediately centrifuged for 10 minutes at 2000 g; plasma was stored at -80°C until analysis. Plasma norepinephrine was determined by means of high-performance liquid chromatography and electrochemical de-

tection after liquid-liquid extraction (38,39). The limit of detection was 10 pg/ml. Inter- and intra-assay variance was <5%. To reduce error variance caused by imprecision of the intra-assay, all samples from one subject were analyzed in the same run.

### Statistical Analyses

Data were analyzed using SPSS (version 13.0) statistical software package (SPSS Inc., Chicago IL). All tests were two-tailed with level of significance of  $p \leq .05$  and level of statistical trends of  $p \leq .10$ . Using the trapezoid formula, we calculated areas under the total response curves, expressed as area under the measured time points with respect to ground for all coagulation measures and norepinephrine (40). Before statistical analyses, all data were tested for normality using the Kolmogorov-Smirnov test. Coagulation values and coagulation areas under the curve (AUCs) were logarithmically transformed to achieve normal distributions. For clarity, we provide untransformed data.

To assess the associations between social support and coagulation activity at baseline and after stress, we first calculated linear regression analyses with the respective coagulation measure as the dependent variable and social support score as a continuous independent variable. We used coagulation baseline measures as the dependent variables to assess associations between social support and coagulation activity at rest. We employed AUC measures of the coagulation parameters to assess the associations between social support and stress-induced coagulation changes. In light of previously reported associations between BMI, MAP, and age with coagulation parameters at rest and in response to stress (30,33,41), we controlled for BMI, MAP, and age. We entered these parameters as predictors in all analyses. All independent variables were simultaneously forced into the regression equations. The optimal total sample size to predict stress reactivity in coagulation parameters was  $n = 59$  for detecting a medium-to-large effect size of  $f^2 = 0.25$  in multiple regression analyses with a power of 0.80 using four predictors. Second, we further tested regression results by performing general linear models with repeated measures for each coagulation parameter as the dependent variable and with social support as the continuous independent variable. In these analyses, we again controlled for BMI, MAP, and age. For illustrative purposes, we categorized the study group based on their social support scores into four groups of subjects with increasing levels of social support termed as follows: lowest (2.75–3.38,  $n = 11$ ), lower (3.50–3.63,  $n = 16$ ), higher (3.75–3.88,  $n = 25$ ), and highest social support scores (4–4,  $n = 11$ ).

To test for associations between social support and norepinephrine stress reactivity, we used general linear modeling with norepinephrine levels as repeated dependent factor and levels of social support as the continuous independent factor, at the same time controlling for age, BMI, and MAP. To assess whether associations between social support and coagulation activity at baseline and after stress are related to norepinephrine levels, we again calculated regression analyses. As dependent variables, we entered those coagulation measures that were significantly associated with social support in the linear model. As independent variables, we simultaneously entered social support and norepinephrine to test for a mediation effect of norepinephrine. To test for a moderating effect of norepinephrine, we simultaneously entered social support, norepinephrine, as well as their interaction term (social support  $\times$  norepinephrine) as independent variables. We always controlled for age, BMI, and MAP. Interaction terms were computed on Z-transformed data rendering means of 0 and standard deviations (SD) of 1. All regression analyses including interaction terms were performed on Z-transformed data. We calculated Pearson's product-moment correlations to assess the associations between social support, age, BMI, and MAP.

## RESULTS

### Subject Characteristics

Social support scores of our 63 study participants ranged between 2.75 and 4 (mean  $\pm$  SD = 3.66  $\pm$  0.32). The mean age was 37  $\pm$  13.7 years, the mean BMI was 24.8  $\pm$  3.04, and MAP was 94.7  $\pm$  10.62 mm Hg. Age, MAP, and BMI were intercorrelated ( $p < .001$ ) but they did not correlate with social support ( $p > .21$ ).

### Relationship Between Social Support and Coagulation Activity at Rest and in Response to Stress Regression Analyses

#### *Social Support and Coagulation Activity at Rest D-Dimer at Rest*

Lower social support scores were significantly associated with higher D-dimer levels at rest ( $\beta = -0.22$ ,  $p = .012$ ;  $\Delta R^2 = .05$ ) independent of MAP ( $p = .11$ ), BMI ( $\beta = 0.47$ ,  $p < .001$ ;  $\Delta R^2 = .15$ ), and age ( $\beta = 0.51$ ,  $p < .001$ ;  $\Delta R^2 = 0.17$ ) with the total model explaining 60% of the variance in D-dimer resting levels.

#### *Fibrinogen at Rest*

Higher fibrinogen at rest was significantly associated with lower social support ( $\beta = -0.35$ ,  $p < .001$ ;  $\Delta R^2 = .11$ ) independent of MAP ( $p = .90$ ), BMI ( $\beta = 0.42$ ,  $p < .001$ ;  $\Delta R^2 = .12$ ), and age ( $\beta = 0.33$ ,  $p = .004$ ;  $\Delta R^2 = .08$ ) with the total model explaining 52% of the variance in fibrinogen resting levels.

#### *FVII:C at Rest*

Social support was not significantly associated with FVII:C at rest ( $p = .96$ ).

### Social Support and Coagulation Levels Between Rest and 20 Minutes After Stress

#### *D-Dimer AUC*

Lower social support was associated with higher D-dimer AUC ( $\beta = -0.21$ ,  $p = .017$ ;  $\Delta R^2 = 0.04$ ) independent of MAP ( $p = .07$ ), BMI ( $\beta = 0.42$ ,  $p < .001$ ;  $\Delta R^2 = .13$ ), and age ( $\beta = 0.56$ ,  $p < .001$ ;  $\Delta R^2 = .21$ ). The whole model explained 59% of the variance in D-dimer AUC.

#### *Fibrinogen AUC*

Independent of MAP ( $p = .95$ ), BMI ( $\beta = 0.38$ ,  $p = .001$ ;  $\Delta R^2 = .10$ ), and age ( $\beta = 0.34$ ,  $p = .005$ ;  $\Delta R^2 = .08$ ), lower social support was associated with higher fibrinogen AUC ( $\beta = -0.36$ ,  $p = .001$ ;  $\Delta R^2 = 0.12$ ). The whole model explained 48% of the observed variance in fibrinogen AUC.

#### *FVII:C AUC*

There were no associations between social support and AUC of FVII:C ( $p = .83$ ).

### General Linear Models

Across all subjects, the TSST elicited significant increases in D-dimer ( $p = .009$ ,  $f = 0.34$ ), fibrinogen ( $p = .012$ ,  $f = 0.32$ ), and FVII:C ( $p = .046$ ,  $f = 0.32$ ) from baseline to immediately after stress at the same time controlling for age, BMI, and MAP. To validate the results from the above regression analyses, we applied general linear models with repeated measures of coagulation factors as dependent variables and social support as a continuous independent variable. After controlling for BMI, MAP, and age, there were significant main effects for social support for repeated D-dimer and fibrinogen levels (D-dimer:  $F(1,58) = 6.0$ ,  $p = .018$ ,  $f = 0.31$ ,

Figure 1A; fibrinogen:  $F(1,58) = 13.8, p < .001, f = 0.47$ , Figure 1B) whereas there were no significant interactions between stress and social support ( $p > .58$ ). Neither main effects nor interaction effects were observed in terms of repeated measurements of FVII:C ( $p > .24$ ).

For illustrative purposes, Figure 1 (panels A and B) shows coagulation responses to the TSST in four groups of subjects with increasing levels of social support (lowest: 2.75–3.38,  $n = 11$ ; lower: 3.50–3.63,  $n = 16$ ; higher: 3.75–3.88,  $n = 25$ ), and highest social support scores (4–4,  $n = 11$ ) based on quartiles (note that subject numbers differ in the four groups because individual PSS values were not balanced above and below the particular cut-off values).




Figure 1. Prothrombotic factor levels in four groups of subjects with lowest (2.75–3.38,  $n = 11$ ), lower (3.50–3.63,  $n = 16$ ), higher (3.75–3.88,  $n = 25$ ), and highest social support (SS) scores (4–4,  $n = 11$ ). (A, B). Values are mean  $\pm$  standard error of the mean. Stress reactivity of D-dimer (A) and fibrinogen (B) across all 63 subjects. We applied general linear models with repeated measures of coagulation factors as dependent variables and social support as continuous independent variable at the same time controlling for body mass index, mean arterial pressure, and age in all analyses. The main effect of social support was significant in terms of D-dimer  $F(1,58) = 6.0, p = .018, f = 0.31$ , (A) and fibrinogen  $F(1,58) = 13.8, p < .001, f = 0.47$ , (B). The figure shows these data for illustrative purposes by four categories of SS scores. TSST = Trier Social Stress Test.

## Norepinephrine, Social Support, and Coagulation Parameters

### Social Support and Norepinephrine Stress Reactivity

The TSST stimulated significant increases in norepinephrine levels ( $F(1/61) = 94.9, p < .001$ ). General linear modeling with norepinephrine as repeated dependent variable and social support as continuous independent variable revealed that higher social support was associated with lower norepinephrine levels before and after stress (main effect:  $F(1/57) = 4.65, p = .035, f = 0.27$ , Figure 2), at the same time controlling for age, BMI, and MAP. There was no interaction between social support and stress ( $p = .33$ ).

### Norepinephrine and Associations Between Social Support and Coagulation Parameters

Table 1 depicts regression results for associations between norepinephrine, social support, and coagulation parameters at rest and in response to stress.

#### At Rest

Regression analyses after controlling for BMI, MAP, and age revealed that resting norepinephrine levels were not significantly related to any of the coagulation parameters ( $p > .60$ ), and was determined to not be a significant factor in the associations between social support and resting levels of D-dimer and fibrinogen (Table 1). This suggests that resting norepinephrine levels do not mediate the observed associations between social support and resting levels of D-dimer and fibrinogen. Additionally, entering baseline norepinephrine, levels, social support, plus the norepinephrine-by-social sup-




Figure 2. Norepinephrine levels in four groups of subjects with lowest (2.75–3.38,  $n = 11$ ), lower (3.50–3.63,  $n = 16$ ), higher (3.75–3.88,  $n = 25$ ), and highest social support (SS) scores (4–4,  $n = 11$ ). Values are mean  $\pm$  standard error of the mean. Stress reactivity of norepinephrine across all 63 subjects. We applied general linear models with repeated measures of norepinephrine as dependent variables and social support as continuous independent variable at the same time controlling for body mass index, mean arterial pressure, and age. Higher social support was associated with lower norepinephrine levels before and after stress (main effect:  $F(1,57) = 4.65, p = .035, f = 0.27$ ). Figure 2 shows these data for illustrative purposes by four categories of SS scores. TSST = Trier Social Stress Test.

**TABLE 1. Hierarchical Regression Analyses for Associations Between Social Support and Coagulation Parameters Controlling for Norepinephrine**

| Variable               | Standardized $\beta$ Coefficient | <i>t</i> | <i>p</i> | R <sup>2</sup> Change |
|------------------------|----------------------------------|----------|----------|-----------------------|
| D-dimer at baseline    |                                  |          |          |                       |
| Age                    | 0.50                             | 4.89     | <.001    | .17                   |
| BMI                    | 0.47                             | 4.63     | <.001    | .15                   |
| MAP                    | -0.16                            | -1.61    | .11      | .02                   |
| NEPI at baseline       | 0.02                             | 0.20     | .84      | .00                   |
| SS                     | -0.22                            | -2.47    | .017     | .04                   |
| Fibrinogen at baseline |                                  |          |          |                       |
| Age                    | 0.33                             | 2.89     | .005     | .07                   |
| BMI                    | 0.41                             | 3.69     | .001     | .12                   |
| MAP                    | 0.01                             | 0.10     | .92      | .00                   |
| NEPI at baseline       | 0.05                             | 0.54     | .60      | .00                   |
| SS                     | -0.33                            | -3.45    | .001     | .10                   |
| D-dimer AUC            |                                  |          |          |                       |
| Age                    | 0.55                             | 5.31     | <.001    | .20                   |
| BMI                    | 0.42                             | 4.12     | <.001    | .12                   |
| MAP                    | -0.19                            | -1.88    | .07      | .03                   |
| NEPI AUC               | 0.03                             | 0.36     | .72      | .00                   |
| SS                     | -0.20                            | -2.29    | .026     | .04                   |
| Fibrinogen AUC         |                                  |          |          |                       |
| Age                    | 0.33                             | 2.78     | .007     | .07                   |
| BMI                    | 0.37                             | 3.22     | .002     | .09                   |
| MAP                    | 0.00                             | 0.02     | .98      | .00                   |
| NEPI AUC               | 0.07                             | 0.68     | .50      | .00                   |
| SS                     | -0.34                            | -3.42    | .001     | .11                   |

BMI = body mass index; MAP = mean arterial blood pressure; NEPI = norepinephrine; AUC = area under the curve; SS = social support score.

port interaction into the regression equation revealed that resting norepinephrine levels did not moderate the relationship between social support and both D-dimer and fibrinogen ( $p$ 's > .69).

### Stress Reactivity

To determine if norepinephrine secretion mediates between social support and stress reactivity of coagulation parameters D-dimer and fibrinogen, regression analyses were recalculated with norepinephrine AUC entered as an independent variable (Table 1). To test for a moderation effect of norepinephrine, its interaction with social support was additionally entered as an independent variable. Controlling for norepinephrine AUC did not significantly affect the observed associations between social support and AUC of D-dimer and fibrinogen. Thus, neither norepinephrine AUC alone nor its interaction with social support were significantly associated with any of the coagulation parameter AUCs ( $p > .50$ ). In sum, norepinephrine does not seem to be a mediator or a moderator of social support effects on coagulation activity.

Notably, results of the PSS subscales "instrumental support" and "emotional support" were similar to the combined PSS scale in all analyses (data not shown).

## DISCUSSION

Research suggested that poor social support is associated with health risks (2), such as CAD (3–6). The mechanisms

involved are not known, although overactivation of hemostasis might be one contributing factor. The main objective of the present study was to investigate whether social support is associated with elevated coagulation factor levels at rest and throughout an acute stress process. The coagulation factors fibrinogen and D-dimer were examined at baseline, representing resting activity, and also for the AUC (i.e., immediately before stress, immediately after stress, and 20 minutes after stress), representing the total stress-response course. The results of our study suggest that low social support is associated with increased fibrinogen and D-dimer levels at rest, and that acute stress and low social support are independently associated with an increased AUC. This may indicate that individuals with reduced social support run an even higher atherothrombotic risk during acute stress. Specifically, effects sizes of stress-induced coagulation increases suggest clinical importance. A prothrombotic state of the blood promotes atherosclerosis development and, after rupture of an atherosclerotic plaque, thrombotic occlusion of a coronary artery leading to an ACS (42,43).

Even though FVII:C was responsive to the TSST, we did not find a significant association between social support and FVII:C both at baseline and throughout the stress period. Clotting factor FVII belongs to the extrinsic system of blood coagulation, whereas fibrinogen is the precursor of fibrin further down in the coagulation cascade after intrinsic and extrinsic coagulation pathways have merged (21). Therefore, it could be that social support is more strongly related to factors involved in the intrinsic pathway of blood coagulation, which reasoning we did, however, not address in the present study. After clot formation, the fibrinolysis system degrades fibrin, whereby fibrin degradation products such as D-dimer are formed. Because it indicates activation of the entire coagulation system (i.e., the step of fibrin formation) and also the fibrinolysis system (i.e., the step of fibrin dissolution), D-dimer is termed a hypercoagulability marker (44). Hypercoagulability markers are more sensitive to change than are individual hemostatic factors (45), providing one explanation for why D-dimer emerged as a significant correlate of social support whereas FVII:C did not. Our findings correspond to previous studies that social support scores were negatively associated with resting fibrinogen levels (19,21) now extending these findings to D-dimer. These may be important findings in relationship to health. Elevated fibrinogen levels are associated with cardiovascular disease risk (46), and are related to both atherogenesis and thrombogenesis. For each SD increase in fibrinogen above the mean, it was suggested that there is an 84% increase in the 5-year risk of ischemic heart disease (47). In addition, increased D-dimer levels are associated with greater risk of myocardial infarction (48), cerebrovascular events (49), and peripheral arterial disease (50).

Acute mental stress activates both the coagulation and the fibrinolysis components of hemostasis to result in net hypercoagulability (19,21). Increased social support has been suggested to act as a buffer to attenuate physiological stress responses, as indexed by cardiovascular, autonomic, and HPA

responses (8,10–13). However, our present findings did not suggest an interaction between social support and stress for hemostatic measures, but social support and stress are independent factors and therefore may be additive. Higher social support predicted lower D-dimer and fibrinogen AUC from baseline to 20 minutes after the stress task. These findings support a previous study showing that higher fibrinogen levels were found in socially isolated participants both before and after a color-word and a mirror-tracing task (27). The present study extends those results to include D-dimer, which is involved with both coagulation and fibrinolysis. Furthermore, an interpersonal stressor, the TSST, was used in the present research, which expands the range of psychological stress that is found to generate such hemostatic responses. Catecholamines play a role in stress-induced hemostatic changes (30), but their role in social support-related changes in coagulation parameters is unclear. To examine if norepinephrine mediates or moderates the association between social support and hemostatic changes at rest and in response to stress, fibrinogen and D-dimer were measured at baseline and throughout the stress task (baseline through 20 minutes post task). Resting norepinephrine levels and the interaction of norepinephrine baseline levels with social support did not predict levels of D-dimer and fibrinogen, suggesting that norepinephrine is not associated with hemostatic reactions to stress in relationship to social support.

Future studies are needed to examine additional pathways that may mediate social support and health, besides the sympathetic reactivity hypothesis, antistress hormones, inflammatory responses, and sympatho-vagal balance may play roles as alternative mediators. For example, oxytocin was found to have potential antistress effects and when combined with social support, there was a reduced cortisol response along with increased calmness and decreased anxiety during the TSST (13). In turn, cortisol changes during stress are associated with total fibrin formation (51), suggesting a potential alternate pathway in which social support may buffer stress-induced hemostatic activity and ultimately health.

Social networks were associated with reduced inflammatory marker interleukin-6 responses in male Framingham Heart Study participants (52), and reactivity in interleukin-6 and D-dimer to stress were positively correlated (51). Additionally, in patients with coronary heart disease (CHD), elevated social support is associated with greater probability of sympatho-vagal balance (i.e., reduced heart rate, MAP, cardiac index, and increased high frequency heart rate variability) during recovery from mental stress (53). In contrast, social isolation was associated with decreased heart rate variability (54) suggesting a reduction in parasympathetic or vagal tone. In women with CHD, we previously observed elevated levels of fibrinogen in association with reduced vagal cardiac control (55).

There are also hypotheses about social relationships suggesting that they exhibit main effects on health but do not necessarily interact with stress (56). Lack of social support due to social isolation had a hazard ratio (HR) of 1.75 for mortality in patients with heart failure (57). Compared with

heart failure patients reporting a high social network, hospital readmission was more frequent among those who had moderate social networks (HR = 1.87; 95% Confidence Interval (CI) = 1.06–3.29;  $p < .05$ ) and low social networks (HR = 1.98; 95% CI = 1.07–3.68;  $p < .05$ ) (58). Similarly, loneliness was significantly associated with elevated systolic BP, but not so with differences in autonomic or endocrine reactivity to stress (59). Comparisons of persons reporting more types of contacts (parents, children, family members, and friends) with those reporting fewer types yielded age- and gender-adjusted HRs of 0.73 (95% CI = 0.64–0.82) for mortality and HRs of 0.75 (95% CI = 0.61–0.91) for CHD (60).

In summary, social support is associated with various factors that might also affect hemostasis and that warrant an investigation as potential pathways linking social support and hemostasis and ultimately health. Alternatively, social relationships may function as a main effect on health. As is often the case, both hypotheses are likely true depending on the situation and factors being measured.

Our study has several strengths, which include recruitment of a sizable group of apparently healthy and unmedicated subjects across a range of perceived social support levels in a natural unselected setting. However, the study has also its limitations. First, its cross-sectional nature does not allow us to interpret the direction of the social support-coagulation-stress link, although we feel it is unlikely that coagulation influences social support. Second, our findings were obtained in a sample of apparently healthy men with BP in the normotensive and mildly hypertensive range and may not be generalized to individuals with more severe hypertension and women. Third, we restricted our analyses to three coagulation parameters. Fourth, the significance of low scores on PSS remains to be elucidated and future studies should include assessments of a person's social network.

In conclusion, social support seems to be important for coagulation parameter levels, such as fibrinogen and D-dimer, which may affect thrombosis and atherosclerotic disease. Lower social support was associated with higher baseline levels of fibrinogen and D-dimer. In addition, lower social support was independently related to an increase of these hemostatic factors throughout the stress response period. Stress-related alterations in hemostasis might act as a potential trigger for the onset of ACS (19). Although social support did not seem to affect the magnitude of the coagulation parameter responses to stress, low social support may have added risk because of its association with greater overall coagulation parameter levels. Furthermore, social support does not seem to drive stress-associated changes in norepinephrine. Therefore, future studies are needed to examine the mechanisms that are involved in the relationship between social support and hemostasis factors in order to develop interventions that can reduce hemostatic risk factors of CHD related to low levels of social support.

## REFERENCES

1. Cobb S. Presidential Address—1976. Social support as a moderator of life stress. *Psychosom Med* 1976;38:300–14.

2. McEwen BS. Physiology and neurobiology of stress and adaptation: central role of the brain. *Physiol Rev* 2007;87:873–904.
3. Lett HS, Blumenthal JA, Babyak MA, Strauman TJ, Robins C, Sherwood A. Social support and coronary heart disease: epidemiologic evidence and implications for treatment. *Psychosom Med* 2005;67:869–78.
4. Strike PC, Steptoe A. Psychosocial factors in the development of coronary artery disease. *Prog Cardiovasc Dis* 2004;46:337–47.
5. Kuper H, Marmot M, Hemingway H. Systematic review of prospective cohort studies of psychosocial factors in the etiology and prognosis of coronary heart disease. *Semin Vasc Med* 2002;2:267–314.
6. Rozanski A, Blumenthal JA, Kaplan J. Impact of psychological factors on the pathogenesis of cardiovascular disease and implications for therapy. *Circulation* 1999;99:2192–217.
7. Andre-Petersson L, Engstrom G, Hedblad B, Janzon L, Rosvall M. Social support at work and the risk of myocardial infarction and stroke in women and men. *Soc Sci Med* 2007;64:830–41.
8. Christenfeld N, Gerin W. Social support and cardiovascular reactivity. *Biomed Pharmacother* 2000;54:251–7.
9. Lepore SJ. Problems and prospects for the social support-reactivity hypothesis. *Ann Behav Med* 1998;20:257–69.
10. Nausheen B, Gidron Y, Gregg A, Tissarchondou HS, Peveler R. Loneliness, social support and cardiovascular reactivity to laboratory stress. *Stress* 2007;10:37–44.
11. Seeman TE, McEwen BS. Impact of social environment characteristics on neuroendocrine regulation. *Psychosom Med* 1996;58:459–71.
12. Kirschbaum C, Klauer T, Filipp SH, Hellhammer DH. Sex-specific effects of social support on cortisol and subjective responses to acute psychological stress. *Psychosom Med* 1995;57:23–31.
13. Heinrichs M, Baumgartner T, Kirschbaum C, Ehlert U. Social support and oxytocin interact to suppress cortisol and subjective responses to psychosocial stress. *Biol Psychiatry* 2003;54:1389–98.
14. Wirtz PH, von Kanel R, Mohiyeddini C, Emini L, Ruedisueli K, Groessbauer S, Ehlert U. Low social support and poor emotional regulation are associated with increased stress hormone reactivity to mental stress in systemic hypertension. *J Clin Endocrinol Metab* 2006;91:3857–65.
15. Steptoe A, Willemsen G. Psychophysiological responsiveness in coronary heart disease. In: Stansfeld SA, Marmot MG, editors. *Stress and the Heart*. London: BMJ Books; 2002:168–180.
16. Linden W, Gerin W, Davidson K. Cardiovascular reactivity: status quo and a research agenda for the new millennium. *Psychosom Med* 2003;65:5–8.
17. McEwen BS. Protective and damaging effects of stress mediators. *N Engl J Med* 1998;338:171–9.
18. Matthews KA, Katholi CR, McCreath H, Whooley MA, Williams DR, Zhu S, Markovitz JH. Blood pressure reactivity to psychological stress predicts hypertension in the CARDIA study. *Circulation* 2004;110:74–8.
19. Thrall G, Lane D, Carroll D, Lip GY. A systematic review of the effects of acute psychological stress and physical activity on haemorrhology, coagulation, fibrinolysis and platelet reactivity: Implications for the pathogenesis of acute coronary syndromes. *Thromb Res* 2007;120:819–47.
20. Strike PC, Perkins-Porras L, Whitehead DL, McEwan J, Steptoe A. Triggering of acute coronary syndromes by physical exertion and anger: clinical and sociodemographic characteristics. *Heart* 2006;92:1035–40.
21. von Kanel R, Mills PJ, Fainman C, Dimsdale JE. Effects of psychological stress and psychiatric disorders on blood coagulation and fibrinolysis: a biobehavioral pathway to coronary artery disease? *Psychosom Med* 2001;63:531–44.
22. Davis MC, Swan PD. Association of negative and positive social ties with fibrinogen levels in young women. *Health Psychol* 1999;18:131–9.
23. Rosengren A, Wilhelmsen L, Welin L, Tsipogianni A, Teger-Nilsson AC, Wedel H. Social influences and cardiovascular risk factors as determinants of plasma fibrinogen concentration in a general population sample of middle aged men. *BMJ* 1990;300:634–8.
24. Orth-Gomer K, Rosengren A, Wilhelmsen L. Lack of social support and incidence of coronary heart disease in middle-aged Swedish men. *Psychosom Med* 1993;55:37–43.
25. Helminen A, Rankinen T, Vaisanen S, Rauramaa R. Social network in relation to plasma fibrinogen. *J Biosoc Sci* 1997;29:129–39.
26. Loucks EB, Berkman LF, Gruenewald TL, Seeman TE. Social integration is associated with fibrinogen concentration in elderly men. *Psychosom Med* 2005;67:353–8.
27. Steptoe A, Kunz-Ebrecht S, Owen N, Feldman PJ, Rumley A, Lowe GD, Marmot M. Influence of socioeconomic status and job control on plasma fibrinogen responses to acute mental stress. *Psychosom Med* 2003;65:137–44.
28. von Kanel R, Kudielka BM, Preckel D, Hanebuth D, Herrmann-Lingen C, Frey K, Fischer JE. Opposite effect of negative and positive affect on stress procoagulant reactivity. *Physiol Behav* 2005;86:61–8.
29. Preckel D, Von Kanel R. Regulation of hemostasis by the sympathetic nervous system: any contribution to coronary artery disease? *Heart Drug* 2004;4:123–30.
30. Wirtz PH, Ehlert U, Emini L, Rudisuli K, Groessbauer S, Mausbach BT, von Kanel R. The role of stress hormones in the relationship between resting blood pressure and coagulation activity. *J Hypertens* 2006;24:2409–16.
31. von Kanel R, Mills PJ, Ziegler MG, Dimsdale JE. Effect of beta2-adrenergic receptor functioning and increased norepinephrine on the hypercoagulable state with mental stress. *Am Heart J* 2002;144:68–72.
32. Wirtz PH, Siegrist J, Rimmel U, Ehlert U. Higher overcommitment to work is associated with lower norepinephrine secretion before and after acute psychosocial stress in men. *Psychoneuroendocrinology* 2008;33:92–9.
33. Wirtz PH, Redwine LS, Baertschi C, Spillmann M, Ehlert U, von Kanel R. Coagulation activity before and after acute psychosocial stress increases with age. *Psychosom Med* 2008;70:476–81.
34. von Kanel R, Kudielka BM, Abd-el-Razik A, Gander ML, Frey K, Fischer JE. Relationship between overnight neuroendocrine activity and morning haemostasis in working men. *Clin Sci (Lond)* 2004;107:89–95.
35. Schulz U, Schwarzer R. Social support and coping with illness: the Berlin Social Support Scales (BSSS). *Diagnostica* 2003;49:73–82.
36. Kirschbaum C, Pirke KM, Hellhammer DH. The “Trier Social Stress Test”—a tool for investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology* 1993;28:76–81.
37. von Kanel R, Preckel D, Zraggen L, Mischler K, Kudielka BM, Haerberli A, Fischer JE. The effect of natural habituation on coagulation responses to acute mental stress and recovery in men. *Thromb Haemost* 2004;92:1327–35.
38. Ehrenreich H, Schuck J, Stender N, Pilz J, Gefeller O, Schilling L, Poser W, Kaw S. Endocrine and hemodynamic effects of stress versus systemic CRF in alcoholics during early and medium term abstinence. *Alcohol Clin Exp Res* 1997;21:1285–93.
39. Smedes F, Kraak JC, Poppe H. Simple and fast solvent extraction system for selective and quantitative isolation of adrenaline, noradrenaline and dopamine from plasma and urine. *J Chromatogr* 1982;231:25–39.
40. Pruessner JC, Kirschbaum C, Meinlschmid G, Hellhammer DH. Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change. *Psychoneuroendocrinology* 2003;28:916–31.
41. Wirtz PH, Ehlert U, Emini L, Rudisuli K, Groessbauer S, von Kanel R. Procoagulant stress reactivity and recovery in apparently healthy men with systolic and diastolic hypertension. *J Psychosom Res* 2007;63:51–8.
42. Falk E, Fernandez-Ortiz A. Role of thrombosis in atherosclerosis and its complications. *Am J Cardiol* 1995;75:3B–11B.
43. Libby P. The molecular mechanisms of the thrombotic complications of atherosclerosis. *J Intern Med* 2008;263:517–27.
44. von Kanel R, Dimsdale JE. Fibrin D-dimer: a marker of psychosocial distress and its implications for research in stress-related coronary artery disease. *Clin Cardiol* 2003;26:164–8.
45. Bauer KA, Rosenberg RD. The pathophysiology of the prethrombotic state in humans: insights gained from studies using markers of hemostatic system activation. *Blood* 1987;70:343–50.
46. Koenig W. Fibrin(ogen) in cardiovascular disease: an update. *Thromb Haemost* 2003;89:601–9.
47. Heinrich J, Balleisen L, Schulte H, Assmann G, van de Loo J. Fibrinogen and factor VII in the prediction of coronary risk. Results from the PROCAM study in healthy men. *Arterioscler Thromb* 1994;14:54–9.
48. Ridker PM, Hennekens CH, Cerskus A, Stampfer MJ. Plasma concentration of cross-linked fibrin degradation product (D-dimer) and the risk of future myocardial infarction among apparently healthy men. *Circulation* 1994;90:2236–40.
49. Smith FB, Lee AJ, Fowkes FG, Price JF, Rumley A, Lowe GD. Hemostatic factors as predictors of ischemic heart disease and stroke in the Edinburgh artery study. *Arterioscler Thromb Vasc Biol* 1997;17:3321–5.
50. Fowkes FG, Lowe GD, Housley E, Rattray A, Rumley A, Elton RA, MacGregor IR, Dawes J. Cross-linked fibrin degradation products, progression of peripheral arterial disease, and risk of coronary heart disease. *Lancet* 1993;342:84–6.

51. von Känel R, Kudielka BM, Hanebuth D, Preckel D, Fischer JE. Different contribution of interleukin-6 and cortisol activity to total plasma fibrin concentration and to acute mental stress-induced fibrin formation. *Clin Sci (Lond)* 2005;109:61–7.
52. Loucks EB, Sullivan LM, D’Agostino RB Sr, Larson MG, Berkman LF, Benjamin EJ. Social networks and inflammatory markers in the Framingham heart study. *J Biosoc Sci* 2006;38:835–42.
53. Lache B, Meyer T, Herrmann-Lingen C. Social support predicts hemodynamic recovery from mental stress in patients with implanted defibrillators. *J Psychosom Res* 2007;63:515–23.
54. Horsten M, Mittleman MA, Wamala SP, Schenck-Gustafsson K, Orth-Gomer K. Social relations and the metabolic syndrome in middle-aged Swedish women. *J Cardiovasc Risk* 1999;6:391–7.
55. von Känel R, Orth-Gomér K. Autonomic function and prothrombotic activity in women after an acute coronary event. *J Womens Health (Larchmt)* 2008;17:1331–7.
56. Cohen S. Keynote presentation at the Eight International Congress of Behavioral Medicine: the Pittsburgh common cold studies: psychosocial predictors of susceptibility to respiratory infectious illness. *Int J Behav Med* 2005;12:123–31.
57. Friedmann E, Thomas SA, Liu F, Morton PG, Chapa D, Gottlieb SS, Sudden Cardiac Death in Heart Failure Trial Investigators. Relationship of depression, anxiety, and social isolation to chronic heart failure outpatient mortality. *Am Heart J* 2006;152:940.e1–8.
58. Rodriguez-Artalejo F, Guallar-Castillon P, Herrera MC, Otero CM, Chiva MO, Ochoa CC, Banegas JR, Pascual CR. Social network as a predictor of hospital readmission and mortality among older patients with heart failure. *J Card Fail* 2006;12:621–7.
59. Hawkey LC, Masi CM, Berry JD, Cacioppo JT. Loneliness is a unique predictor of age-related differences in systolic blood pressure. *Psychol Aging* 2006;21:152–64.
60. Barefoot JC, Gronbaek M, Jensen G, Schnohr P, Prescott E. Social network diversity and risks of ischemic heart disease and total mortality: findings from the Copenhagen city heart study. *Am J Epidemiol* 2005; 161:960–7.