

Passerine migration strategies and body mass variation along geographic sectors across East Africa, the Middle East and the Arabian Peninsula

Elizabeth Yohannes · Herbert Biebach ·
Gerhard Nikolaus · David J. Pearson

Received: 15 November 2007 / Revised: 12 October 2008 / Accepted: 17 October 2008 / Published online: 19 November 2008
© The Author(s) 2008. This article is published with open access at Springerlink.com

Abstract Using the body mass estimates of 12 long-distance migrating Palearctic passerine species monitored at successive sites across the Eastern Africa flyway, we tested whether birds modulate their body mass according to specific seasonal demands across different geographic sectors. We compared body mass estimates across latitudinal distances and geographic sectors in Europe, the desert, Northeast Africa and East Africa. Our results show that, depending on the species and season considered, the average body mass increase or decrease is variable at and among different geographic sectors. By comparing the variation in body mass between different ecological sectors, we were able to show when and where migrants accumulate their migratory fuel reserves during migration.

Keywords Body mass · Eastern Africa · Migration strategy · Palearctic passerines

Introduction

Most organisms that live in seasonal environments undertake seasonal journeys of varying distances. World wide, millions of birds leave northern latitude breeding grounds every autumn to migrate to the south and undergo a return migration during the spring. Some species travel a little over 100 km, while others cover longer distances and can be on migration for about half a year (Alerstam 1990). To make these trips, birds need to cope with the large energy costs of long-distance flight, particularly when crossing ecological barriers, such as oceans, mountains and deserts.

The eco-physiological mechanisms adopted by long-distance migrating Palearctic birds for crossing ecological barriers have attracted the interest of ornithologists for more than 45 years (Moreau 1961, 1972). Different ecological theories have been proposed to describe the seasonal migrations and flights for crossing such barriers (particularly the Sahara Desert). To date, the authors of each study have argued—to varying degrees—that birds use flight strategies based on the incorporation of step-wise and/or intermittent flights. These include a non-stop flight, short/long intermittent flight, migration oriented along a given route/site, broader front, stopover at oases, nocturnal flights, using shade to avoid heat in the day, loop migration, among others (see, for example, Bairlein et al. 1983; Bairlein 1985, 1991; Biebach et al. 1986; Biebach 1990; Pearson et al. 1988; Schmaljohann et al. 2006).

Sustained flight over long-distance journeys and across ecological barriers requires substantial migratory fuel deposition and energy accumulation. In the Palearctic–African avian migration system, the majority of the passerine migrants which cross the Sahara Desert are known to deposit large fuel loads before crossing this barrier (see Fry et al. 1970; Finlayson 1981; Fransson et al. 2005, 2006).

Communicated by F. Bairlein.

E. Yohannes (✉)
Von-der-Tann Str. 7, 82346 Andechs, Germany
e-mail: yohannes@orn.mpg.de

H. Biebach
Max-Planck Institute for Ornithology, Von-der-Tann Str. 7,
82346 Andechs, Germany

G. Nikolaus
Padingbutteler Strich 36, 27632 Padingbutteler, Germany

D. J. Pearson
Lupin Close Reydon, Southwold, Suffolk IP18 6NW, UK

Hypothetical migration strategy models that are based on migratory fuel accumulation derived from studies along the Western Africa flyway (Zink 1973, 1982; Bairlein 1991; Schaub and Jenni 2000a, b; Fransson et al. 2001; Ottosson et al. 2005) indicate at least four different types of autumn strategies along a migration sector between northern Europe and North Africa:

1. birds which show a steady increase in body mass along the migration route (e.g. Garden Warblers *Sylvia borin* and Pied Flycatchers *Ficedula hypoleuca*);
2. birds which start to increase body mass towards the south, but well before reaching the Sahara (e.g. Sedge Warblers *Acrocephalus schoenobaenus*);
3. birds which do not show a steadily increase in body mass towards the south, but accumulate the required energy to fly over the Sahara Desert in North Africa just before crossing it (e.g. Reed Warblers *Acrocephalus scirpaceus* and Whitethroats *Sylvia communis*);
4. birds which do not steadily increase their body mass towards the south and do not accumulate excess energy that could be used to cross the Sahara Desert in North Africa but which regularly stop and fuel up in the Sahara (e.g. Spotted Flycatchers *Muscicapa striata*).

The difference in energy-accumulating strategies may largely depend on the availability of suitable stopover sites, spatio-temporal variation in food and species differences in food preferences (e.g. exclusively insectivores flycatchers vs. warblers that might also feed on fruits and berries). The strategies described above mainly support organization of the journey before and during the crossing of the Sahara Desert in autumn. However, our knowledge of the organization of fuel accumulation for migrating and traversing ecological barriers in spring is very limited.

Although many investigations of migratory Palearctic passerines along the Eastern African flyway have provided data on the body mass of birds at one particular site, very few (e.g. Bairlein 1997, 1998; Schaub and Jenni 2000a, b) have reported comparative results from two or more successive study sites. Indeed, most of the available studies were aimed at one single species and at a limited number of study sites. Furthermore, unlike the situation for the Western Palearctic migration route (e.g. Bairlein 1997, 1998), there has, to date, been no scientific network that has monitored birds along the Eastern Africa, the Middle Eastern and Arabian flyways to make a detailed comparative study of migration along this route in the autumn and spring. Thus, it is not well known when and where Palearctic migrants prepare for migration between different locations in the autumn and spring migration (e.g. Berthold 1996) along this route.

In this study, we investigated when and where Palearctic migrants prepare and accumulate their migration energy budget during the autumn and spring migrations. For

discussion purposes we have segmented the migration routes into four different sectors: Europe, Desert, Northeast Africa (NE Africa) and Eastern Africa. We used body mass measurements of 12 migratory passerine birds monitored at successive sites in Europe, Eastern Africa, the Middle East and Arabia. Our approaches were twofold: first, we compared the general and specific migratory strategies adopted by different species in different seasons. Second, by comparing the variation in strategies between different ecological sectors along the migration route, we tested whether birds modulate their energy-budget (body mass) according to specific seasonal and ecological demands en-route.

Unlike the western population, migrating bird species originating from the eastern populations have to cover a longer distance of inhospitable area. Based on known information and from a theoretical migration strategy, different types of strategies can be hypothesized.

During the autumn birds may start accumulating energy reserves in the north (e.g. Europe) and show a steady increase of body mass towards the south before crossing the desert. Birds cross the desert without refueling and hence have lower body mass upon reaching regions south of the Sahara. Alternatively, birds may use an intermittent ‘stop and refuel’ strategy and exhibit no steady increase or decrease in body mass during their migration across Europe and the desert. Species that do not apply a ‘stop and refuel’ strategy during migration across the desert may exhibit a steady decline of body mass. These birds accumulate larger energy stores upon reaching NE and Eastern Africa, prior to their southward migration to the wintering grounds.

In the spring, birds are expected to accumulate large energy reserves in NE Africa immediately before crossing the desert. To accomplish this, birds may accumulate energy stores at each stopover site before crossing the desert in Eastern and NE Africa. This would result in a steady increase in body mass as they move northward, while energy reserves would decline during the desert crossing. Birds could also accumulate substantial energy reserves in Eastern Africa but lose body mass during their northbound journey to NE Africa and the desert. When faced with a situation in which such a choice must be made, birds may tend to find high-quality stopover areas in NE Africa close to the desert or, alternatively, to apply the ‘stop and refuel’ strategy while crossing the desert.

Methods

Body mass data and species

Autumn and spring body mass values of migratory birds were obtained from both published and unpublished

resources from several ringing sites (Fig. 1) and migration studies in NE and Eastern Africa. These include data from the Sudan (Nikolaus and Pearson 1982; Nikolaus 1983), Ethiopia (Ash 1980, 1981; J. Ash, unpublished; B. Biebach, unpublished), Uganda (Pearson 1972, and unpublished; M. Fogden, unpublished; M. Wilson, unpublished), Kenya (Pearson and Backhurst 1976a, b, and unpublished; Britton and Britton 1977, and unpublished) and Tanzania (L. and N. Baker, unpublished; B. Biebach and U. Bauchinger, unpublished). We considered data gathered from Egypt (Moreau 1961; Watson 1971; Goodman and Meininger 1989; B. Biebach and U. Bauchinger, unpublished), Eritrea (Smith 1957) and Somalia (Archer and Godman 1961; Ash and Miskell 1998; J. Ash, unpublished). We also used published and unpublished data sources from the Middle East and the Arabian regions, including Israel (Safriel 1968; Safriel and Lavee 1988; Morgan and Shirihai 1997), Jordan (J. Ash and R. Spencer, unpublished; D. Pearson, unpublished), Kuwait (Cleere et al. 2000, and unpublished),

Oman and Saudi Arabia (G. Nikolaus, J. Ash, M. Evans and S. Newton, unpublished). Published migration information was also included in the analysis (e.g. Csörgö and Lövei 1995; Fransson 1995; Chernestov 1996; Lindström et al. 1996; Morgan and Shirihai 1997; Schmidt 1964; Bairlein 1997; Honza et al. 2000).

The selection of ‘candidate species’ for this study was based (but not exclusively dependent) on Pearson et al. (1988) and Pearson and Lack (1992). We tended to include species that differ in their general spatio-temporal migration pattern (Cramp and Perrins 1992, 1993). Our main focus was to include both long- and short-distance trans-Saharan migrants (that migrate and winter south and north of the equator, respectively) of the Eastern African flyway/migration route (taken here as the route through Sudan, Ethiopia, Eritrea, Egypt, Somalia, Kenya, Uganda and Tanzania) with different breeding origins and wintering locations. Broadly, we applied four criteria for the choice of ‘candidate species’ (modified after Pearson et al. 1988 and Pearson and Lack 1992):

1. species wintering entirely in NE Africa (e.g. Lesser Whitethroat).
2. species that mainly winter in NE Africa and are uncommon south of 5°S (e.g. Nightingale and Barred Warbler).
3. species that mainly winter along the equator and south of it (e.g. Willow Warbler, Garden Warbler, Spotted Flycatcher, Great Reed Warbler, Sedge Warbler and Whitethroat).
4. species wintering entirely south of the equator but with very few wintering north of 5°S (e.g. Thrush Nightingale, Marsh Warbler and Red-Backed Shrike).

The categorizations are not mutually exclusive, and it is evident that some species, such as the Great Reed Warbler, Sedge Warbler, and Whitethroat winter in a region that covers a wider section from the northern to the southern tropics. In addition, we selectively included species that are characterized by having a ‘two-stage’ southward migration (a rapid southward migration followed by an extended stopover in NE Africa before resuming migration to southern Africa; e.g. Marsh warbler) and/or that show a ‘loop migration’ (spring northward passage that follows a more easterly course than during the autumn passage; e.g. Red-backed Shrike) on their northward journey (Pearson et al. 1988).

We used body mass data of 12 species: the Great Reed Warbler *Acrocephalus arundinaceus*, Marsh Warbler *A. palustris*, Sedge Warbler, Red-backed Shrike *Lanius colurio*, Thrush Nightingale *Luscinia luscinia*, Nightingale *L. megarhynchos*, Spotted Flycatcher, Willow Warbler *Phylloscopus trochilus*, Garden Warbler, Whitethroat, Lesser Whitethroat *Sylvia curruca* and Barred Warbler *Sylvia*




Fig. 1 Geographic distribution of study sites and migration sectors. Triangle, hexagon and square symbols indicate study sites in the autumn, spring, and both autumn and spring, respectively. NE Africa Northeast Africa

nisoria collected from more than 100 sites located in more than 15 different countries.

Some ringing sites in East Africa were used as wintering grounds by some birds and, therefore, both wintering and migrating birds were present. The distinction between migratory and wintering birds was carried out according to the timing of migration described by Ash (1981), Pearson et al. (1988), Pearson and Lack (1992), Cramp and Perrins (1992, 1993) and Pearson (personal communication) (Table 1).

Since the body mass measurements considered in this study were gathered from several studies undertaken in different years by various investigators, the data were not collected in a wholly standardized way. Corrections for effects of possible sources of bias could not be carried out.

Data analysis

For the 12 study species, seasonal body mass data in the autumn ($n = 38,094$ birds) from 57 study sites and spring ($n = 24,592$ birds) from 54 study sites were calculated. Our approaches were twofold: first, we calculated the mean \pm standard error (SE) of body mass values at successive latitudinal geographic position of each study site.

Second, we made an extended comparison of the results by dividing the Eastern flyway latitudinal regions into four different sectors (Fig. 1):

1. European sector: study sites located north of 40°N;
2. Desert sector: localities in the Middle East and North Africa between 35°N and 15°N, including Kuwait, Jordan, Israel, Saudi Arabia, Oman, Egypt, northern and central Sudan and Eritrea;
3. NE Africa sector: regions along the Eastern African flyway located between 15°N and 4°N (central and southern Sudan, Ethiopia, Somalia, northern Kenya, northern and central Uganda);
4. Eastern Africa sector: localities below 4°N and 10°S (study sites in southern Kenya, southern Uganda and Tanzania).

We used nonparametric statistics (Spearman correlation with Bonferroni correction for a significance level following multiple comparisons) to test for an association between body mass and latitude for each species and season. Due to restricted body mass data from birds on migration during the spring migration in Europe, spring comparisons are made only for sectors in Africa and Arabia. For each species, variation in body mass along each

Table 1 Study locations in Eastern Africa and earliest arrival (autumn) and departure (spring) dates considered (as defined in Methods) for each species

Species	Study sites and migratory period				
	Ethiopia	Central Kenya	Uganda	Ngulia, Kenya	Tanzania
Marsh Warbler	<Nov 30	<Dec 21		<Dec 18	<Jan 30 >Apr 5
Sedge Warbler	<Jan 22 >Apr 15	<Dec 27 >Apr 5	<Jan 20 >Mar 17	<Dec 21	
Red-backed Shrike			<Nov 11	<Dec 7	
Thrush Nightingale	<Dec 5	>Mar 27 >Apr 20		<Dec 15	>Mar 11 <Jan 12 >Mar 14
Nightingale	<Nov 29 >Feb 27			<Dec 12	
Spotted Flycatcher	<Dec 5 >Apr 23				
Willow Warbler	<Dec 6 >Mar 25	<Dec 20 >Mar 3	<Dec 17 >Mar 5	<Dec 20	<Dec 25 >Mar 10
Garden Warbler	<Nov 25 >Apr 12	<Jan 5 >Mar 25	<Dec 20 >Mar 8	<Dec 30	<Jan 21 >Mar 10
Whitethroat	<Nov 27 >Mar 15	<Nov 25 >Mar 31		<Dec 18	<Dec 25 >Mar 10
Lesser Whitethroat	<Oct 10 >Mar 5				
Barred Warbler				<Dec 18	
	>Mar 12	>Mar 21			

The signs < and > indicate before and after, respectively, the dates given in the Table

migration sector was compared separately by applying the Mann–Whitney *U* test with a Bonferroni-adjusted significance level of 0.004.

Results

Plots of the mean body mass values in response to latitude are given in Fig. 2a, b. There was no significant association between latitude and body mass in autumn (for all species

investigated). Significant negative associations were observed for the Marsh Warbler ($r = -0.90, P = 0.003$) and Whitethroat ($r = -0.87, P = 0.003$) crossing the desert during the spring migration; the other ten species considered showed no significant association between latitude and body mass during the spring migration.

Tables 2, 3 show sector-wise mean \pm SE body mass values for each species. With the exception of the Sedge and Garden Warbler, body mass comparisons between pairs of geographic sectors showed no significance




Fig. 2 a Autumn, b spring. Mean body mass \pm standard error (SE)




Fig. 2 continued

differences (for all the other species considered in this study). For the Sedge Warbler, sector-wise autumn body mass comparisons revealed significantly higher body mass values in Europe compared to values in NE ($P = 0.002$) and Eastern Africa ($P < 0.001$). The Garden Warbler showed a significantly higher body mass in the Eastern African sector than in the NE African sector ($P < 0.001$).

Body mass values recorded from the desert were not significantly different than those calculated from study sites located in Europe, NE Africa or Eastern Africa in the autumn. Only the Willow Warbler showed a significant increase in body mass values (during spring migration) in the NE African sector compared to the Eastern Africa ($P = 0.003$) and Desert sectors ($P = 0.001$).




Fig. 2 continued

Discussion

Body mass variation among latitudes and migration sectors

With the exception of the Desert sector during the spring migration, body mass did not show a significant association with latitude during the autumn and spring migration. This implies that birds do not show a substantial increase or decrease in energy reserve per latitude in a given sector. Most species showed an increase in body mass along the

two African sectors but a decrease during their northward migration across the desert in the spring (Table 3). It has been reported that nearly all long-distance trans-Saharan migrant passerines landing in the Sahara Desert have both body masses well above the breeding level and sufficient energy reserves to cross the desert without refueling. Most of the birds checked at an oasis were not critically short of energy reserves, and only a few of them used the desert oases to regain fat (Bairlein et al. 1983; Bairlein 1985; Biebach 1985; Biebach et al. 1986; Biebach 1988; Bairlein 1992; Biebach 1995). This result suggests that energy




Fig. 2 continued

reserves are either deposited before the birds start crossing the Sahara or at stopover sites in the desert.

After crossing the desert, most species showed a substantial decrease in body mass in NE Africa compared to values registered in Europe. The Sedge Warbler, Red-backed Shrike, Thrush Nightingale, Nightingale, Willow Warbler, Garden Warbler, Whitethroat and Barred Warbler showed a marked decrease in body mass in this sector (Table 2). Across the Eastern African flyway, several bird

species perform step migration whereby they cross the Saharan or Arabian deserts without delay and arrive for extended stopover in NE Africa. Within this stopover period, many birds accomplish a partial or complete moult. Migratory passerines with such a stepwise autumn migration and moult include the Marsh Warbler, Whitethroat and Great Reed Warbler (Pearson 1973, 1975; Pearson and Backhurst 1976a; Curry-Lindahl 1981; Pearson 1982; Dowsett-Lemaire and Dowsett 1987; Pearson et al. 1988).

Table 2 Changes in mean body mass during the autumn migration for 12 species of passerine migrants at four geographic sectors

Species	Geographic sectors							
	Europe		Desert		NE Africa		East Africa	
	Mean ± SE	<i>n</i>	Mean ± SE	<i>n</i>	Mean ± SE	<i>n</i>	Mean ± SE	<i>n</i>
Great Reed Warbler	31.71 ± 1.04	1201	28.22 ± 1.14	398	29.98 ± 0.96	21	29.57 ± 1.31	32
Marsh Warbler	12.40 ± 0.34	351	11.85 ± 0.54	2699	10.93 ± 0.15	50	11.57 ± 0.22	5622
Sedge Warbler	12.25 ± 0.13	765	11.08 ± 0.68	635	10.93 ± 0.15	37	11.02 ± 0.16	58
Red-backed Shrike	31.64 ± 1.37	444	26.83 ± 1.38	1011	26.73 ± 0.28	14	28.70 ± 0.90	804
Thrush Nightingale	27.38 ± 0.30	358	22.33 ± 1.45	119	22.22 ± 0.51	167	22.27 ± 0.47	2707
Nightingale	25.72 ± 0.84	38	–	126	20.69 ± 0.48	544	22.19 ± 0.98	60
Spotted Flycatcher	15.29 ± 0.24	484	14.20 ± 4.54	774	14.61 ± 0.47	29	14.89 ± 0.49	653
Willow Warbler	9.04 ± 0.23	2758	8.90 ± 0.56	2022	8.17 ± 0.16	61	8.52 ± 0.18	1165
Garden Warbler	19.64 ± 0.65	1052	19.60 ± 0.90	1095	17.11 ± 0.32	125	19.04 ± 0.22	1140
Whitethroat	15.27 ± 0.35	478	14.36 ± 0.86	904	14.03 ± 0.12	246	14.64 ± 0.19	3697
Lesser Whitethroat	12.27 ± 0.46	87	11.98 ± 0.52	1941	11.07 ± 0.10	95	–	–
Barred Warbler	31.24 ± 1.54	109	23.79 ± 2.40	336	24.22 ± 0.46	23	24.03 ± 0.38	559

SE, Standard error

Table 3 Changes in mean body mass during the spring migration for 12 species of passerine migrants at three geographic sectors

Species	Geographic sectors					
	East Africa		NE Africa		Desert	
	Mean ± SE	<i>n</i>	Mean ± SE	<i>n</i>	Mean ± SE	<i>n</i>
Great Reed Warbler	32.10 ± 1.09	94	34.07 ± 1.08	71	29.27 ± 1.18	1094
Marsh Warbler	11.79 ± 0.23	64	14.47 ± 1.26	22	11.67 ± 0.45	792
Sedge Warbler	11.96 ± 0.23	789	12.92 ± 0.57	152	10.86 ± 0.22	3210
Red-backed Shrike	27.08 ± 0.70	108	30.66 ± 1.19	281	26.16 ± 1.66	277
Thrush Nightingale	23.21 ± 0.80	154	29.63 ± 0.83	24	23.52 ± 1.68	269
Nightingale	23.23 ± 0.92	20	23.52 ± 1.64	34	21.44 ± 1.08	332
Spotted Flycatcher	14.31 ± 0.52	11	18.18 ± 1.06	209	14.75 ± 0.51	702
Willow Warbler	8.63 ± 0.1	1979	10.33 ± 0.50	567	7.94 ± 0.22	3394
Garden Warbler	20.13 ± 0.23	482	18.91 ± 0.70	201	17.92 ± 1.07	1227
Whitethroat	14.98 ± 0.24	636	15.84 ± 1.02	125	14.82 ± 0.62	1670
Lesser Whitethroat	–	–	11.98 ± 0.48	77	11.48 ± 0.45	5309
Barred Warbler	24.91 ± 1.12	24	23.09 ± 0.33	57	24.82 ± 0.78	135

However, individuals of other species may also perform moult in these areas. In such cases, a moult-related decline in body mass might be exhibited.

Along the NE Africa migration sector, slight gains in body mass were estimated for the Garden Warbler compared to those observed in Europe, the desert and Eastern Africa (Table 2). As indicated earlier by Ash (1980, 1994), some bird species arrive in NE Africa with a moderate body mass and show a substantial increase in body mass before departure to more southerly located wintering grounds.

In the spring, most of the study species gained considerable mass at the northeasterly sites in Africa (from which they take off for trans-desert migration to the northern latitude) followed by a substantial mass loss during migration across the deserts (Table 3). Birds caught in Eastern Ethiopia, Somalia and the Farasan Islands of Saudi Arabia revealed pronounced body mass increases relative to birds caught at study sites located in Eastern Africa. For example, on average, the Marsh Warblers in Jijiga, 09°22'N 42°51'E eastern Ethiopia, were 4.22 g heavier than birds in Kifufu (03°09'S 39°12'E), Tanzania. Marsh

Warblers were not observed feeding at the stopover site in Jijiga (H. Biebach, personal communication and personal observation); therefore, the increase in body mass must have been achieved at feeding sites located south of this site.

Based on our analysis, the four strategies demonstrated earlier (e.g. by Schaub and Jenni 2000a) for crossing the desert by migratory birds of the Western African migration route (i.e. steady increase in body mass along the migration route, increase in body mass towards the south before reaching the Sahara or just before crossing the desert and regular stop-and-fuel strategies) are not directly applied by birds migrating along the Eastern African flyway/migration route (taken here as the route through Egypt, Sudan, Eritrea, Ethiopia, Somalia, Kenya, Uganda and Tanzania). We are aware that, unlike the data from Schaub and Jenni (2000a), our results are derived from a dataset (gathered from different sources) that do not account for fluctuations or unidirectional changes due to factors that affect body mass estimates. Hence, some of the estimates may have given a biased conclusion. Nevertheless, our results, which are based on a large number of birds and study sites, provide valuable information on the general trend of passerine migration strategies over a wider geographic range in both seasons.

Schaub and Jenni (2000a) note that they did not sample birds from the eastern population of Sedge Warbler, Whitethroats and Spotted Flycatcher. However, our results (which include data from the eastern population) indicate that while migrating along the Eastern African flyway, some species (e.g. Sedge and Garden Warbler) may have adopted a slightly different strategy than those reported earlier (e.g. Schaub and Jenni 2000a, b).

Similar to the results shown for change in body mass during the migration from Western Europe to Northern Africa, the Spotted Flycatcher did not show a significant change along the Eastern flyway. Elsewhere, observations and recoveries of Spotted Flycatchers have indicated that North Africa is used as a stopover site before crossing the Sahara (Moreau 1972; Bundy 1976; Sultana and Gauci 1982; Flint and Stewart 1983; Fransson 1986). In a study at an Egyptian oasis, only very few Spotted Flycatchers stayed several days and gained some mass before continuing migration to the south (Biebach 1985; Goodman and Meininger 1989), while all of the others had high masses and departed after a short stopover. Along the entire migration route considered in this study, the highest autumn mass for the Spotted flycatcher was reported from the stopover sites in the deserts. This implies that birds may intermittently stop and refuel while crossing the desert.

Similarly, the western population of Sedge Warbler already starts to accumulate a larger fuel store in Central Europe, long before arriving at the northern edge of the

Sahara (Bibby and Green 1981; Schaub and Jenni 2000a). Along the eastern route, this species does accumulate its energy store in Europe, but it still maintains a steady body mass towards the south before crossing the desert. Similarly, Whitethroats showed no significant increase in body mass in the east, while in the west, it accumulates the larger energy reserves in North Africa before its flight across the Sahara.

These differences could mainly be accounted for by the fact that the eastern populations have to cover a longer distance of inhospitable area (including the Arabian Desert and the Red Sea followed by the Sahara Desert). As such, the different landscape pattern may place constraints that require a different type of migration strategy. Thus, it could be speculated that different populations might have evolved different migration strategies of energy storage during migration. Using body mass measurement of Garden Warblers caught on southbound migration across SW Europe, both Bairlein (1991) and Schaub and Jenni (2000a) independently found a similar pattern of a relatively larger increase in body mass in eastern birds than in western birds (comparing 10° east and west) during the autumn migration from the breeding ground in Europe to North Africa. These researchers argue that the western population of Garden Warblers migrates through the Iberian peninsula and crosses a shorter stretch of the Mediterranean Sea, while the Eastern population is required to cross the eastern section of the Mediterranean Sea and then crosses the Sahara Desert (Biebach 1996); this latter route provides limited opportunity for refueling and, hence, larger energy reserves needs to be stored.

Several factors may affect the body mass measured at various study sites. These include body size, time and intensity of moult, time of day and year as well as inter-year variation, method and effort of capture, geographical location, food availability in the study site, errors and accuracy of measurements by various investigators, among many others (Clark 1979; Murphy 1996; Schaub and Jenni 2000a). In addition, different populations may be associated at different study sites and in different seasons. Similarly, other factors, such as sampling bias towards lower body mass values, the arrival and departure of the birds caught at each site, the stopover duration, fuel deposition rate of the birds at each site, variation between different years and weather conditions, contribute to the variation in body mass.

For some of the species, body mass values differed between geographic sectors, indicating that birds possibly modulate their energy budget depending on their current location (geographic sector) en-route. This is in agreement with a recent appraisal and study of spatio-temporal precision in bird migration. For instance, in an experimental study, Fransson et al. (2001) and Kullberg et al. (2003,

2007) illustrated that birds may use geomagnetic cues as external indicators for successful migration, and they explain that these cues (simultaneously operating with bird's orientation behavior and endogenous rhythm) provide information about the geographical position when such information is required, for instance, for a fueling decision. Given this, depending on the species and the route taken, a certain relationship would be expected between body mass and latitude of a given geographic area.

Palaearctic birds that migrate through eastern and northeastern Africa are likely to belong to populations from different breeding grounds. Some passerines cross the Sahara to reach breeding grounds in the eastern Mediterranean while others presumably migrate over a broader front to move into Southwest Asia through the Arabian regions (Moreau 1972). These two subpopulations may differ in terms of the timing of migration and fuel deposition (and hence extent of fattening), which in turn may be related to the physical and geographic differences between the migration routes.

Some birds bound for the breeding areas in the north-northeast may start to increase body mass as far south as the equator. For example, Pearson (1971) has shown that in the spring, the heaviest Sedge and Marsh Warblers encountered in Uganda, Lake Victoria had sufficient fat reserves for migrating directly to the Middle East. However, a considerable number of birds caught in late spring in Uganda had moderate body masses. These birds may keep their reserves by frequently feeding, while moving either to more northerly sites or to further staging areas close to the desert. The extent and rate of fattening may also vary between sites (Pearson 1971, 1978).

Many locations in Eastern and NE Africa serve as important fattening areas for Palaearctic migratory birds in both south- and northbound migration seasons. In the autumn, birds show an increase in body mass in NE Africa before migrating to more southerly wintering grounds. In the spring, birds gain considerable mass in NE Africa and immediately to the south of the Arabian/Sahara Desert before takeoff for trans-desert migration. Thus, the quality of these stopover sites may very well exert a significant influence over the success of the journey as well as the reproductive fitness of the birds in their breeding grounds.

Conclusion

Our results are based on body mass estimates of 12 Palaearctic migratory passerines monitored at successive sites in Europe, Eastern Africa, the Middle East and Arabia. We show that, in general, passerine migrants adopt different patterns of accumulating energy reserves for crossing ecological barriers and migration. These patterns vary depending on the geographic sector and seasons (autumn

and spring). Moreover, different populations or subpopulations may follow specific strategies that particularly fulfill the eco-physiological demands of each geographic sectors en-route.

Zusammenfassung

Zugstrategien und Variation der Körpermasse von Singvögeln entlang geographischer Sektoren in Ostafrika, im Mittleren Osten und in Arabien

Wir überprüften, ob 12 langstreckenziehende Singvogelarten ihre Körpermasse entsprechend bestimmten saisonalen Erfordernissen längs ihren Zugrouten ändern. Dazu verwendeten wir Körpermasse-Daten, die an Orten auf verschiedenen Breitengraden und in verschiedenen Sektoren entlang des ostafrikanischen Zugweges in Europa, der Sahara und in Nordost- und Ostafrika gesammelt worden waren. Die durchschnittlichen Zu- und Abnahmen der Körpermasse variierten sowohl innerhalb einzelner Sektoren als auch zwischen verschiedenen Sektoren je nach Vogelart und Saison. Durch den Vergleich der Variation der Körpermassen in verschiedenen ökologischen Sektoren können wir zeigen, wann und wo Langstreckenzieher während des Zuges ihre Fettreserven auffüllen.

Acknowledgments This study comprises an analysis of data collected from a number of investigations. We thank all the authorities, investigators, ringers and numerous field assistants involved in the projects. Special thanks are extended to John Ash for kindly allowing us to use his dataset. Our sincere thanks are also expressed to Prof. Eberhard Gwinner, the late director of the Max-Planck Research Centre for Ornithology and Prof. Ellen Thaler who supported the study in several ways. The Austrian Academic Exchange Service, Ministry of Foreign Affairs and Max-Planck Institute for Ornithology provided financial supports. Comments and suggestions by Graham Grieve, Richard E. Johnson, Pavel Ktitorov, Holger Schielzeth and three anonymous reviewers helped to improve the manuscript.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Alerstam T (1990) Bird migration. Cambridge University Press, Cambridge
- Archer G, Godman EM (1961) The birds of British Somaliland and the Gulf of Aden, vol 3–4. Oliver & Boyd, London
- Ash JS (1980) Migrational status of Palaearctic birds in Ethiopia. In: Proc 4th Pan-African Ornithol Congress. Seychelles, pp 199–208
- Ash JS (1981) Bird-ringing results and ringed bird recoveries in Ethiopia. *Scopus* 5:85–101

- Ash JS (1994) Weights of migrant Palearctic *Sylvia* warblers in Ethiopia. *Scopus* 17:119–123
- Ash JS, Miskell JE (1998) *Birds of Somalia*. Pica Press, Robertsbridge
- Bairlein F (1985) Body weights and fat deposition of Palearctic passerine migrants in the central Sahara. *Oecologia* 66:144–146. doi:10.1007/BF00378566
- Bairlein F (1991) Body mass of Garden Warbler (*Sylvia borin*) on migration: a review of field data. *Vogelwarte* 36:48–61
- Bairlein F (1992) Recent prospects on trans-Saharan migration of songbirds. *Ibis* 134:41–46
- Bairlein F (1997) Spatio-temporal course, ecology and energetics of western Palearctic–African songbird migration. Summary report 1994–1996. Institut für Vogelforschung, Wilhelmshaven
- Bairlein F (1998) The European–African songbird migration network: new challenges for large-scale study of bird migration. *Biol Cons Fauna* 102:13–27
- Bairlein F, Beck P, Feiler W, Querner U (1983) Autumn weights of some Palearctic passerine migrants in the Sahara. *Ibis* 125:404–407. doi:10.1111/j.1474-919X.1983.tb03131.x
- Berthold P (1996) Control of bird migration. Chapman & Hall, London
- Bibby CJ, Green RE (1981) Autumn migration strategies of Reed and Sedge Warblers. *Ornis Scand* 12:1–12. doi:10.2307/3675898
- Biebach H (1985) Sahara stopover in migratory Flycatchers: fat and food affect the time program. *Experientia* 41:695–697. doi:10.1007/BF02007727
- Biebach H (1988) Ecophysiology of resting Willow Warblers (*Phylloscopus trochilus*) crossing the Sahara. In: Proc 19th Int Ornithol Congress. Ottawa, pp 2162–2168
- Biebach H (1990) Strategies of trans-Saharan migrants. In: Gwinner E (ed) *Bird migration: physiology and ecophysiology*. Springer, Berlin Heidelberg New York, pp 352–367
- Biebach H (1995) Stopover of migrants flying across the Mediterranean Sea and the Sahara. *Isr J Zool* 41:387–392
- Biebach H (1996) Energetics of winter and migratory fattening. In: Carey C (ed) *Avian energetics and nutritional ecology*. Chapman & Hall, New York, pp 280–323
- Biebach H, Friedrich W, Heine G (1986) Interaction of bodymass, fat, foraging and stopover period in trans-Saharan migrating passerine birds. *Oecologia* 69:370–379. doi:10.1007/BF00377059
- Britton PL, Britton BA (1977) An April fall of Palearctic migrants at Ngulia. *Scopus* 1:109–111
- Bundy G (1976) The birds of Libya. An annotated check-list. B. O. U. Check-list No. 1. British Ornithologists' Union, London
- Chernestov N (1996) Preliminary hypotheses on migration of the Sedge Warbler (*Acrocephalus schoenobaenus*) in the Eastern Baltic. *Vogelwarte* 38(4):201–210
- Clark GA Jr (1979) Body weights of birds: a review. *Condor* 81:193–202. doi:10.2307/1367288
- Cleere N, Kelly D, Pilcher CWT (2000) Result from late autumn ringing project in Kuwait, 1995. *Ring. Migr* 20:186–190
- Cramp S, Perrins CM (1992) *Handbook of the birds of Europe, the Middle East and North Africa: the birds of the western Palearctic. Warblers*, vol VI. Oxford University Press, Oxford
- Cramp S, Perrins CM (1993) *Handbook of the birds of Europe the Middle East and North Africa: the birds of the western Palearctic. Flycatchers to shrikes*, vol 7. Oxford University Press, Oxford
- Csörgö T, Lövei GL (1995) Autumn migration and recurrence of the Thrush Nightingale *Luscinia luscinia* at a stopover site in central Hungary. *Ardeola* 42(1):57–68
- Curry-Lindahl K (1981) *Bird migration in Africa*, vol 1. Academic Press, London
- Dowsett-Lemaire F, Dowsett RJ (1987) European Reed and Marsh Warblers in Africa: migration patterns, moult and habitat. *Ostrich* 58:65–85
- Finlayson JC (1981) Seasonal distribution, weights and fat of passerine migrants at Gibraltar. *Ibis* 123:88–95. doi:10.1111/j.1474-919X.1981.tb00176.x
- Flint PR, Stewart PF (1983) The birds of Cyprus. B. O. U. Check-list no. 6. British Ornithologists' Union, London
- Fransson T (1986) The migration and wintering area of Nordic spotted flycatcher, *Muscicapa striata* (English summary). *Var Faglarld* 45:5–18
- Fransson T (1995) Timing and speed of migration in North and West European populations of *Sylvia* warblers. *J Avian Biol* 26:39–48. doi:10.2307/3677211
- Fransson T, Jakobsson S, Johannsson P, Kullberg C, Lind J, Vallin A (2001) Magnetic cues trigger extensive refuelling. *Nature* 414:35–36. doi:10.1038/35102115
- Fransson T, Jakobsson S, Kullberg C (2005) Nonrandom distribution of ring recoveries from trans-Saharan migrants indicates species-specific stopover sites. *J Avian Biol* 36:6–11. doi:10.1111/j.0908-8857.2005.03471.x
- Fransson T, Jakobsson S, Kullberg C, Mellroth R, Pettersson T (2006) Fuelling in front of the Sahara desert in autumn—an overview of Swedish field studies of migratory birds in the eastern Mediterranean. *Ornis Svec* 16:74–83
- Fry CH, Ash JS, Ferguson-Lees J (1970) Spring weights of some Palearctic migrants at Lake Chad. *Ibis* 112:58–82. doi:10.1111/j.1474-919X.1970.tb00076.x
- Goodman SM, Meininger PL (1989) *The birds of Egypt*. Oxford University Press, Oxford
- Honza M, Literak I, Pavelka J, Formanek J (2000) Postbreeding occurrence of the Marsh Warbler *Acrocephalus palustris* in reed bed areas in the Czech Republic and its migration to Africa. *Ökol Vögel* 22:119–129
- Kullberg C, Lind J, Fransson T, Jakobsson S, Vallin A (2003) Magnetic cues and time of season affect fuel deposition in migratory thrush nightingales (*Luscinia luscinia*). *Proc R Soc Lond Ser B* 270:373–378. doi:10.1098/rspb.2002.2273
- Kullberg C, Henshaw I, Jakobsson S, Johannsson P, Fransson T (2007) Fuelling decisions in migratory birds: geomagnetic cues override the seasonal effect. *Proc R Soc Lond Ser B* 274:2145–2151
- Lindström Å, Hedenström A, Pettersson J (1996) The autumn migration of Willow Warbler *Phylloscopus trochilus* in Sweden: results from a nation-wide co-operative project. *Ornis Svec* 6:145–172
- Moreau RE (1961) Problems of Mediterranean–Saharan migration. *Ibis* 103:373–427. doi:10.1111/j.1474-919X.1961.tb02454.x
- Moreau RE (1972) *The Palearctic–African bird migration systems*. Academic Press, London
- Morgan JH, Shirihai H (1997) Passerines and Passerine migration in Eilat. *Int Birdw Cent Eilat Tech Publ* 6(1):1–50
- Murphy ME (1996) Energetics and nutrition of moult. In: Carey C (ed) *Avian energetics and nutritional ecology*. Chapman & Hall, New York, pp 158–198
- Nikolaus G (1983) An important passerine ringing site near the Sudan Red Sea coast. *Scopus* 7:15–18
- Nikolaus G, Pearson DJ (1982) Autumn passage of Marsh Warblers *Acrocephalus palustris* and Sprossers *Luscinia luscinia* on the Sudan Red Sea coast. *Scopus* 6:17–19
- Ottosson U, Waldenström J, Hjort C, McGregor R (2005) Garden Warbler *Sylvia borin* migration in sub-Saharan West Africa: phenology and body mass changes. *Ibis* 147:750–757. doi:10.1111/j.1474-919X.2005.00460.x
- Pearson DJ (1971) Weights of some Palearctic migrants in southern Uganda. *Ibis* 113:173–184. doi:10.1111/j.1474-919X.1971.tb05142.x
- Pearson DJ (1972) The wintering and migration of some Palearctic passerines at Kampala, Southern Uganda. *Ibis* 114:43–60. doi:10.1111/j.1474-919X.1972.tb02588.x

- Pearson DJ (1973) Moults of some Palearctic warblers wintering in Uganda. *Bird Study* 20:24–36
- Pearson DJ (1975) The timing of complete moult in the Great Reed Warbler *Acrocephalus arundinaceus*. *Ibis* 117:506–509. doi: [10.1111/j.1474-919X.1975.tb04244.x](https://doi.org/10.1111/j.1474-919X.1975.tb04244.x)
- Pearson DJ (1978) The genus *Sylvia* in Kenya and Uganda. *Scopus* 2:63–71
- Pearson DJ (1982) The migration and wintering of Palearctic *Acrocephalus* warblers in Kenya and Uganda. *Scopus* 6:49–59
- Pearson DJ, Backhurst GC (1976a) The Southward migration of Palearctic birds over Ngulia, Kenya. *Ibis* 118:78–105. doi: [10.1111/j.1474-919X.1976.tb02012.x](https://doi.org/10.1111/j.1474-919X.1976.tb02012.x)
- Pearson DJ, Backhurst GC (1976b) Palearctic passerine migration at Kariobangi, Nairobi. *EANHS Bull*: 23–28
- Pearson DJ, Lack P (1992) Migration patterns and habitat use by passerine and near-passerine migrant birds in eastern Africa. *Ibis* 134:89–98
- Pearson DJ, Nikolaus G, Ash JS (1988) The Southward migration of Palearctic passerines through northeast and east tropical Africa: a review. *Proc 4th Pan-African Ornithol Congress. Seychelles*, pp 243–262
- Safriel UN (1968) Bird migration in Eilat, Israel. *Ibis* 110:283–320. doi: [10.1111/j.1474-919X.1968.tb00039.x](https://doi.org/10.1111/j.1474-919X.1968.tb00039.x)
- Safriel UN, Lavee D (1988) Weight changes of cross-desert migrants at an oasis—do energetic considerations along determine the length of stopover? *Oecologia* 76:611–619
- Schaub M, Jenni L (2000a) Body mass of six long-distance migrant passerine species along the autumn migration route. *J Ornithol* 141:441–460
- Schaub M, Jenni L (2000b) Fuel deposition of three passerine bird species along the migration route. *Oecologia* 122:306–317. doi: [10.1007/s004420050036](https://doi.org/10.1007/s004420050036)
- Schmaljohann H, Liechti F, Bruderer B (2006) Songbird migration across the Sahara: the non-stop hypothesis rejected! *Proc R Soc Lond B Biol Sci* 274:735–739. doi: [10.1098/rspb.2006.0011](https://doi.org/10.1098/rspb.2006.0011)
- Schmidt E (1964) Untersuchungen an einigen Holunder fressenden Singvögeln in Ungarn. *Zool Abh Mus Tierkde Dresden* 27:11–28
- Smith KD (1957) An annotated Check-list of the birds of Eritrea. *Ibis* 120:480–497
- Sultana J, Gauci C (1982) A new guide to the birds of Malta. Malta Ornithological Society, Valletta
- Watson GE (1971) A serological and ectoparasite survey of migratory birds in northeast Africa. Smithsonian Institution, Washington D.C.
- Zink G (1973) Der Zug europäischer Singvögel—ein Atlas der Wiederfunde beringter Vögel. Lief. 1. Vogelzug, Möggingen
- Zink G (1982) Der Zug europäischer Singvögel—ein Atlas der Wiederfunde beringter Vögel. Lief. 3. Vogelzug, Möggingen