

Amygdala–Hippocampal Connectivity Changes During Acute Psychosocial Stress: Joint Effect of Early Life Stress and Oxytocin

Yan Fan^{*,1,2,3}, Karin Pestke¹, Melanie Feeser^{1,2,3}, Sabine Aust^{1,2,3}, Jens C Pruessner^{4,5}, Heinz Böker⁶, Malek Bajbouj^{1,2,3,7} and Simone Grimm^{1,2,3,6,7}

¹Cluster of Excellence 'Languages of Emotion', Freie Universität Berlin, Berlin, Germany; ²Department of Psychiatry, Campus Benjamin Franklin, Charité Berlin, Berlin, Germany; ³Dahlem Institute for Neuroimaging of Emotion, Freie Universität Berlin, Berlin, Germany; ⁴Douglas Mental Health Institute, Department of Psychiatry, McGill University, Montreal, Canada; ⁵Montreal Neurological Institute, McGill University, Montreal, Canada; ⁶Department of Psychiatry, Psychotherapy and Psychosomatics, Hospital of Psychiatry, University of Zurich, Zurich, Switzerland

Previous evidence shows that acute stress changes both amygdala activity and its connectivity with a distributed brain network. Early life stress (ELS), especially emotional abuse (EA), is associated with altered reactivity to psychosocial stress in adulthood and moderates or even reverses the stress-attenuating effect of oxytocin (OXT). The neural underpinnings of the interaction between ELS and OXT remain unclear, though. Therefore, we here investigate the joint effect of ELS and OXT on transient changes in amygdala-centered functional connectivity induced by acute psychosocial stress, using a double-blind, randomized, placebo-controlled, within-subject crossover design. Psychophysiological interaction analysis in the placebo session revealed stress-induced increases in functional connectivity between amygdala and medial prefrontal cortex, posterior cingulate cortex, putamen, caudate and thalamus. Regression analysis showed that EA was positively associated with stress-induced changes in connectivity between amygdala and hippocampus. Moreover, hierarchical linear regression showed that this positive association between EA and stress-induced amygdala–hippocampal connectivity was moderated after the administration of intranasal OXT. Amygdala–hippocampal connectivity in the OXT session correlated negatively with cortisol stress responses. Our findings suggest that altered amygdala–hippocampal functional connectivity during psychosocial stress may have a crucial role in the altered sensitivity to OXT effects in individuals who have experienced EA in their childhood.

INTRODUCTION

Appropriate neural and physiological responses to transient shifts of homeostasis during stress are crucial for our survival and well-being (McEwen, 2003). The deterioration of homeostasis renders the individual vulnerable to the cumulative burden of stress. Cumulative stress early in life comprises a spectrum of adverse social experiences that occur before puberty, including sexual, physical, and emotional forms of abuse, as well as physical and emotional neglect (Pechtel and Pizzagalli, 2011). Ample evidence supports that early life stress (ELS) has enduring impact on the neuroendocrine system and leads to dysregulated cortisol responses to psychosocial stress in adulthood (Strüber *et al*,

2014). Such ELS-associated changes in stress reactivity have been associated with the development of depressive and anxiety disorders later in life (McEwen, 2003). However, the neural mechanisms underlying the impact of ELS on stress reactivity have not yet been elucidated.

Stress reactivity is regulated in the central nervous system through the coordinated circuitry coupling among limbic and prefrontal brain regions, such as the amygdala, hippocampus, and medial prefrontal cortex (mPFC) (Herman *et al*, 2005). As a key locus in the limbic–prefrontal circuit, the amygdala is reciprocally interconnected with regions involved in autonomic–neuroendocrine control, memory, and salience processing (Freese and Amaral, 2009). Amygdala activity and connectivity signal the emotional significance of the perceived stimuli and prepare our brain and body for suitable responses (LeDoux, 2000). Changes in functional connectivity (FC) between amygdala and other limbic–prefrontal regions have been demonstrated during and after stress induction (Hermans *et al*, 2011; Vaisvaser *et al*, 2013; Veer *et al*, 2011). Particularly, enhanced amygdala–hippocampal FC has been found both in the immediate responses to and in the prolonged aftermath of

*Correspondence: Y Fan, Klinik für Psychiatrie und Psychotherapie, Charité—Universitätsmedizin Berlin, Campus Benjamin Franklin, Hindenburgdamm 30, Berlin 12203, Germany, Tel: +49 (0)30 450 517778, Fax: +49 (0)30 450 517942, E-mail: yan.fan@fu-berlin.de or yan.fan@charite.de

⁷These authors contributed equally to this work.

stress (Vaisvaser *et al*, 2013). The hippocampus inhibits the hypothalamic–pituitary–adrenal (HPA) axis and receives negative feedback from its hormonal end product cortisol (Herman *et al*, 2005). Amygdala–hippocampal FC is inversely associated with cortisol levels (Vaisvaser *et al*, 2013) and predicts the capacity of the HPA axis to restore homeostasis after perturbation (Kiem *et al*, 2013). These previous findings demonstrate that amygdala–hippocampal FC has a critical role in responses to transient stress, especially for the restoration and maintenance of homeostasis via neuroendocrine control. It remains unclear though whether not only transient but also sustained/cumulative stress early in life is associated with altered amygdala–hippocampal FC.

ELS is related to changes in gray and white matter in stress regulation regions, such as amygdala, hippocampus, and mPFC. ELS-associated deficits in structural connectivity among these regions are also evident, supporting network abnormalities in the limbic–prefrontal circuit (Hart and Rubia, 2012). Recent studies have found ELS-associated decreases in resting-state FC (rs-FC) within the limbic–prefrontal circuit (Burghy *et al*, 2012; Fan *et al*, 2014; Herringa *et al*, 2013). Although rs-FC is postulated to reflect trait-like individual differences in circuitry integrity, task-evoked FC changes (eg, measured by psychophysiological interactions (PPIs)) are supposed to reflect transient changes in brain network coupling in response to specific task demands. Gee *et al* (2013) reported changes in amygdala–mPFC FC during emotion perception in adolescents who have experienced ELS. In line with the recent discovery of stress-dependent epigenetic changes in the hippocampus (Klengel *et al*, 2013), Holz *et al* (2015) investigated subjects with a genetic variant that makes them more sensitive to stress-dependent epigenetic modifications in the hippocampus and reported enhanced amygdala–hippocampus FC during emotion perception in these individuals. These findings further support that task-evoked FC changes in the limbic–prefrontal circuit are also influenced by ELS. For this reason, stress-induced changes in amygdala-centered FC are of primary importance for understanding the impact of ELS on stress reactivity. However, so far no study has investigated the influence of ELS on the transient FC shift within the limbic–prefrontal circuit during acute psychosocial stress.

Furthermore, ELS has long-lasting effects on the development of the oxytocinergic system, in which the secretion of oxytocin (OXT) inhibits both basal and stress-induced HPA axis activity (Neumann *et al*, 2000). Intranasal administration of OXT attenuates cortisol stress responses. However, in subjects who have experienced ELS, this stress-attenuating effect of OXT is diminished or even reversed (Grimm *et al*, 2014). The neural substrates involved in the altered sensitivity to OXT after ELS remain unclear. ELS is associated with decreased cerebrospinal fluid (CSF) OXT level in adulthood (Heim *et al*, 2009). Rodent models show that ELS reduces the expression of OXT receptors in the amygdala (Hill *et al*, 2014). Recent studies on fear and salience arousal show that OXT reduces amygdala activation and alters amygdala-centered FC to brain regions involved in autonomic–neuroendocrine control (Riem *et al*, 2012). In our recent study, contrary to the stress-attenuating effect in control subjects, OXT enhanced cortisol stress responses and stress-induced deactivation in limbic areas in subjects who experienced ELS (Grimm *et al*, 2014). It remains unclear

whether stress-induced changes in amygdala-centered FC are also modulated by the joint effect of OXT and ELS.

The primary aim of the present study was to investigate the effect of ELS on stress-induced changes in amygdala-centered FC. Task-evoked FC changes between amygdala and other limbic–prefrontal regions have been demonstrated during transient acute stress (Hermans *et al*, 2011; Vaisvaser *et al*, 2013; Veer *et al*, 2011). It has also been shown that ELS impacts structural connectivity and trait-like rs-FC within the limbic–prefrontal circuit (Burghy *et al*, 2012; Fan *et al*, 2014; Hart and Rubia, 2012; Herringa *et al*, 2013). Here we hypothesize that ELS influences stress-induced FC changes from amygdala to mPFC and hippocampus. Our second aim was to explore the interaction between ELS and OXT on stress-induced amygdala FC shift. Grimm *et al* (2014) found that the hippocampus is involved in the interaction effects of ELS and OXT on stress reactivity. We therefore hypothesize a joint effect of EA and OXT on stress-induced FC changes between amygdala and hippocampus.

MATERIALS AND METHODS

Assessment of ELS

ELS was measured using the Childhood Trauma Questionnaire (CTQ; Bernstein and Fink, 1998). The CTQ is a 28-item self-report questionnaire that retrospectively assesses five types of adverse childhood experiences: emotional neglect (EN), emotional abuse (EA), physical neglect (PN), physical abuse (PA), and sexual abuse (SA). Scores range from 5 to 25 for each subscale, with high scores indicating strong exposure to the corresponding type of maltreatment.

It has been found that different subtypes of ELS have differential impact on stress reactivity and stress-related affective disorders (Carpenter *et al*, 2009; Strüber *et al*, 2014). For example, unlike other subtypes of ELS, EA is a particularly strong predictor for blunt HPA reactivity, dampened cortisol stress responses, and internalizing psychopathology, such as major depression and social anxiety (Carpenter *et al*, 2009; Grimm *et al*, 2014; Keyes *et al*, 2012). Strong effects of EA have also been found on trait-like rs-FC within the limbic–prefrontal circuit, as well as on OXT concentrations in the CSF (Fan *et al*, 2014; Heim *et al*, 2009). For this reason, we focused on EA as a main predictor in the multiple regression analyses.

Subjects

Thirty-two healthy young male participants (aged 21–37 years, mean age 28.3 ± 4.7) were recruited from a preexisting community-dwelling sample ($N = 541$; Fuge *et al*, 2014). The short version of the Structured Clinical Interview for Diagnostic and Statistical Manual of Mental Disorders (Wittchen *et al*, 1997) was used to screen for history of psychiatric or neurological disease. Verbal intelligence was assessed using a German word recognition test (Lehrl *et al*, 1995; Schmidt and Metzler, 1992), which is functionally equivalent to the widely used National Adult Reading Test (Nelson and O'Connell, 1978). Inclusion criteria were age 18–40 years, absence of present and past diagnosis of psychiatric or neurologic disease, absence of major or unstable general medical conditions, IQ score (verbal intelligence)

> 90, and ability to participate in MRI study procedures. This study belongs to a project probing OXT effects on resting-state network (Fan *et al*, 2014) and stress reactivity (Grimm *et al*, 2014). The study was conducted in accordance with the latest version of the Declaration of Helsinki and approved by the Institutional Review Board of the Germany Psychological Society. All subjects gave written informed consent before screening and were reimbursed for participation.

Study Design

In a double-blind, placebo-controlled, within-subject cross-over design, participants received either OXT (Syntocinon Spray; Novartis, Basel, Switzerland) or a placebo (PLA; sodium chloride solution) intranasally in two separate experimental sessions. The sequence of OXT and PLA sessions was counter-balanced across subjects. We adopted a standardized protocol used in previous studies (Gamer *et al*, 2010; Labuschagne *et al*, 2010) where participants self-administered a single intranasal dose of 24 IU OXT/PLA (three puffs per nostril, each puff with 4 IU, or 6.82 mg) under the supervision of the experimenter 45 min before each fMRI scanning session. Participants abstained from alcohol during the 24 h and from exercise, caffeine, and food for 1 h before the experiment.

Induction of Psychosocial Stress

In each scanning session, psychosocial stress was induced using the Montreal Imaging Stress Task (MIST; Pruessner *et al*, 2008), in which the subjects completed a series of challenging mental arithmetic tasks. The MIST utilizes a block design and consists of three runs (each lasts for 7 min). Within each run, three conditions (ie, 0.5 min rest, 1 min control, and 2 min stress) were presented in this order and repeated once. In contrast to the control condition, the difficulty and time limit of the mental arithmetic tasks were manipulated in the stress condition, and the subjects were given negative feedbacks. Psychosocial stress was hence induced via elements of uncontrollability and social evaluative threat (for full description of the MIST, see Supplementary Methods, as well as Grimm *et al*, 2014).

Psychological and Physiological Measures

Before scanning started, the subjects completed the German versions of the Spielberger State-Trait Anxiety Inventory (STAI; Spielberger and Sydeman, 1994), the NEO five-factor inventory (Costa and McCrae, 1985), and the Rosenberg Self-Esteem scale (Rosenberg, 1965). To validate the stress induction, feelings of negativity and uncontrollability were assessed immediately before and after the fMRI scanning sessions via the state part of the STAI (Spielberger and Sydeman, 1994). Saliva samples were collected with the Salivette sampling device (Sarstedt) before (ie, 45 min after OXT/PLA administration) and immediately after the MIST task (for full description, see Grimm *et al*, 2014). Saliva-derived cortisol was analyzed using a time-resolved fluorescence immunoassay (Dressendorfer *et al*, 1992). The subjective report of state anxiety and salivary cortisol levels before and after stress induction were compared by paired-

sample *T*-tests. Cortisol stress responses were calculated as the difference between salivary cortisol levels before and after psychosocial stress.

MRI Acquisition

Structural and functional MRI data were acquired on a Siemens Trio 3T scanner using a 12-channel radio-frequency (RF) head coil. T1-weighted structural images were acquired with the following parameters: 176 sagittal slices covering the whole brain, repetition time (TR) = 1900 ms, echo time (TE) = 2.52 ms, flip angle = 9°, 256 × 256 matrix, voxel size 1 × 1 × 1 mm³. In the OXT and PLA sessions, respectively, 3 runs of 220 volumes of T2*-weighted echo-planar images were acquired with the following parameters: 37 axial slices of 3 mm thickness covering the whole brain, repetition time (TR) = 2000 ms, echo time (TE) = 30 ms, flip angle = 70°, 64 × 64 matrix, field of view = 192 × 192 mm², in-plane resolution = 3 × 3 mm².

fMRI Preprocessing

Functional images were preprocessed using MATLAB 2012 (The Mathworks, Natick, MA, USA) and SPM8 (Statistical parametric mapping software, SPM; Wellcome Department of Imaging Neuroscience, London, UK; <http://www.fil.ion.ucl.ac.uk>). The images were corrected for the acquisition time differences between slices, realigned to the first volume to correct for head motion between volumes, then normalized into standard stereotactic space (template provided by the Montreal Neurological Institute) using the normalization parameters estimated by T1 structural image unified segmentation, and finally smoothed with a Gaussian kernel of 8 mm full-width-half-maximum.

FC Analysis: PPI

Whole-brain-wise PPI analyses (Friston *et al*, 1997) were performed to examine how amygdala-centered FC pattern is modulated by acute psychosocial stress. The PPI analysis is a specific form of moderated multiple regression model. The predictors of this regression model include the time course of the experiment (psychological factor), the time course of a seed region (physiological factor), an interaction term of both, and covariates. Significant interaction effects between the psychological and physiological factors reveal how activity in a brain region covariates with activity in the seed region in response to the experimental manipulation.

We used the standard implementation of PPI in SPM8 and the generalized PPI toolbox (gPPI; <http://www.nitrc.org/projects/gppi>) introduced by McLaren *et al* (2012), which allows including more than one experimental term in the multiple regression model. In this manner, the psychological factors, ie, three experimental conditions (rest, control, stress) of the MIST, are represented by three separate regressors in the PPI model. Generalized PPI hence measures how FC strength changes for each task condition relative to the implicit baseline (eg, fixation) and is shown to be more sensitive and accurate for assessing pair-wise connectivity differences between conditions (eg, stress > control) when there are more than two conditions in the task.

Two seed regions were drawn in the left and right amygdala according to the intersection between an anatomical mask (Automated Anatomical Labeling; Tzourio-Mazoyer *et al*, 2002) and the result of a meta-analysis of 163 fMRI studies on emotional processing (Wager *et al*, 2012). For each participant and each seed region, the physiological factor is represented by the first eigenvariate of the BOLD signal within the seed region, which were high-pass filtered with a cutoff of 128 s, prewhitened to account for serial autocorrelations of order 1 (AR(1)), adjusted for an F-contrast over all experimental conditions, and deconvolved using the canonical Haemodynamic Response Function. Interaction terms were calculated separately for each experimental condition as the product between the vector of the condition and the physiological factor. The resulting contrast image 'stress vs control' reveals stress-induced FC change to the corresponding seed region.

Statistical Analysis

First, to identify changes in FC pattern during acute psychosocial stress, a voxel-wise one-sample *T*-test was performed as second-level analysis on the contrast 'stress vs control' in the placebo session. To investigate the effect of EA on stress-induced FC changes in the placebo session, we performed a multiple regression analysis on the contrast image 'stress vs control' with EA as the main predictor and controlled for the following three covariates: age, IQ, and experiment sequence. All regressors were included in one model. Mean-centered predictors were used to reduce multicollinearity. Results are reported at a threshold of cluster-level FWE-corrected $p < 0.05$, cluster size > 50 voxels.

In brain regions where a main effect of EA was found, we built regions of interest (ROIs) and extracted FC strength for the stress and control conditions as beta values from the PPI model (see Supplementary Methods). Stress-induced FC changes were calculated as the differences in FC strength between the stress and control conditions. Previous studies have associated ELS with reduced gray matter volume (GMV) in limbic areas (Teicher *et al*, 2012) and with personality traits, such as neuroticism, trait anxiety, and self-esteem (Grimm *et al*, 2014). We further explored the influence of GMV and personality traits on the main effect of EA, using ROI-based multiple regression model (main predictor: EA; covariates: age, IQ, experimental sequence, GMV in seed region, GMV in the corresponding ROI, neuroticism, trait anxiety, and self-esteem).

To examine whether OXT and its joint effect with EA also modulates stress-induced FC change in these regions, we extracted stress-induced FC changes for the placebo and OXT sessions separately and performed a series of hierarchical multiple regression (HMR) analyses. For each ROI, five covariates were entered in the first step: age, IQ, experimental sequence, GMV in seed region, and GMV in the corresponding ROI. In the second step, EA and OXT were entered as main predictors. In the final step, an interaction OXT \times EA was entered. All predictors were mean centered. Before model estimation, we calculated Mahalanobis distance to make sure that multivariate outliers were excluded. We further identified data points that exerted an

excessive influence on the HMR model, using Cook's distances following each model estimation. Data points with a Cook's distance exceeding the sample specific cutoff (four/sample size) were removed from the analysis, and the HMR model was re-estimated. We report results from the final HMR models free of outliers and influential data points. To examine an interaction effect, we tested whether successive regression steps significantly increased the variance explained by the model (ΔR^2).

RESULTS

Psychological and Physiological Measures

One subject was identified as an outlier because of a score of 21 on the SA subscale. SA has been associated with distinct patterns of stress reactivity compared with other ELS subtypes (Carpenter *et al*, 2009). For this reason, we excluded this subject and performed analyses on the data from 31 subjects (mean age 28.2 ± 4.7 , mean IQ 112.5 ± 15.5 , four left handed). Scores on the EN, EA, and PN subscales range from the none/low to severe ELS defined by previous normative data (Scher *et al*, 2001) (ie, -1 SD \leq the 50th percentile of the norm; $+1$ SD \geq the 90th percentile of the norm; see Table 1). None of the subjects scored high on the subscale of PA and

Table 1 Demographic and Psychometric Measures

	Mean (SD)	Range
Age, years	28.2 (4.7)	21–37
IQ	112.5 (15.5)	92–145
<i>CTQ</i>		
Total score	37.8 (13.9)	25–84
Emotional neglect	10.5 (4.9)	5–23
Emotional abuse	8.5 (4.6)	5–23
Physical neglect	7.5 (3.7)	5–20
Physical abuse	6.3 (2.5)	5–15
Sexual abuse	5.1 (0.3)	5–6
<i>NEO-FFI</i>		
Neuroticism	19.1 (8.5)	4–36
Extroversion	29.3 (6.5)	12–40
Openness	32.8 (8.9)	11–46
Agreeableness	30.9 (6.1)	18–40
Conscientiousness	30.1 (7.4)	11–40
Rosenberg self-esteem scale	40.0 (5.9)	23–50
<i>STAI</i>		
Trait anxiety	36.5 (7.0)	25–48
State anxiety at baseline	35.0 (5.3)	26–49
State anxiety after stress (PLA)	39.8 (8.0)	24–63
State anxiety after stress (OXT)	41.2 (11.0)	22–64

Abbreviations: CTQ, Childhood Trauma Questionnaire; OXT, oxytocin; PLA, placebo.

SA subscale (ie, +1 SD not exceeding the 90th percentile of normative data; see Table 1).

Partial correlation analyses controlling for age and IQ showed that EA scores correlated positively with neuroticism ($r(23)=0.55$, $p<0.01$) and trait anxiety ($r(23)=0.40$, $p<0.05$) and negatively with self-esteem ($r(23)=-0.65$, $p<0.001$). We observed higher state anxiety levels ($t(28)=3.33$, $p<0.01$) and salivary cortisol levels ($t(18)=2.55$, $p<0.05$) after stress induction in the PLA session. EA correlated positively with state anxiety levels after stress induction regardless of the administration of OXT or PLA ($r_{\text{parital}}(50)=0.32$, $p<0.05$). HMR revealed a significant OXT \times EA interaction effect on the cortisol stress responses ($\beta=0.430$, $t(35)=2.83$, $p<0.001$). The full model with the interaction term was significant ($F(6,35)=2.493$, $p<0.05$), and adding the OXT \times EA interaction as an additional predictor explained a significant proportion of variance in cortisol stress responses ($\Delta R^2=0.16$, F Change(1,35)=8.01, $p<0.01$). *Post hoc* analysis showed a negative correlation between EA scores and cortisol stress responses in the PLA session ($r_{\text{parital}}(14)=-0.43$, $p=0.098$) and a positive correlation in the OXT session ($r_{\text{parital}}(18)=0.48$, $p<0.05$) (see Supplementary Figure S1).

Effect of Psychosocial Stress on Amygdala FC

To investigate the effect of psychosocial stress on amygdala coupling, we performed a PPI analysis for each subject's fMRI data in the placebo session. Whole-brain one-sample *T*-tests at group level revealed brain regions where activity was more positively coupled with that of bilateral amygdalae during psychosocial stress than during the control condition. Among these are the pregenual anterior cingulate cortex (pgACC), mPFC, posterior cingulate cortex (PCC), left anterior insula, caudate, putamen, right thalamus, inferior parietal lobule (IPL), and bilateral cerebellum (see Supplementary Table S1, Supplementary Figure S2).

Effect of EA on Stress-Induced Amygdala FC Shift

Voxel-wise multiple regression in the placebo session further revealed regions where such stress-induced connectivity changes were modulated by EA. Specifically, as severity of EA increased, there was stronger stress-induced FC ascent between bilateral amygdalae and bilateral hippocampus extending to parahippocampal gyrus (PHG), bilateral fusiform gyrus (FFG), and cerebellum (see Table 2, Figure 1a).

In subsequent ROI-based multiple regression analysis controlling for age, IQ, experimental sequence, GMV, neuroticism, trait anxiety, and self-esteem, the main effect of EA on stress-induced FC enhancement remained significant for FC between the right amygdala seed and bilateral hippocampus extending to PHG, left FFG, and bilateral cerebellum (all β 's >0.64 , all p 's <0.004), as well as for FC between the left amygdala seed and right PHG, right FFG, and left precuneus (all β 's >0.67 , all p 's <0.004). We also observed a significant association between neuroticism and stress-induced FC enhancement between the right amygdala seed and bilateral hippocampus/PHG (all β 's <-0.45 , all p 's <0.05).

Table 2 Brain Regions in Which Stress-Induced Connectivity Changes were Predicted by Emotional Abuse

Region	Seed	
	Left amygdala	Right amygdala
Left Hipp/PHG		-30, -39, -12 Z=3.9
Right Hipp	39, -24, -15 Z=3.29	30, -33, -18 Z=4.28
Right PHG	24, -33, -15 Z=3.57	36, -39, -15 Z=4.07
Left fusiform gyrus		-33, -39, -21 Z=4.08
Right fusiform gyrus	42, -42, -21 Z=4.11	
Left precuneus	-15, -75, 45 Z=3.99	
Left cerebellum	-6, -57, -30 Z=4.18	-6, -57, -27 Z=4.55
Right cerebellum		30, -48, -18 Z=3.7

Abbreviations: Hipp, hippocampus; PHG, parahippocampal gyrus. Results are reported at a threshold of cluster-level FWE-corrected $p<0.05$, cluster size >50 voxels. Coordinates are reported according to the MNI space.

Joint Effect of EA and OXT on Stress-Induced Amygdala FC Shift

To examine the main effect of OXT and the joint effect of OXT and EA on the stress-induced connectivity shift between the amygdala and hippocampus, a ROI was drawn in the right hippocampus where the effect of EA was observed. Stress-induced changes in amygdala-hippocampal FC were extracted for the placebo and OXT sessions separately. No significant main effect of OXT was found in the right hippocampus. HMR showed a significant OXT \times EA interaction ($\beta=-0.27$, $t(44)=-2.13$, $p<0.05$). The full model with the interaction term was significant ($R^2=0.42$, $F(8,44)=3.90$, $p<0.01$), and the OXT \times EA interaction explained a significant proportion of variance in amygdala-hippocampal FC ($\Delta R^2=0.06$, F Change(1,44)=4.53, $p<0.05$). This ROI analysis further revealed that, in the placebo session, increased EA scores were associated with a positive shift in hippocampal-amygdala FC during psychosocial stress; while in the OXT session there was no significant correlation between EA scores and stress-induced FC changes between hippocampus and amygdala (see Figure 1b).

Similar OXT \times EA interactions were also found in stress-induced FC changes from right amygdala to right PHG ($\beta=-0.26$, $t(43)=-2.23$, $p<0.05$), from left amygdala to left fusiform gyrus ($\beta=-0.34$, $t(44)=-3.27$, $p<0.01$), as well as from bilateral amygdalae to left cerebellum (for l Amg, $\beta=-0.41$, $t(44)=-2.88$, $p<0.01$; for r Amg, $\beta=-0.47$, $t(45)=-3.97$, $p<0.001$; for detailed results of the HMR models, please see Supplementary Table S2).

Correlation analyses with the physiological data showed that stress-induced FC changes between the right amygdala




Figure 1 (a) Voxel-wise multiple regression analysis of the placebo session showed that emotional abuse positively predicted stress-induced connectivity enhancement between the amygdala and the illustrated regions (cluster-level FWE-corrected $p < 0.05$, cluster size > 50 voxels). (b) Results of the hierarchical multiple regression estimating the joint effect of oxytocin and emotional abuse on stress-induced connectivity shift between right amygdala and right hippocampus. (c) Cortisol stress response correlated negatively with the stress-induced connectivity shift between right amygdala and right hippocampus *Comu Ammonis* (CA) areas during the oxytocin session but not the placebo session. Amg, amygdala; Cb, cerebellum; Hipp, hippocampus; OXT, oxytocin; PCu, precuneus; PLA, placebo.

and right hippocampus correlated negatively with cortisol stress responses during the OXT session ($r(21) = -0.453$, $p < 0.05$) but not during the placebo session ($r(16) = 0.187$, $p > 0.45$; Fischer's $Z = 1.98$, $p < 0.05$; see Figure 1c). Similar correlation pattern was also found in stress-induced FC changes between the right amygdala and right PHG (for OXT session, $r(20) = -0.697$, $p < 0.001$; for PLA session, $r(15) = 0.137$, $p > 0.60$; Fischer's $Z = 2.83$, $p < 0.001$).

DISCUSSION

The first aim of the present study was to examine the impact of EA on stress-induced changes in amygdala-centered FC. Consistent with previous findings (Hermans *et al*, 2011), psychosocial stress induced increased FC strength between the amygdala and distributed stress-related cortical-subcortical regions when compared with the control condition.

Among these are regions associated with salience processing and autonomic-neuroendocrine control (ie, mPFC/pgACC, insula, caudate, putamen, and thalamus), as well as attentional (re)orientation and executive control (ie, MCC, PCC/precuneus, IPL, and cerebellum). Moreover, early life experience of EA positively predicted stress-induced FC enhancement between bilateral amygdalae and the right hippocampus/PHG. This linear association remained significant after controlling for individual differences in age, IQ, experimental sequence, GMV of corresponding areas, neuroticism, self-esteem, and trait anxiety. ELS has previously been associated with limbic hyper-responsiveness to negative facial cues during early stage of emotion processing (Dannowski *et al*, 2013). Impaired structural connectivity within the limbic-prefrontal circuit has also been observed in subjects who have experienced ELS (Hart and Rubia, 2012). Recently, ELS has also been associated with decreased amygdala-mPFC and hippocampal-mPFC FC during

resting state (Burghy *et al*, 2012; Fan *et al*, 2014; Herringa *et al*, 2013), suggesting ELS-associated aberration in the trait-like functional integrity of the limbic–prefrontal circuit. Adding to this evidence, our finding highlights the association between EA and transient task-evoked changes in amygdala–hippocampal FC in response to acute psychosocial stress.

The amygdala–hippocampal pathway is crucially involved in the regulation of HPA axis (Herman *et al*, 2005). Kiem *et al* (2013) reported that stronger amygdala–hippocampal rs-FC is associated with dampened cortisol responses to Dex/CRH test, which indicates decreased efficiency of the HPA axis to re-establish homeostasis after perturbation. Vaisvaser *et al* (2013) found prolonged increases in amygdala–hippocampal FC even 2 h after psychosocial stress, which was associated with blunted cortisol stress responses. These findings associate increased amygdala–hippocampal coupling with dysregulated HPA stress reactivity, especially aberrant glucocorticoid negative feedback. Substantial evidence associates ELS with dysregulated cortisol stress responses and HPA reactivity (Strüber *et al*, 2014). It has recently been found that ELS can induce epigenetic modification in the hippocampus, which is related to deficiency in HPA regulation via glucocorticoid negative feedback (Klengel *et al*, 2013). Consistent with these previous findings, the ELS-associated enhancement in stress-induced amygdala–hippocampal FC may mediate the impact of ELS on neuroendocrine control. The stress regulation circuit and the neural circuit for learning and memory largely overlap in the amygdala and hippocampus and enhanced connectivity between these two regions has an important role in both encoding and consolidation of emotional memory (Richter-Levin and Akirav, 2000). Acute stress alters hippocampal activity during encoding and affects memory performance (Henckens *et al*, 2009). The ELS-associated enhancement in stress-induced amygdala–hippocampal FC changes may therefore indicate an impact of ELS on memory during stress.

It should be noted that although we observed stress-induced changes in FC between bilateral amygdalae and the mPFC/pgACC, these were not influenced by EA even at ROI level (see Supplementary Material). At a first glance, this seems inconsistent with previous findings that ELS is associated with decreased amygdala–prefrontal FC during resting state (Burghy *et al*, 2012; Fan *et al*, 2014; Herringa *et al*, 2013). However, it is important to emphasize that task-evoked FC changes reflect transient changes in brain network coupling in response to specific task demands. The stress-induced FC change assesses reactivity of the stress regulation circuit in response to a transient challenge, which is complimentary to the trait-like measurement of task-independent FC during resting state. Cousijn *et al* (2012) recently showed that stress induction modulates both tonic and phasic activity in the amygdala but only phasic activity in the hippocampus. The ELS-associated enhancement in stress-induced amygdala–hippocampal FC is likely to reflect the impact of ELS on phasic FC changes in the amygdala–hippocampal pathway in response to acute stress, while rs-FC changes between amygdala and the mPFC/pgACC may reflect the influence of ELS on more tonic network properties in the amygdala–prefrontal pathway.

Furthermore, we found a joint effect of OXT and EA on the stress-induced changes in amygdala–hippocampal FC. Specifically, OXT induced a positive shift in stress-induced FC changes in subjects with lower EA scores but a negative shift in subjects with higher EA scores. The changes in amygdala–hippocampal FC in the OXT session were negatively associated with cortisol stress responses. This concurred with the physiological measure that cortisol stress responses in the OXT condition were reduced in subjects with lower EA scores but were increased in subjects with higher EA scores. Previous findings demonstrated that OXT modulates HPA stress responses, and this stress-attenuating effect is mediated via the inhibition of central–medial amygdala (Huber *et al*, 2005). It has further been reported that the stress-attenuating effect of OXT is altered in subjects who have experienced ELS (Bakermans-Kranenburg and van IJzendoorn, 2013). Our recent study revealed that alteration in stress-induced limbic deactivation is involved in the ELS-dependent sensitivity to OXT (Grimm *et al*, 2014). In addition to these findings, we demonstrated ELS-dependent OXT effects on stress-induced changes in amygdala–hippocampal FC. Considering the inverse association between amygdala–hippocampal FC and cortisol stress responses, our finding is in concordance with the altered stress-attenuating effect of OXT in subjects who have experienced ELS. To our knowledge, this is the first study associating stress-induced phasic FC changes in the amygdala–hippocampal pathway with ELS-dependent sensitivity to OXT.

It is worth noticing that the oxytocinergic system itself is highly sensitive to the impact of ELS (Heim *et al*, 2009; Hill *et al*, 2014). From an early developmental stage, the endogenous secretion of OXT and the expression of OXT receptors (especially in the amygdala) are modulated by the amount and quality of maternal affectionate behavior (Hill *et al*, 2014). Other than the ELS-associated enhancement of stress-induced amygdala–hippocampal FC, ELS-associated dysfunction in the oxytocinergic system itself may also be an intermediate factor for altered OXT effects.

It has been proposed that OXT is advantageous for the treatment of stress-related psychiatric disorders such as social phobia due to its stress-attenuating effect (see review in Bakermans-Kranenburg and van IJzendoorn, 2013). However, recent evidence reveals that OXT is not always beneficial depending on various modulating factors, eg, the personal history of ELS (Bakermans-Kranenburg and van IJzendoorn, 2013). Regarding the neural mechanism of the ELS-dependent sensitivity to OXT, we have previously shown differential OXT effects on pgACC stress responses depending on resting-state amygdala–pgACC FC before stress induction (Fan *et al*, 2014). Our findings in the present study provide further support for the differential effects of OXT on task-evoked amygdala–hippocampus FC during stress. As both trait-like basal rs-FC and stress-induced phasic FC changes within the limbic–prefrontal circuit are modulated by ELS, these findings emphasize the necessity to consider the tonic basal state and phasic functional reactivity of the limbic–prefrontal circuit when assessing the stress-attenuating effect of OXT.

Although our study provides new insights regarding the joint effect of EA and OXT on stress-induced changes in amygdala–hippocampal FC, several limitations need to be considered. The CTQ assesses ELS based on retrospective

self-report and is hence susceptible to subjectivity bias in perception of early environment and in reconstructive memory. We did not include any 'objective' measure of ELS (eg, medical records). However, it was mainly on the EA and EN subscale that our subjects reported moderate-to-severe ELS (Bernstein and Fink, 1998; Scher *et al*, 2001). Such information on emotional maltreatment is unlikely to be reflected in medical records. In addition, many subjects were not in contact with their families, so that family interviews would have been difficult to arrange. As the perception and recall of ELS are influenced by various factors, including personality traits and coping skills, it is important to acknowledge the subjectivity in our measurement of EA when interpreting the findings. Previous studies showed differential effects of ELS on stress reactivity in a gender- and age-specific manner, depending on the specific type and severity of ELS, as well as the presence/absence of concurrent stress and psychiatric symptoms (Heim and Binder, 2012; Strüber *et al*, 2014). In the present study, we recruited only healthy young male subjects and used EA as the main predictor in our multiple regression models. Although this helps to delineate the effect of EA from that of other subtypes of ELS and from the influence of psychiatric symptoms, one should be cautious when generalizing our findings to female samples or subjects with concurrent psychiatric symptoms. As we did not obtain any longitudinal data, it is not known whether the subjects will develop psychiatric disorders later in life. Future studies are needed to differentiate individuals vulnerable and resilient to the detrimental influence of ELS. It is also important to point out that the pharmacokinetics of OXT and its effect at molecular biology level remain unknown. The exact effect of OXT on limbic-prefrontal network may be much more complicated on a micro-circuitry level. For example, it has been suggested that OXT inhibits activity in the amygdala via its interaction with the GABAergic and serotonergic system within different nuclei of the amygdala complex (Strüber *et al*, 2014). Although our findings associate the limbic-prefrontal circuit with the stress-attenuating effect of OXT on a macro-network scale, further cellular biological research is needed to elucidate the effect of OXT on a micro-circuitry level.

In conclusion, we found that EA is positively associated with enhanced stress-induced FC strength between the amygdala and hippocampus, whereas this association is not significant after the administration of intranasal OXT. Stress-induced amygdala-hippocampal FC is crucial in the neural mechanism underlying the impact of ELS on stress reactivity and may also be involved in the ELS-dependent stress-attenuating effect of OXT.

FUNDING AND DISCLOSURE

This work was supported by the German Research Foundation (DFG, Cluster of Excellence 'Languages of Emotion'). The authors declare no conflict of interest.

ACKNOWLEDGMENTS

We thank the team of Dahlem Institute for Neuroimaging of Emotion for their technical support, Jue Wang for helpful

suggestions with the data analysis, and Anne Weigand and Matti Gärtner for comments on the manuscript.

REFERENCES

- Bakermans-Kranenburg MJ, van IJzendoorn MH (2013). Sniffing around oxytocin: review and meta-analyses of trials in healthy and clinical groups with implications for pharmacotherapy. *Transl Psychiatry* 3: e258.
- Bernstein DP, Fink L (1998). *Childhood Trauma Questionnaire: A retrospective self-report manual*. The Psychological Corporation: San Antonio, TX, USA San Antonio, TX, USA.
- Burghy CA, Stodola DE, Ruttle PL, Molloy EK, Armstrong JM, Oler JA *et al* (2012). Developmental pathways to amygdala-prefrontal function and internalizing symptoms in adolescence. *Nat Neurosci* 15: 1736–1741.
- Carpenter LL, Tyrka AR, Ross NS, Khoury L, Anderson GM, Price LH (2009). Effect of childhood emotional abuse and age on cortisol responsivity in adulthood. *Biol Psychiatry* 66: 69–75.
- Costa PT, McCrae RR (1985). *The NEO Personality Inventory manual*. Psychological Assessment Resources: Odessa, FL, USA.
- Cousijn H, Rijpkema M, Qin S, van Wingen GA, Fernandez G (2012). Phasic deactivation of the medial temporal lobe enables working memory processing under stress. *Neuroimage* 59: 1161–1167.
- Dannlowski U, Kugel H, Huber F, Stuhrmann A, Redlich R, Grotegerd D *et al* (2013). Childhood maltreatment is associated with an automatic negative emotion processing bias in the amygdala. *Hum Brain Mapp* 34: 2899–2909.
- Dressendorfer RA, Kirschbaum C, Rohde W, Stahl F, Strasburger CJ (1992). Synthesis of a cortisol-biotin conjugate and evaluation as a tracer in an immunoassay for salivary cortisol measurement. *J Steroid Biochem Mol Biol* 43: 683–692.
- Fan Y, Herrera-Melendez AL, Pestke K, Feeser M, Aust S, Otte C *et al* (2014). Early life stress modulates amygdala-prefrontal functional connectivity: Implications for oxytocin effects. *Hum Brain Mapp* 35: 5328–5339.
- Freese JL, Amaral DG (2009) Neuroanatomy of the primate amygdala. In: Whalen PJ, Phelps EA (eds). *The Human Amygdala*. The Guilford Press: New York, NY, USA, pp 3–42.
- Friston KJ, Buechel C, Fink GR, Morris J, Rolls E, Dolan RJ (1997). Psychophysiological and modulatory interactions in neuroimaging. *Neuroimage* 6: 218–229.
- Fuge P, Aust S, Fan Y, Weigand A, Gärtner M, Feeser M *et al* (2014). Interaction of early life stress and corticotropin-releasing hormone receptor gene: effects on working memory. *Biol Psychiatry* 76: 888–894.
- Gamer M, Zurowski B, Buchel C (2010). Different amygdala subregions mediate valence-related and attentional effects of oxytocin in humans. *Proc Natl Acad Sci USA* 107: 9400–9405.
- Gee DG, Gabard-Durnam LJ, Flannery J, Goff B, Humphreys KL, Telzer EH *et al* (2013). Early developmental emergence of human amygdala-prefrontal connectivity after maternal deprivation. *Proc Natl Acad Sci USA* 110: 15638–15643.
- Grimm S, Pestke K, Feeser M, Aust S, Weigand A, Wang J *et al* (2014). Early life stress modulates oxytocin effects on limbic system during acute psychosocial stress. *Soc Cogn Affect Neurosci* 9: 1828–1835.
- Hart H, Rubia K (2012). Neuroimaging of child abuse: a critical review. *Front Hum Neurosci* 6: 52.
- Heim C, Binder EB (2012). Current research trends in early life stress and depression: review of human studies on sensitive periods, gene-environment interactions, and epigenetics. *Exp Neurol* 233: 102–111.
- Heim C, Young LJ, Newport DJ, Mletzko T, Miller AH, Nemeroff CB (2009). Lower CSF oxytocin concentrations in women with a history of childhood abuse. *Mol Psychiatry* 14: 954–958.

- Henckens MJ, Hermans EJ, Pu Z, Joels M, Fernandez G (2009). Stressed memories: how acute stress affects memory formation in humans. *J Neurosci* **29**: 10111–10119.
- Herman JP, Ostrander MM, Mueller NK, Figueiredo H (2005). Limbic system mechanisms of stress regulation: hypothalamo-pituitary-adrenocortical axis. *Prog Neuropsychopharmacol Biol Psychiatry* **29**: 1201–1213.
- Hermans EJ, van Marle HJ, Ossewaarde L, Henckens MJ, Qin S, van Kesteren MT *et al* (2011). Stress-related noradrenergic activity prompts large-scale neural network reconfiguration. *Science* **334**: 1151–1153.
- Herrington RJ, Birn RM, Ruttle PL, Burghy CA, Stodola DE, Davidson RJ *et al* (2013). Childhood maltreatment is associated with altered fear circuitry and increased internalizing symptoms by late adolescence. *Proc Natl Acad Sci USA* **110**: 19119–19124.
- Hill KT, Warren M, Roth TL (2014). The influence of infant-caregiver experiences on amygdala Bdnf, OXTr, and NPY expression in developing and adult male and female rats. *Behav Brain Res* **272C**: 175–180.
- Holz NE, Buchmann AF, Boecker R, Blomeyer D, Baumeister S, Wolf I *et al* (2015). Role of FKBP5 in emotion processing: results on amygdala activity, connectivity and volume. *Brain Struct Funct* **200**: 1355–1368.
- Huber D, Veinante P, Stoop R (2005). Vasopressin and oxytocin excite distinct neuronal populations in the central amygdala. *Science* **308**: 245–248.
- Keyes KM, Eaton NR, Krueger RF, McLaughlin KA, Wall MM, Grant BF *et al* (2012). Childhood maltreatment and the structure of common psychiatric disorders. *Br J Psychiatry* **200**: 107–115.
- Kiem SA, Andrade KC, Spoomaker VI, Holsboer F, Czisch M, Samann PG (2013). Resting state functional MRI connectivity predicts hypothalamus-pituitary-axis status in healthy males. *Psychoneuroendocrinology* **38**: 1338–1348.
- Klengel T, Mehta D, Anacker C, Rex-Haffner M, Pruessner JC, Pariante CM *et al* (2013). Allele-specific FKBP5 DNA demethylation mediates gene-childhood trauma interactions. *Nat Neurosci* **16**: 33–41.
- Labuschagne I, Phan KL, Wood A, Angstadt M, Chua P, Heinrichs M *et al* (2010). Oxytocin attenuates amygdala reactivity to fear in generalized social anxiety disorder. *Neuropsychopharmacology* **35**: 2403–2413.
- LeDoux JE (2000). Emotion circuits in the brain. *Annu Rev Neurosci* **23**: 155–184.
- Lehrl S, Triebig G, Fischer B (1995). Multiple choice vocabulary test MWT as a valid and short test to estimate premorbid intelligence. *Acta Neurol Scand* **91**: 335–345.
- McEwen BS (2003). Mood disorders and allostatic load. *Biol Psychiatry* **54**: 200–207.
- McLaren DG, Ries ML, Xu G, Johnson SC (2012). A generalized form of context-dependent psychophysiological interactions (gPPI): a comparison to standard approaches. *Neuroimage* **61**: 1277–1286.
- Nelson HE, O'Connell A (1978). Dementia: the estimation of premorbid intelligence levels using the New Adult Reading Test. *Cortex* **14**: 234–244.
- Neumann ID, Wigger A, Torner L, Holsboer F, Landgraf R (2000). Brain oxytocin inhibits basal and stress-induced activity of the hypothalamo-pituitary-adrenal axis in male and female rats: partial action within the paraventricular nucleus. *J Neuroendocrinol* **12**: 235–243.
- Pechtel P, Pizzagalli DA (2011). Effects of early life stress on cognitive and affective function: an integrated review of human literature. *Psychopharmacology (Berl)* **214**: 55–70.
- Pruessner JC, Dedovic K, Khalili-Mahani N, Engert V, Pruessner M, Buss C *et al* (2008). Deactivation of the limbic system during acute psychosocial stress: evidence from positron emission tomography and functional magnetic resonance imaging studies. *Biol Psychiatry* **63**: 234–240.
- Richter-Levin G, Akirav I (2000). Amygdala-hippocampus dynamic interaction in relation to memory. *Mol Neurobiol* **22**: 11–20.
- Riem MM, van IMH, Tops M, Boksem MA, Rombouts SA, Bakermans-Kranenburg MJ (2012). No laughing matter: intranasal oxytocin administration changes functional brain connectivity during exposure to infant laughter. *Neuropsychopharmacology* **37**: 1257–1266.
- Rosenberg M (1965). *Society and the Adolescent Self-Image*. Princeton University Press: Princeton, NJ, USA Princeton, NJ, USA.
- Scher CD, Stein MB, Asmundson GJ, McCreary DR, Forde DR (2001). The childhood trauma questionnaire in a community sample: psychometric properties and normative data. *J Trauma Stress* **14**: 843–857.
- Schmidt KH, Metzler P (1992). *Wortschatztest (WST)*. Beltz Test GmbH: Weinheim, Germany Weinheim, Germany.
- Spielberger CD, Sydeman SJ (1994). *State-Trait Anxiety Inventory and State-Trait Anger Expression Inventory*. Lawrence Erlbaum Associates: Hillsdale, NJ, USA.
- Strüber N, Strüber D, Roth G (2014). Impact of early adversity on glucocorticoid regulation and later mental disorders. *Neurosci Biobehav Rev* **38**: 17–37.
- Teicher MH, Anderson CM, Polcari A (2012). Childhood maltreatment is associated with reduced volume in the hippocampal subfields CA3, dentate gyrus, and subiculum. *Proc Natl Acad Sci USA* **109**: E563–E572.
- Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N *et al* (2002). Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* **15**: 273–289.
- Vaisvaser S, Lin T, Admon R, Podlipsky I, Greenman Y, Stern N *et al* (2013). Neural traces of stress: cortisol related sustained enhancement of amygdala-hippocampal functional connectivity. *Front Hum Neurosci* **7**: 313.
- Veer IM, Oei NY, Spinhoven P, van Buchem MA, Elzinga BM, Rombouts SA (2011). Beyond acute social stress: increased functional connectivity between amygdala and cortical midline structures. *Neuroimage* **57**: 1534–1541.
- Wager TD, Barrett LF, Bliss-Moreau E, Lindquist KA, Duncan S, Kober H *et al* (2012) The neuroimaging of emotion In: Lewis M, Haviland-Jones JM, Barrett LF (eds). *Handbook of Emotions*. The Guilford Press: New York, NY, USA, pp 249–271.
- Wittchen HU, Wunderlich U, Gruschwitz S, Zaudig M (1997). *SKID-I. Strukturiertes Klinisches Interview für DSM-IV. Achse I: Psychische Störungen*. Hogrefe: Göttingen, Germany Göttingen, Germany.