

DETECTING OPTIMALITY AND EXTRACTING SOLUTIONS IN POLYNOMIAL OPTIMIZATION WITH THE TRUNCATED GNS CONSTRUCTION

MARÍA LÓPEZ QUIJORNA

ABSTRACT. A basic closed semialgebraic subset of \mathbb{R}^n is defined by simultaneous polynomial inequalities $p_1 \geq 0, \dots, p_m \geq 0$. We consider Lasserre's relaxation hierarchy to solve the problem of minimizing a polynomial over such a set. These relaxations give an increasing sequence of lower bounds of the infimum. In this paper we provide a new certificate for the optimal value of a Lasserre relaxation be the optimal value of the polynomial optimization problem. This certificate is that a modified version of an optimal solution of the Lasserre relaxation is a generalized Hankel matrix. This certificate is more general than the already known certificate of an optimal solution being flat. In case we have optimality we will extract the potential minimizers with a truncated version of the Gelfand-Naimark-Segal construction on the optimal solution of the Lasserre relaxation. We prove also that the operators of this truncated construction commute if and only if the matrix of this modified optimal solution is a generalized Hankel matrix. This generalization of flatness will bring us to reprove a result of Curto and Fialkow on the existence of quadrature rule if the optimal solution is flat and a result of Xu and Mysovskikh on the existence of a Gaussian quadrature rule if the modified optimal solution is generalized Hankel matrix. At the end, we provide a numerical linear algebraic algorithm for detecting optimality and extracting solutions of a polynomial optimization problem.

1. NOTATION

Throughout this paper, we suppose $n \in \mathbb{N} = \{1, 2, \dots\}$ and abbreviate (X_1, \dots, X_n) by \underline{X} . We let $\mathbb{R}[\underline{X}]$ denote the ring of real polynomials in n indeterminates. We denote $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$. For $\alpha \in \mathbb{N}_0^n$, we use the standard notation :

$$|\alpha| := \alpha_1 + \dots + \alpha_n \text{ and } \underline{X}^\alpha := X_1^{\alpha_1} \dots X_n^{\alpha_n}$$

For a polynomial $p \in \mathbb{R}[\underline{X}]$ we denote $p = \sum_{\alpha} p_{\alpha} \underline{X}^{\alpha}$ ($a_{\alpha} \in \mathbb{R}$). For $d \in \mathbb{N}_0$, by the notation $\mathbb{R}[\underline{X}]_d := \{\sum_{|\alpha| \leq d} a_{\alpha} \underline{X}^{\alpha} \mid a_{\alpha} \in \mathbb{R}\}$ we will refer to the vector space of polynomials with degree less or equal to d . Polynomials all of whose monomials have exactly the same degree $d \in \mathbb{N}_0$ are called d -forms. They form a finite dimensional vector space that we will denote by:

$$\mathbb{R}[\underline{X}]_{=d} := \left\{ \sum_{|\alpha|=d} a_{\alpha} \underline{X}^{\alpha} \mid a_{\alpha} \in \mathbb{R} \right\}$$

so that

$$\mathbb{R}[\underline{x}]_d = \mathbb{R}[\underline{X}]_0 \oplus \dots \oplus \mathbb{R}[\underline{X}]_d.$$

We will denote by $s_k := \dim \mathbb{R}[\underline{X}]_k$ and by $r_k := \dim \mathbb{R}[\underline{X}]_{=k}$. For $d \in \mathbb{N}_0$ we denote $\mathbb{R}[\underline{X}]_d^*$ the dual space of $\mathbb{R}[\underline{X}]_d$ i.e. the set of linear forms from $\mathbb{R}[\underline{X}]_d$ to \mathbb{R} and for $\ell \in \mathbb{R}[\underline{X}]_{2d}^*$ we denote by $\ell' := \ell|_{\mathbb{R}[\underline{X}]_{2d-2}}$ the restriction of the linear form ℓ to the space $\mathbb{R}[\underline{X}]_{2d-2}$. For $d \in \mathbb{N}_0$ and $a \in \mathbb{R}^n$ we denote $ev_a \in \mathbb{R}[\underline{X}]_d^*$ the linear form such that for all $p \in \mathbb{R}[\underline{X}]_d$, $ev_a(p) = p(a)$.

Date: April 6, 2017.

2010 Mathematics Subject Classification. Primary: 90C22, 90C26; Secondary: 44A60, 65D32.

Key words and phrases. Moment relaxation, Lasserre relaxation, polynomial optimization, semidefinite programming, quadrature, truncated moment problem, GNS construction.

2. INTRODUCTION

Let polynomials $f, p_1, \dots, p_m \in \mathbb{R}[\underline{X}]$ with $m \in \mathbb{N}_0$ be given. A polynomial optimization problem involves finding the infimum of f over the so called basic closed semialgebraic set S , defined by:

$$(1) \quad S := \{x \in \mathbb{R}^n \mid p_1(x) \geq 0, \dots, p_m(x) \geq 0\}$$

and also, if it is possible, a polynomial optimization problem involves extracting optimal points or minimizers i.e. elements in the set:

$$S^* := \{x^* \in S \mid \forall x \in S \ f(x^*) \leq f(x)\}$$

So from now on we will denote as (P) , to refer us to the above defined polynomial optimization problem, that is to say:

$$(2) \quad (P) \text{ minimize } f(x) \text{ subject to } x \in S$$

The optimal value of (P) , i.e. the infimum of $f(x)$ where x ranges over all feasible solutions S will be denoted by P^* , that is to say:

$$(3) \quad P^* := \inf\{f(x) \mid x \in S\} \in \{-\infty\} \cup \mathbb{R} \cup \{\infty\}$$

Note that $P^* = +\infty$ if $S = \emptyset$ and $P^* = -\infty$ if and only if f is unbounded from below on S , for example if $S = \mathbb{R}^n$ and f is of odd degree.

For $d \in \mathbb{N}_0$ let us define:

$$(4) \quad V_d := (1, X_1, X_2, \dots, X_n, X_1^2, X_1X_2, \dots, X_1X_n, \\ X_2^2, X_2X_3, \dots, X_n^2, \dots, X_n^d)^T$$

as a basis for the vector space of polynomials in n variables of degree at most d . Then

$$V_d V_d^T = \begin{pmatrix} 1 & X_1 & X_2 & \cdots & X_n^d \\ X_1 & X_1^2 & X_1X_2 & \cdots & X_1X_n^d \\ X_2 & X_1X_2 & X_2^2 & \cdots & X_2X_n^d \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ X_n^d & X_1X_n^d & X_2X_n^d & \cdots & X_n^{2d} \end{pmatrix} \in \mathbb{R}[\underline{X}]_{2d}^{s_d \times s_d}$$

Let us substitute for every monomial $\underline{X}^\alpha \in \mathbb{R}[\underline{X}]_{2d}$ a new variable Y_α . This matrix has the following form:

$$(5) \quad M_d := \begin{pmatrix} Y_{(0,\dots,0)} & Y_{(1,\dots,0)} & Y_{(0,1,\dots,0)} & \cdots & Y_{(0,\dots,1)} \\ Y_{(1,\dots,0)} & Y_{(2,\dots,0)} & Y_{(1,1,\dots,0)} & \cdots & Y_{(1,\dots,d)} \\ Y_{(0,1,\dots,0)} & Y_{(1,1,\dots,0)} & Y_{(0,2,\dots,0)} & \cdots & Y_{(0,1,\dots,d)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ Y_{(0,\dots,d)} & Y_{(1,\dots,d)} & Y_{(0,1,\dots,d)} & \cdots & Y_{(0,\dots,2d)} \end{pmatrix} \in \mathbb{R}[\underline{Y}]_1^{s_d \times s_d}$$

Definition 2.1. Every matrix $M \in \mathbb{R}^{s_d \times s_d}$ with the same shape than the matrix (5) is called a generalized Hankel matrix of order d . We denote the affine linear space of generalized Hankel matrix of order d by:

$$H_d := \{M_d(y) \mid y \in \mathbb{R}^{s_{2d}}\}$$

For $p \in \mathbb{R}[\underline{X}]_k$ denote $d_p := \lfloor \frac{k-\text{deg } p}{2} \rfloor$ and consider the following symmetric matrix:

$$(6) \quad pV_{d_p}^t V_{d_p} = \begin{pmatrix} p & pX_1 & pX_2 & \cdots & pX_n^{d_p} \\ pX_1 & pX_1^2 & pX_1X_2 & \cdots & pX_1X_n^{d_p} \\ pX_2 & pX_2X_1 & pX_2^2 & \cdots & pX_2X_n^{d_p} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ pX_n^{d_p} & pX_1X_n^{d_p} & pX_n^{d_p}X_2 & \cdots & pX_n^{2d_p} \end{pmatrix} \in \mathbb{R}[\underline{X}]_k^{s_{d_p} \times s_{d_p}}$$

Definition 2.2. For $p \in \mathbb{R}[X]_k$ the localizing matrix of p of degree k is the matrix resulting from substitute every monomial X^α such that $|\alpha| \leq k$ in (6) for a new variable Y_α . We denote this matrix by $M_{k,p} \in \mathbb{R}[Y]_1^{s_{d_p} \times s_{d_p}}$.

Definition 2.3. For a $t \times t$ real symmetric matrix A , the notation $A \succeq 0$ means that A is positive semidefinite, i.e. $a^T A a \geq 0$ for all $a \in \mathbb{R}^t$.

In order to give further characterizations of positive semidefiniteness, let us remember a very well know theorem in linear algebra.

Reminder 2.4. Suppose $A \in \mathbb{R}^{t \times t}$ is symmetric. Then there is a diagonal matrix $D \in \mathbb{R}^{t \times t}$ and $U \in \mathbb{R}^{t \times t}$ orthogonal matrix, i.e. $UU^T = U^T U = I_t$, such that $U^T A U = D$

Reminder 2.5. Let $A \in \mathbb{R}^{t \times t}$ symmetric. The following are equivalent:

- (1) $A \succeq 0$.
- (2) All eigenvalues of A are nonnegative.
- (3) There exists $B \in \mathbb{R}^{t \times t}$ such that $A = B^T B$.

Proof. (1) \implies (2). Suppose $A \succeq 0$ and take $\lambda \in \mathbb{R}$ and eigenvalue of A such that $Av = \lambda v$ for $v \in \mathbb{R}^t$ eigenvector of A , then $v \neq 0$. implying that $0 \leq v^T A v = v^T \lambda v = \lambda \|v\|^2$ then $\lambda \geq 0$.

(2) \implies (3). Suppose that all eigenvalues of A are nonnegative then by 2.4 there exists D diagonal matrix with nonnegative entries $\lambda_i \geq 0$ and U orthogonal matrix such that $U^T A U = D$. Take $B := R U^T \in \mathbb{R}^{t \times t}$ where R is the diagonal matrix which entries are $\sqrt{\lambda_i}$.

(3) \implies (1). Suppose there is $B \in \mathbb{R}^{t \times t}$ such that $A = B^T B$, take $a \in \mathbb{R}^t$ then $a^T A a = a^T B^T B a = (Ba)(Ba) = \|Ba\|^2 \geq 0$. \square

Definition 2.6. Let (P) be a polynomial optimization problem as in (2) and let $k \in \mathbb{N}_0 \cup \{\infty\}$ such that $f, p_1, \dots, p_m \in \mathbb{R}[X]_k$. The Moment relaxation (or Lasserre relaxation) of (P) of degree k is the following semidefinite optimization problem:

$$(7) \quad (P_k) \text{ minimize } \sum_{|\alpha| \leq k} f_\alpha y_\alpha \text{ subject to}$$

$$M_{k,1}(y) \succeq 0, \quad y_{(0,\dots,0)} = 1, \quad M_{k,p_i}(y) \succeq 0$$

the optimal value of (P_k) that is to say, the infimum over all

$$y = (y_{(0,\dots,0)}, \dots, y_{(0,\dots,k)}) \in \mathbb{R}^{s_k}$$

that ranges over all feasible solutions of (P_k) is denoted by $P_k^* \in \{-\infty\} \cup \mathbb{R} \cup \{\infty\}$.

Given a polynomial optimization problem (P) as in (2) and $M := M_d(y) \in \mathbb{R}^{s_d \times s_d}$ an optimal solution of (P_{2d}) , it is always possible to find a matrix $W_M \in \mathbb{R}^{s_d \times r_d}$ such that M can be decomposed in a block matrix of the following form (see 4.8 below for a proof):

$$M = \left(\begin{array}{c|c} A_M & A_M W_M \\ \hline W_M^T A_M & C_M \end{array} \right)$$

This useful result can be also found in [23] and in [3, Lemma 2.3]. Define the following matrix:

$$\widetilde{M} := \left(\begin{array}{c|c} A_M & A_M W_M \\ \hline W_M^T A_M & W_M^T A_M W_M \end{array} \right)$$

In this paper we prove that \widetilde{M} is well-defined, that is to say it does not depend from the election of W_M , and assuming that $W_M^T A_M W_M$ is a generalized Hankel matrix we will use a new method to find a decomposition:

$$(8) \quad \widetilde{M} = \sum_{i=1}^r \lambda_i V_d(a_i) V_d(a_i)^T$$

where $r := \text{rank } M$, $a_1, \dots, a_r \in \mathbb{R}^n$ and $\lambda_1 > 0, \dots, \lambda_r > 0$. In this paper we will show that for some polynomial optimization problems if we have that $W_M^T A_M W_M$ is generalized Hankel and the nodes are contained in S , even if M is not flat i.e. $W_M^T A_M W_M \neq C_M$ (see the definition in 5.17), we can still claim optimality, that is to say that a_1, \dots, a_r are global minimizers. We will also see some examples to discard optimality or in other

words to discard that M has a factorization as in (8), see 6.7. Let us advance two results concerning optimality.

Theorem 2.7. Let (P) be a polynomial optimization problem as in (2) and suppose that $M_d(y) \in \mathbb{R}^{s_d \times s_d}$ is an optimal solution of (P_{2d}) and $\widetilde{M_d(y)}$ is a generalized Hankel matrix. Then there are $a_1, \dots, a_r \in \mathbb{R}^n$ points and $\lambda_1 > 0, \dots, \lambda_r > 0$ weights such that:

$$(9) \quad \widetilde{M_d(y)} = \sum_{i=1}^r \lambda_i V_d(a_i) V_d(a_i)^T$$

where $r = \text{rank } A_M$. Moreover if $\{a_1, \dots, a_r\} \subseteq S$ and $f \in \mathbb{R}[\underline{X}]_{2d-1}$ then a_1, \dots, a_r are global minimizers of (P) and $P^* = P_{2d}^* = f(a_i)$ for all $i \in \{1, \dots, r\}$.

Proof. The correspondence given in 3.5 together with the Theorem 7.1 will give us the proof. \square

Remark 2.8. Let (P) be a polynomial optimization problem without constraints. Suppose $M_d(y) \in \mathbb{R}^{s_d \times s_d}$ is an optimal solution of (P_{2d}) with $\widetilde{M_d(y)}$ a generalized Hankel matrix and that $f \in \mathbb{R}[\underline{X}]_{2d-1}$. Applying Theorem 2.7 we get the decomposition (9), and since we can ensure that $a_1, \dots, a_r \subseteq S = \mathbb{R}^n$ then they are global minimizers of (P) and $P^* = P_{2d}^* = f(a_i)$ for all $i \in \{1, \dots, r\}$.

Example 2.9. Let us consider the following polynomial optimization problem taken from [8, Problem 4.7]:

$$\begin{aligned} & \text{minimize} && f(x) = -12x_1 - 7x_2 + x_2^2 \\ & \text{subject to} && -2x_1^4 + 2 - x_2 = 0 \\ & && 0 \leq x_1 \leq 2 \\ & && 0 \leq x_2 \leq 3 \end{aligned}$$

We get the optimal value $P_4^* = -16.7389$ associated to the following optimal solution:

$$(10) \quad M_2(y) = \begin{array}{c} \begin{array}{cccccc} & 1 & X_1 & X_2 & X_1^2 & X_1 X_2 & X_2^2 \\ \begin{array}{l} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \end{array} & \begin{pmatrix} 1.0000 & 0.7175 & 1.4698 & 0.5149 & 1.0547 & 2.1604 \\ 0.7175 & 0.5149 & 1.0547 & 0.3694 & 0.7568 & 1.5502 \\ 1.4698 & 1.0547 & 2.1604 & 0.7568 & 1.5502 & 3.1755 \\ 0.5149 & 0.3694 & 0.7568 & 0.2651 & 0.5430 & 1.1123 \\ 1.0547 & 0.7568 & 1.5502 & 0.5430 & 1.1123 & 2.2785 \\ 2.1604 & 1.5502 & 3.1755 & 1.1123 & 2.2785 & 8.7737 \end{pmatrix} \end{array} \end{array}$$

and the modified moment matrix of $M_2(y)$ is the following:

$$(11) \quad \widetilde{M_2(y)} = \left(\begin{array}{ccc|ccc} 1.0000 & 0.7175 & 1.4698 & 0.5149 & 1.0547 & 2.1604 \\ 0.7175 & 0.5149 & 1.0547 & 0.3694 & 0.7568 & 1.5502 \\ 1.4698 & 1.0547 & 2.1604 & 0.7568 & 1.5502 & 3.1755 \\ \hline 0.5149 & 0.3694 & 0.7568 & 0.2651 & 0.5430 & 1.1123 \\ 1.0547 & 0.7568 & 1.5502 & 0.5430 & 1.1123 & 2.2785 \\ 2.1604 & 1.5502 & 3.1755 & 1.1123 & 2.2785 & \mathbf{4.6675} \end{array} \right)$$

We get that $\widetilde{M_2(y)}$ is a generalized Hankel matrix and $f \in \mathbb{R}[X_1, X_2]_3$ to conclude optimality, according with Theorem 2.7, it remains to calculate the factorization (8) and check if the points are in S . We will see in Section 5 in 5.20 how to compute this factorization, in this case, it is easy to see that:

$$\widetilde{M_2(y)} = V_2(\alpha, \beta) V_2(\alpha, \beta)^T$$

where $\alpha := 0.7175$ and $\beta := 1.4698$. One can verify that $(\alpha, \beta) \in S$ and therefore we can conclude that $P_4^* = P^* = -16.7389$ is the optimal value and (α, β) is a minimizer.

Theorem 2.10. Let (P) be a polynomial optimization problem given as in (2) and suppose that the p_i from (1) are all of degree at most 1 (so that S is a polyhedron). Suppose that $M_d(y) \in \mathbb{R}^{s_d \times s_d}$ is an optimal solution of (P_{2d}) and that $\widetilde{M}_d(y)$ is a generalized Hankel matrix. Then there are $a_1, \dots, a_r \in S$ and $\lambda_1 > 0, \dots, \lambda_r > 0$ weights such that:

$$\widetilde{M}_d(y) = \sum_{i=1}^r \lambda_i v_d(a_i) v_d(a_i)^T$$

Moreover if $f \in \mathbb{R}[X]_{2d-1}$ then a_1, \dots, a_r are global minimizers of (P) and $P^* = P_{2d}^* = f(a_i)$ for all $i = 1, \dots, r$.

Proof. The correspondence given in Corollary 3.5 together with the Theorem 7.3 will give us the result. \square

Example 2.11. Let us consider the following polynomial optimization problem, taken from [10, page 18], whose objective function is the Mzotkin polynomial [14, Prop.1.2.2]:

$$\begin{aligned} \text{minimize} \quad & f(x) = x_1^4 x_2^2 + x_1^2 x_2^4 - 3x_1^2 x_2^2 + 1 \\ \text{subject to} \quad & -2 \leq x_1 \leq 2 \\ & -2 \leq x_2 \leq 2 \end{aligned}$$

We get the optimal value $P_8^* = 6.2244 \cdot 10^{-9}$ from the following optimal solution of (P_8) :

$$(12) \quad M := M_{8,1}(y) = \left(\begin{array}{c|c} A_M & A_M W_M \\ \hline W_M^T A_M & C_M \end{array} \right)$$

where:

$$(13) \quad A_M = \begin{matrix} & 1 & X_1 & X_2 & X_1^2 & X_1 X_2 & X_2^2 & X_1^3 & X_1^2 X_2 & X_1 X_2^2 & X_2^3 \\ \begin{matrix} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \\ X_1^3 \\ X_1^2 X_2 \\ X_1 X_2^2 \\ X_2^3 \end{matrix} & \begin{pmatrix} 1.0000 & -0.0005 & -0.0004 & 1.0000 & -0.0000 & 1.0000 & -0.0005 & -0.0004 & -0.0005 & -0.0004 & -0.0004 \\ -0.0005 & 1.0000 & -0.0000 & -0.0005 & -0.0004 & -0.0005 & 1.0000 & -0.0000 & -0.0000 & 1.0000 & -0.0000 \\ -0.0004 & -0.0000 & 1.0000 & -0.0004 & -0.0005 & -0.0005 & -0.0004 & -0.0000 & -0.0004 & -0.0000 & 1.0000 \\ 1.0000 & -0.0005 & -0.0004 & 1.0000 & -0.0000 & 1.0000 & -0.0005 & -0.0004 & -0.0005 & -0.0004 & -0.0004 \\ -0.0000 & -0.0004 & -0.0005 & -0.0000 & 1.0000 & -0.0000 & -0.0004 & -0.0005 & -0.0005 & -0.0004 & -0.0005 \\ 1.0000 & -0.0005 & -0.0004 & 1.0000 & -0.0000 & -0.0000 & 1.0000 & -0.0005 & -0.0004 & -0.0005 & -0.0004 \\ -0.0005 & 1.0000 & -0.0000 & -0.0005 & -0.0004 & -0.0005 & 1.0001 & -0.0000 & 1.0001 & -0.0000 & -0.0000 \\ -0.0004 & -0.0000 & 1.0000 & -0.0004 & -0.0005 & -0.0004 & -0.0000 & 1.0001 & -0.0000 & 1.0001 & 1.0001 \\ -0.0005 & 1.0000 & -0.0000 & -0.0005 & -0.0004 & -0.0005 & 1.0001 & -0.0000 & 1.0001 & -0.0000 & -0.0000 \\ -0.0004 & -0.0000 & 1.0000 & -0.0004 & -0.0005 & -0.0004 & -0.0000 & 1.0001 & -0.0000 & 1.0001 & 1.0001 \end{pmatrix} \end{matrix}$$

$$(14) \quad W_M = \begin{pmatrix} 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \text{ and } C_M = \begin{matrix} & X_1^4 & X_1^3 X_2 & X_1^2 X_2^2 & X_1 X_2^3 & X_2^4 \\ \begin{matrix} X_1^4 \\ X_1^3 X_2 \\ X_1^2 X_2^2 \\ X_1 X_2^3 \\ X_2^4 \end{matrix} & \begin{pmatrix} 6.4115 & -0.0000 & 2.0768 & -0.0000 & 1.7719 \\ -0.0000 & 2.0768 & -0.0000 & 1.7719 & -0.0000 \\ 2.0768 & -0.0000 & 1.7719 & -0.0000 & 2.0768 \\ -0.0000 & 1.7719 & -0.0000 & 2.0768 & -0.0000 \\ 1.7719 & -0.0000 & 2.0768 & -0.0000 & 6.4115 \end{pmatrix} \end{matrix}$$

In this case:

$$W^* A_M W = \begin{pmatrix} 1.0000 & -0.0000 & 1.0000 & -0.0000 & 1.0000 \\ -0.0000 & 1.0000 & -0.0000 & 1.0000 & -0.0000 \\ 1.0000 & -0.0000 & 1.0000 & -0.0000 & 1.0000 \\ -0.0000 & 1.0000 & -0.0000 & 1.0000 & -0.0000 \\ 1.0000 & -0.0000 & 1.0000 & -0.0000 & 1.0000 \end{pmatrix}$$

is a Hankel matrix, what implies that \widetilde{M} is generalized Hankel and since we are minimizing over a polyhedron defined by linear polynomials by Theorem 2.10 $P_8^* = P^*$.

The goal of this paper is to find optimality conditions and extracting global minimizers from an optimal solution of the moment relaxation. That is to say given a polynomial optimization problem (P) as in (2) and an optimal solution of the moment relaxation (P_k) as in 2.6, find conditions to conclude if the optimal value is also the optimal value of the original polynomial optimization problem, i.e. $P^* = P_k^*$ and in this case extracting global minimizers. In the first section we outline Lasserres approach [11] to solve polynomial optimization problems with the language of linear forms, at the end of this section we will reformulate the problem of optimality, that is to say we reformulate the problem of

finding a decomposition of the modified moment matrix as in (9) to the problem of finding a commutative truncated version of the Gelfand-Naimark-Segal construction for a linear form $L \in \mathbb{R}[\underline{X}]_{2d}^*$ with $d \in \mathbb{N}_0 \cup \{\infty\}$, which take nonnegative values in $\sum \mathbb{R}[\underline{X}]_d^2$. The truncated GNS construction for this linear form will be defined in Section 4 and at the end of this section we give a proof of the very useful result of Smul'jan [23] using the inner product defined in the truncated GNS construction. In Section 5 we prove that if the truncated GNS multiplication operators of the optimal solution commute we are able to get the factorization (8) or in other words we find a Gaussian quadrature rule 5.11 representation for the linear form. In this section we will also prove that the commutativity of the truncated GNS operators is a more general fact than the very well know flatness condition, that is the case $C_M = W_M^T A_M W_M$, but the reverse it does not always hold (see (10), 5.19, 46, (47), (50) for examples), at the end of this section we review a result of Curto and Fialkow for the characterization of linear forms with quadrature rule on the whole space with minimal number of nodes. In Section 6 we prove the main result, which is that the truncated GNS multiplication operators of M commute if and only if $W_M^T A_M W_M$ is a Hankel matrix. This fact will help us to detect optimality in polynomial optimization problems and to slightly generalize some classical results of Dunkl, Xu, Mysovskikh, Möller and Putinar [7, Theorem 3.8.7], [16, 17], [18, pages 189-190] on Gaussian quadrature rules, with underlying ideas of [18]. In the last section we group all the results about optimality and global minimizers for an optimal solution of the moment relaxation, at the end we also give an algorithm for detecting the optimality and extracting minimizers with numerical examples.

3. FORMULATION OF THE PROBLEM

To solve polynomial optimization problems we use the very well known moment relaxations defined in 2.6. An introduction in to moment relaxations can also be found for instance in: [13], [11] and [21]. Likewise we will give the equivalent definition using linear forms instead of matrices in 3.4. We will now outline Lasserre's [11] approach to solve this problem. This method constructs a hierarchy of semidefinite programming relaxations, which are generalization of linear programs, and possible to solve efficiently, see [22] and [13] for an introduction. In each relaxation of degree k we build convex set, obtained through the linearization of a equivalent polynomial optimization problem of (P) defined in (2). This equivalent formulation of the problem consists in adding infinitely many redundant inequalities of the form $p \geq 0$ for all $p \in \sum \mathbb{R}[\underline{X}]^2 p_i \cap \mathbb{R}[\underline{X}]_k$ (with the notation $\sum \mathbb{R}[\underline{X}]^2 p_i$ we mean the set of all finite sums of elements of the form $p^2 p_i$, for $p \in \mathbb{R}[\underline{X}]$). The set of this redundant inequalities builds a cone, which is a set containing 0, closed under addition and closed under multiplication for positive scalars. The cone generated for this redundant inequalities is called truncated quadratic module generated by the polynomials p_1, \dots, p_m , as we see in Definition 3.1. This relaxations give us an increasing sequence of lower bounds of the infimum P^* , as you can see in 3.9. Lasserre proved that this sequence converge asymptotically to the infimum if we assume some arquimedean property in the cone generated for the redundant inequalities, see [21, Theorem 5] for a proof.

Definition 3.1. Let $p_1, \dots, p_m \in \mathbb{R}[\underline{X}]$ and $k \in \mathbb{N}_0 \cup \{\infty\}$. We define the k -truncated quadratic module M , generated by p_1, \dots, p_m as:

$$(15) \quad \begin{aligned} M_k(p_1, \dots, p_m) := & \mathbb{R}[\underline{X}]_k \cap \sum \mathbb{R}[\underline{X}]^2 + \mathbb{R}[\underline{X}]_k \cap \sum \mathbb{R}[\underline{X}]^2 p_1 \\ & + \dots + \mathbb{R}[\underline{X}]_k \cap \sum \mathbb{R}[\underline{X}]^2 p_m \subseteq \mathbb{R}[\underline{X}]_k \end{aligned}$$

where here $\mathbb{R}[\underline{X}]_\infty := \mathbb{R}[\underline{X}]$. We use the notation $M(p_1, \dots, p_m) := M_\infty(p_1, \dots, p_m)$, to refer to the quadratic module generated by the polynomials $p_1, \dots, p_m \in \mathbb{R}[\underline{X}]$.

Remark 3.2. Note that:

$$\mathbb{R}[\underline{X}]_k \cap \sum \mathbb{R}[\underline{X}]^2 p = \left\{ \sum_{i=1}^l h_i^2 p \mid h_i \in \mathbb{R}[\underline{X}], 2 \deg(h_i) \leq k - \deg(p) \right\}$$

For a proof this see [21, Page 5].

Lemma 3.3. Let $k \in \mathbb{N}$, $p \in \mathbb{R}[\underline{X}]_k \setminus \{0\}$ and $d := \lfloor \frac{k - \deg(p)}{2} \rfloor$. Let $L \in \mathbb{R}[\underline{X}]_k^*$. Then it holds:

$$(16) \quad L(\sum \mathbb{R}[\underline{X}]_k \cap \mathbb{R}[\underline{X}]^2 p) \subseteq \mathbb{R}_{\geq 0} \iff M_{k,p}(y) \succeq 0$$

Proof. Let us set the matrices $A_\alpha \in \mathbb{R}^{s_d \times s_d}$ for $|\alpha| \leq k$, as the matrices such that:

$$pV_d V_d^T = \sum_{|\alpha| \leq k} \underline{X}^\alpha A_\alpha \in \mathbb{R}[\underline{X}]_k^{s_d \times s_d}.$$

and $y_\alpha := L(\underline{X}^\alpha)$ for $|\alpha| \leq k$.

$$\begin{aligned} L(\sum \mathbb{R}[\underline{X}]_k \cap \mathbb{R}[\underline{X}]^2 p) \subseteq \mathbb{R}_{\geq 0} &\stackrel{3.2}{\iff} \forall h \in \mathbb{R}[\underline{X}]_d, L(h^2 p) \geq 0 \\ &\iff \forall H \in \mathbb{R}^{s_d}, L((H^T V_d)(V_d^T H)p) \geq 0 \\ &\iff \forall H \in \mathbb{R}^{s_d}, L(H^T p V_d V_d^T H) \geq 0 \\ &\iff \forall H \in \mathbb{R}^{s_d}, L(H^T (\sum_{|\alpha| \leq k} \underline{X}^\alpha A_\alpha) H) \geq 0 \\ &\iff \forall H \in \mathbb{R}^{s_d}, L(\sum_{|\alpha| \leq k} \underline{X}^\alpha H^T A_\alpha H) \geq 0 \\ &\stackrel{L \text{ is linear}}{\iff} \forall H \in \mathbb{R}^{s_d}, \sum_{|\alpha| \leq k} L(\underline{X}^\alpha) H^T A_\alpha H \geq 0 \\ &\iff \forall H \in \mathbb{R}^{s_d}, H^T (\sum_{\alpha} y_\alpha A_\alpha) H \geq 0 \\ &\iff \sum_{|\alpha| \leq k} y_\alpha A_\alpha \succeq 0 \iff M_{k,p}(y) \succeq 0 \end{aligned}$$

□

Due to Lemma 3.3 the following definition of moment relaxation using linear forms is equivalent to the definition given in 2.6

Definition 3.4. Let (P) be a polynomial optimization problem given as in 2 and let $k \in \mathbb{N}_0 \cup \{\infty\}$ such that $f, p_1, \dots, p_m \in \mathbb{R}[\underline{X}]_k$. The moment relaxation (or Lasserre relaxation) of (P) of degree k is the semidefinite optimization problem:

$$(P_k) \quad \begin{array}{lll} \text{minimize} & L(f) & \text{subject to} \\ & L \in \mathbb{R}[\underline{X}]_k^* & \\ & L(1) = 1 & \\ & L(M_k(p_1, \dots, p_m)) \subseteq \mathbb{R}_{\geq 0} & \end{array}$$

the optimal value of (P_k) i.e., the infimum over all $L(f)$ where L ranges over all optimal solutions of (P_k) is denoted by $P_k^* \in \{-\infty\} \cup \mathbb{R} \cup \{\infty\}$.

Corollary and Notation 3.5. Let $d \in \mathbb{N}_0$. The correspondence:

$$\begin{array}{l} L \mapsto (L(\underline{X}^{\alpha+\beta}))_{|\alpha|, |\beta| \leq d} \\ \left(\begin{array}{ll} \mathbb{R}[\underline{X}]_{2d} & \rightarrow \mathbb{R} \\ \underline{X}^\alpha & \mapsto y_\alpha \end{array} \right) \leftarrow M_d(y) \end{array}$$

defines a bijection between the linear forms $L \in \mathbb{R}[\underline{X}]_{2d}^*$ such that $L(\sum \mathbb{R}[\underline{X}]_d^2) \subseteq \mathbb{R}_{\geq 0}$ and the set of positive semidefinite generalized Hankel matrices of order d i.e. $H_d \cap \mathbb{R}_{\geq 0}^{s_d \times s_d}$. Let $L \in \mathbb{R}[\underline{X}]_{2d}^*$ such that $L(\sum \mathbb{R}[\underline{X}]_d^2) \subseteq \mathbb{R}_{\geq 0}$ we denote $M_L := (L(\underline{X}^{\alpha+\beta}))_{|\alpha|, |\beta| \leq d}$ and let $M_d(y) \succeq 0$ for $y \in \mathbb{R}^{s_d}$ we denote:

$$L_{M_d(y)} : \mathbb{R}[\underline{X}]_{2d} \longrightarrow \mathbb{R}, \underline{X}^\alpha \mapsto y_\alpha.$$

Proof. The well-definedness of both maps follows from Lemma 3.3. Now, let $L \in \mathbb{R}[\underline{X}]_{2d}^*$ such that $L(\sum \mathbb{R}[\underline{X}]_d^2) \subseteq \mathbb{R}_{\geq 0}$ then:

$$L_{M_L} : \mathbb{R}[\underline{X}]_{2d} \longrightarrow \mathbb{R}, \underline{X}^\alpha \mapsto L(\underline{X}^\alpha)$$

since notice that $M_L = (L(\underline{X}^{\alpha+\beta}))_{|\alpha|,|\beta|\leq d} = M_d(L(0), \dots, L(X_n^{2d}))$. Hence $L_{M_L} = L$. On the other side, let $M_d(y) \succeq 0$ for $y \in \mathbb{R}^{s_d}$ then:

$$M_{L_{M_d(y)}} = (L_{M_d(y)}(\underline{X}^{\alpha+\beta}))_{|\alpha|,|\beta|\leq d} = (y_{\alpha+\beta})_{|\alpha|,|\beta|\leq d} = M_d(y).$$

□

Notation 3.6. We denote the following isomorphism of vector spaces by:

$$\text{poly} : \mathbb{R}^{s_d} \longrightarrow \mathbb{R}[\underline{X}]_d, a \mapsto a^T V_d$$

Proposition 3.7. Let $d \in \mathbb{N}_0$ and $L \in \mathbb{R}[\underline{X}]_{2d}^*$ then:

$$L(pq) = P^T M_L Q$$

where $P := \text{poly}^{-1}(p)$ and $Q := \text{poly}^{-1}(q)$.

Proof. As usual let us set the matrices $A_\alpha \in \mathbb{R}^{s_d \times s_d}$ for $|\alpha| \leq 2d$ as the matrices such that:

$$V_d V_d^T = \sum_{|\alpha| \leq k} \underline{X}^\alpha A_\alpha \in \mathbb{R}[\underline{X}]_k^{s_d \times s_d}.$$

Then:

$$\begin{aligned} L(pq) &= L(P^T V_d V_d^T Q) = L(P^T \sum_{|\alpha| \leq 2d} \underline{X}^\alpha A_\alpha Q) = \\ &= \sum_{|\alpha| \leq 2d} L(\underline{X}^\alpha) P^T A_\alpha Q = P^T \left(\sum_{|\alpha| \leq 2d} L(\underline{X}^\alpha) A_\alpha \right) Q = P^T M_L Q \end{aligned}$$

□

Definition 3.8. Let $L \in \mathbb{R}[\underline{X}]_d^*$. A quadrature rule for L on $U \subseteq \mathbb{R}[\underline{X}]_d$ is a function $w : N \rightarrow \mathbb{R}_{>0}$ defined on a finite set $N \subseteq \mathbb{R}^n$, such that:

$$(17) \quad L(p) = \sum_{x \in N} w(x) p(x)$$

for all $p \in U$. A quadrature rule for L is a quadrature for L on $\mathbb{R}[\underline{X}]_d$. We call the elements of N the nodes of the quadrature rule.

Proposition 3.9. Let (P) be the polynomial optimization problem given in (2) with $f, p_1, \dots, p_m \in \mathbb{R}[\underline{X}]_k$. Then the following holds:

- (i) $P^* \geq P_\infty^* \geq \dots \geq P_{k+1}^* \geq P_k^*$.
- (ii) Let $L \in \mathbb{R}[\underline{X}]_k^*$ with $L(1) = 1$. Suppose L has a quadrature rule with nodes in S , then L is a feasible solution of (P_k) with $L(f) \geq P_k^*$.
- (iii) Suppose (P_k) has an optimal solution L^* , which has a quadrature rule on $\mathbb{R}[\underline{X}]_l$ for some $l \in \{1, \dots, k\}$ with $f \in \mathbb{R}[\underline{X}]_l$ and the nodes are in S . Then $L^*(f) = P_k^*$, moreover we have $P_k^* = P_{k+m}^*$ for $m \geq 0$ and the nodes of the quadrature rule are global minimizers of (P) .
- (iv) In the situation of (iii), suppose moreover that (P) has an unique global minimizer x^* , then $L^*(f) = f(x^*)$ and $x^* = (L^*(X_1), \dots, L^*(X_n))$.

Proof. (i) $P^* \geq P_\infty^*$ since if x is a feasible solution for (P) then $\text{ev}_x \in \mathbb{R}[\underline{X}]^*$ is a feasible solution for P_∞ with the same value, that is $f(x) = \text{ev}_x(f)$. It remains to prove $P_l^* \geq P_k^*$ for $l \in \mathbb{N}_{\geq k} \cup \{\infty\}$. For this let L be a feasible solution of (P_l) , as $M_k(p_1, \dots, p_m) \subseteq M_l(p_1, \dots, p_m)$ then $L|_{\mathbb{R}[\underline{X}]_k}$ is a feasible solution of (P_k) with the same optimal value.

(ii) Suppose L has a quadrature rule with nodes $a_1, \dots, a_N \in S$ and weights $\lambda_1 > 0, \dots, \lambda_N > 0$. From $L(1) = 1$ we get $\sum_{i=1}^N \lambda_i = 1$ and since the nodes are in S it holds $L(M_k(p_1, \dots, p_m)) \subseteq \mathbb{R}_{\geq 0}$. Hence L is a feasible solution of (P_k) . Moreover the following holds:

$$P_k^* = L(1)P_k^* = \sum_{i=1}^N \lambda_i P_k^* \leq \sum_{i=1}^N \lambda_i f(a_i) = L(f)$$

where the inequality follows from the fact that $P_k^* \leq f(x)$ for all $x \in S$.

(iii) Suppose L^* is an optimal solution of (P_k) then $L^*(f) = P_k^* \leq P^*$ using (i) and on

other side since $L^*(1) = 1$ and L^* has a quadrature rule on $\mathbb{R}[\underline{X}]_l$ with nodes in S and $f \in \mathbb{R}[\underline{X}]_l$, there exist $a_1, \dots, a_N \in S$ nodes, and $\lambda_1 > 0, \dots, \lambda_N > 0$ weights, such that:

$$(18) \quad P_k^* = L^*(f) = \sum_{i=1}^N \lambda_i f(a_i) \geq \sum_{i=1}^N \lambda_i P^* = P^*$$

Therefore $L^*(f) = P^*$, and since $P_k^* = P^*$ we get equality everywhere in (i) and we can conclude that $P^* = P_{k+m}^*$ for $m \geq 0$. It remains to show that the nodes are global minimizers of (P) , but this is true since in (18) we have equality everywhere, and if we factor out we get $\sum_{i=1}^N \lambda_i (f(a_i) - P^*) = 0$, as $\lambda_i > 0$ and $f(a_i) - P^* \geq 0$ for all $i \in \{1, \dots, N\}$, implying $f(a_i) = P^*$ for all $i \in \{1, \dots, N\}$.

(iv) Using (iii) we have that $L^*(f) = P^* = f(x^*)$, and continuing with the same notation as in the proof of (iii) we got by unicity of the minimizer x^* , that $a_i = x^*$ for all $i \in \{1, \dots, N\}$. This implies that $L^* = \text{ev}_{x^*}$ on $\mathbb{R}[\underline{X}]_l^*$, and evaluating in the polynomials $X_1, \dots, X_N \in \mathbb{R}[\underline{X}]_1$ we got that:

$$L^*(X_i) = \text{ev}_{x^*}(X_i) = x_i^* \text{ for all } i \in \{1, \dots, N\}.$$

That is to say, $x^* = (L^*(X_1), \dots, L^*(X_n))$. \square

We can now reformulate our problem as:

Given $d \in \mathbb{N}_0$ and $L \in \mathbb{R}[\underline{X}]_{2d+2}^*$ such that $L(\sum \mathbb{R}[\underline{X}]_{d+1}^2) \subseteq \mathbb{R}_{\geq 0}$, we would like to obtain for all $p \in \mathbb{R}[\underline{X}]_{2d+2}$:

- Nodes $x_1, \dots, x_r \in \mathbb{R}^n$ and weights $\lambda_1, \dots, \lambda_r > 0$ such that:

$$L(p) = \sum_{i=1}^r \lambda_i p(x_i)$$

in other words:

- $x_{1,1}, \dots, x_{1,n}, \dots, x_{r,1}, \dots, x_{r,n} \in \mathbb{R}$ and $a_1, \dots, a_r \in \mathbb{R}$ such that:

$$L(p) = \sum_{i=1}^r a_i^2 p(x_{i,1}, \dots, x_{i,n})$$

again with other words:

- $x_{1,1}, \dots, x_{1,n}, \dots, x_{r,1}, \dots, x_{r,n} \in \mathbb{R}$ and $a_1, \dots, a_r \in \mathbb{R}$ such that:

$$L(p) = \left\langle \begin{pmatrix} p(x_{1,1}, \dots, x_{1,n}) & & \\ & \ddots & \\ & & p(x_{r,1}, \dots, x_{r,n}) \end{pmatrix} \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix}, \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix} \right\rangle$$

again written differently:

- $x_{1,1}, \dots, x_{1,n}, \dots, x_{r,1}, \dots, x_{r,n} \in \mathbb{R}$ and $a \in \mathbb{R}^r$ such that:

$$L(p) = \left\langle p \left[\begin{pmatrix} x_{1,1} & & \\ & \ddots & \\ & & x_{r,1} \end{pmatrix}, \dots, \begin{pmatrix} x_{1,n} & & \\ & \ddots & \\ & & x_{r,n} \end{pmatrix} \right] \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix}, \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix} \right\rangle$$

again with less words:

- Diagonal matrices $D_1, \dots, D_n \in \mathbb{R}^{r \times r}$ and $a \in \mathbb{R}^n$ such that:

$$L(p) = \langle p(D_1, \dots, D_n)a, a \rangle$$

Reminder 3.10. Let $r, n \in \mathbb{N}$ and $M_1, \dots, M_n \in \mathbb{R}^{r \times r}$ symmetric commuting matrices. Then there exist an orthogonal matrix $P \in \mathbb{R}^{r \times r}$ such that $P^t M_i P$ is a diagonal matrix for all $i \in \{1, \dots, n\}$.

Using this theorem we can continue with our reformulation of the problem: given $d \in \mathbb{N}_0$ and $L \in \mathbb{R}[\underline{X}]_{2d+2}^*$ such that $L(\sum \mathbb{R}[\underline{X}]_{d+1}^2) \subseteq \mathbb{R}_{\geq 0}$, to find a quadrature rule for L is the

same as to find commuting symmetric matrices $M_1, \dots, M_n \in \mathbb{R}^{r \times r}$ and a vector $a \in \mathbb{R}^r$ such that:

$$(19) \quad L(p) = \langle p(M_1, \dots, M_n)a, a \rangle$$

We end the reformulation of the problem once and for all with the languages of endomorphisms, instead of matrices. That is to say: given $d \in \mathbb{N}_0$ and $L \in \mathbb{R}[\underline{X}]_{2d+2}^*$ such that $L(\sum \mathbb{R}[\underline{X}]_{d+1}^2) \subseteq \mathbb{R}_{\geq 0}$, we would like to obtain a finite dimensional euclidean vector space V , commuting self-adjoint endomorphisms M_1, \dots, M_n of V and $a \in V$ such that:

$$(20) \quad L(p) = \langle p(M_1, \dots, M_n)a, a \rangle$$

Remark 3.11. Gelfand, Naimark and Segal gave a solution for the case we allow to the space V to be infinite dimensional and the linear form to be strictly positive in the sums of squares, that is to say, in the case we are given a linear form $L \in \mathbb{R}[\underline{X}]^*$ such that $L(p^2) > 0$ for all $p \neq 0$. The solution was given by defining the inner product:

$$(21) \quad \langle p, q \rangle := L(pq)$$

and defining the self adjoint operators M_i , for all $i \in \{1, \dots, n\}$, on the infinite dimensional vector space $\mathbb{R}[\underline{X}]$, in the following way:

$$M_i : \mathbb{R}[\underline{X}] \longrightarrow \mathbb{R}[\underline{X}], p \mapsto X_i p$$

Taking $a := 1 \in \mathbb{R}[\underline{X}]$ we have the searched equality (20).

From now on we will assume we are given a linear form $L \in \mathbb{R}[\underline{X}]_{2d+2}^*$ for $d \in \mathbb{N}_0 \cup \{\infty\}$ such that $L(\sum \mathbb{R}[\underline{X}]_{d+1}^2) \subseteq \mathbb{R}_{\geq 0}$ or what is the same due to 3.5 and 3.3 M_L is positive semidefinite, unless L is defined explicitly in other way.

4. TRUNCATED GNS-CONSTRUCTION

In this section we will explain how we can define the euclidean vector space and multiplication operators required in (20) from this positive semidefinite linear form L , in a similar way as in the Gelfand-Neimark-Segal construction 3.11.

First, we will get rid of the problem that, $L(p^2) = 0$ does not imply $p = 0$ for every $p \in \mathbb{R}[\underline{X}]_{d+1}$, that is to say (21) does not define an inner product if the linear form is positive semidefinite. By grouping together the polynomials with this property we will be able to define an inner product, on a quotient space. As a consequence, we will obtain an euclidean vector space. With respect to the multiplication operators, we will need to do the orthogonal projection on the class of polynomials with one degree less, in such a way that when we do the multiplication for the variable X_i we are not out of our ambient space. This construction was already done in [18].

Definition and Notation 4.1. We define and denote the truncated GNS kernel of L :

$$U_L := \{p \in \mathbb{R}[\underline{X}]_{d+1} \mid L(pq) = 0 \text{ for all } q \in \mathbb{R}[\underline{X}]_{d+1}\}$$

Proposition 4.2. The truncated GNS kernel of L is a vector subspace in $\mathbb{R}[\underline{X}]_{d+1}$. Moreover:

$$(22) \quad U_L = \{p \in \mathbb{R}[\underline{X}]_{d+1} \mid L(p^2) = 0\}$$

Proof. The fact that U_L is a vector subspace follows directly from the linearity of L . Let us prove the equality (22). For this let us denote $A := \{p \in \mathbb{R}[\underline{X}]_{d+1} \mid L(p^2) = 0\}$. The inclusion $U_L \subseteq A$ is trivial. For the other inclusion we will demonstrate first, due to L is positive semidefinite and linear, that the Cauchy-Schwarz inequality holds:

$$(23) \quad L(pq)^2 \leq L(p^2)L(q^2)$$

Indeed, for all $t \in \mathbb{R}$ and $p, q \in \mathbb{R}[\underline{X}]_{d+1}$ it holds:

$$0 \leq L((p + tq)^2) = L(p^2) + 2tL(pq) + t^2L(q^2)$$

Therefore the polynomial $r := L(p^2) + 2XL(pq) + X^2L(q^2) \in \mathbb{R}[X]_2$ is non negative, i.e. $r(x) \geq 0$ for all $x \in \mathbb{R}$. In the case $L(q^2) \neq 0$, the discriminant of r has to be less or equal to zero i.e. $4L(pq)^2 - 4L(p^2)L(q^2) \leq 0$ and we get the searched inequality (23). In the case $L(q^2) = 0$, then $L(pq) = 0$ and trivially we get also the inequality (23). As a consequence

if $p \in A$ then $L(p^2) = 0$, and this implies due to (23), $L(pq) = 0$ for all $q \in \mathbb{R}[\underline{X}]_d$, and therefore $p \in U_L$. \square

Definition and Notation 4.3. We define and denote the GNS representation space of L , as the following quotient of vector spaces:

$$(24) \quad V_L := \frac{\mathbb{R}[\underline{X}]_{d+1}}{U_L}$$

For every $p \in \mathbb{R}[\underline{X}]_{d+1}$ we will write \bar{p}^L to refer us to the class of p in V_L . We define and denote the GNS inner product of L , in the following way:

$$(25) \quad \langle \bar{p}^L, \bar{q}^L \rangle_L := L(pq)$$

for every $p, q \in \mathbb{R}[\underline{X}]_{d+1}$.

Proposition 4.4. $(V_L, \langle \cdot, \cdot \rangle_L)$, is a symmetric bilinear form:

Proof. Let us prove first that $\langle \cdot, \cdot \rangle_L$ is well defined. To do this take $p_1, q_1, p_2, q_2 \in \mathbb{R}[\underline{X}]_{d+1}$ with $\bar{p}_1^L = \bar{p}_2^L$ and $\bar{q}_1^L = \bar{q}_2^L$ then:

$$\begin{aligned} \langle \bar{p}_1^L, \bar{q}_1^L \rangle_L = \langle \bar{p}_2^L, \bar{q}_2^L \rangle_L &\iff L(p_1q_1) = L(p_2q_2) \iff L(p_1q_1) - L(p_2q_2) = 0 \\ &\iff L(p_1q_1) + L(-p_2q_1) - L(-p_2q_1) - L(p_2q_2) = 0 \\ &\iff L((p_1 - p_2)q_1) - L(p_2(q_2 - q_1)) = 0 \end{aligned}$$

The last equality holds since $p_1 - p_2, q_2 - q_1 \in U_L$. The bilinearity and symmetry is trivial. $\langle \cdot, \cdot \rangle_L$ is positive semidefinite since $L(\sum \mathbb{R}[\underline{X}]_{d+1}^2) \subseteq \mathbb{R}_{\geq 0}$. It remains to prove that $\langle \cdot, \cdot \rangle_L$ is even positive definite. Indeed, for all $p \in \mathbb{R}[\underline{X}]_{d+1}$ with $\langle \bar{p}^L, \bar{p}^L \rangle_L = 0$ then $L(p^2) = 0$ and then $p \in U_L$ as we have shown in 22. \square

Definition and Notation 4.5. For $i \in \{1, \dots, n\}$, we define the i -th truncated GNS multiplication operator of L as the following map between euclidean vector subspaces of V_L , and denote by $M_{L,i}$:

$$(26) \quad M_{L,i} : \Pi(V_L) \longrightarrow \Pi_L(V_L), \bar{p}^L \mapsto \Pi(\overline{pX_i^L}) \text{ for } p \in \mathbb{R}[\underline{X}]_d$$

where Π_L is the orthogonal projection map of V_L into the vector subspace $\{\bar{p}^L \mid p \in \mathbb{R}[\underline{X}]_d\}$ with respect to the inner product $\langle \cdot, \cdot \rangle_L$. We will call and denote the subvector space:

$$(27) \quad T_L := \Pi(V_L) = \{\bar{p}^L \mid p \in \mathbb{R}[\underline{X}]_d\}$$

of V_L , the GNS-truncation of L .

Proposition 4.6. The i -th truncated GNS multiplication operator of L is a self-adjoint endomorphism of T_L .

Proof. Let us demonstrate first that the i -th truncated GNS multiplication operator of L is well defined. $M_{L,i}$ is well defined if and only if $M_{L,i}(\bar{p}^L) = \bar{0}^L$ for all $p \in U_L \cap \mathbb{R}[\underline{X}]_d$ if and only if $\Pi_L(\overline{X_i p^L}) = \bar{0}^L$ for all $p \in U_L \cap \mathbb{R}[\underline{X}]_d$. Since $\Pi_L(\overline{X_i p^L}) \in T_L$ we can choose $q \in \mathbb{R}[\underline{X}]_d$ such that $\bar{q}^L = \Pi_L(\overline{X_i p^L})$ and then:

$$\begin{aligned} L(q^2) &= \langle \bar{q}, \bar{q} \rangle_L = \langle \Pi_L(\overline{X_i p^L}), \Pi_L(\overline{X_i p^L}) \rangle_L \stackrel{\Pi_L \circ \Pi_L = \Pi_L}{=} \langle \Pi_L(\overline{X_i p^L}), \overline{X_i p^L} \rangle_L \\ &= \langle \bar{q}^L, \overline{X_i p^L} \rangle_L = L(q(X_i p)) = L((qX_i)p) \stackrel{p \in U_L}{=} 0 \end{aligned}$$

Therefore $\Pi_L(\overline{X_i p^L}) = \bar{0}^L$ for all $p \in U_L$. Let us see now that $M_{L,i}$ are self-adjoint endomorphisms, for this let $p, q \in \mathbb{R}[\underline{X}]_d$ then:

$$\begin{aligned} \langle M_{L,i}(\bar{p}^L), \bar{q}^L \rangle_L &= \langle \Pi_L(\overline{X_i p^L}), \bar{q}^L \rangle_L = \langle \overline{X_i p^L}, \Pi_L(\bar{q}^L) \rangle_L = \langle \overline{X_i p^L}, \bar{q}^L \rangle_L = L((X_i p)q) \\ &= L(p(X_i q)) = \langle \bar{p}^L, \overline{X_i q^L} \rangle_L = \langle \Pi_L(\bar{p}^L), \overline{X_i q^L} \rangle_L = \langle \bar{p}^L, \Pi_L(\overline{X_i q^L}) \rangle_L = \langle \bar{p}^L, M_{L,i}(\bar{q}^L) \rangle_L \end{aligned}$$

\square

Remark 4.7. The GNS construction for $L \in \mathbb{R}[\underline{X}]^*$ with $L(\sum \mathbb{R}[\underline{X}]^2) \subseteq \mathbb{R}_{\geq 0}$ is the same as the original (3.11) modulo U_L . The GNS representation space of L and the GNS truncation of L are the same $\frac{\mathbb{R}[\underline{X}]}{U_L}$, where:

$$(28) \quad U_L = \{p \in \mathbb{R}[\underline{X}] \mid L(p^2) \geq 0\}$$

The truncated GNS multiplication operators of L commute, since $\frac{\mathbb{R}[\underline{X}]}{U_L}$ is a commutative ring. One can easily prove that U_L is an ideal. Indeed it is clear that if $p, q \in U_L$ then $L((p+q)^2) = 0$, and if $p \in U_L$ and $q \in \mathbb{R}[\underline{X}]$ then $L(p^2q^2) = L(p(pq^2)) \stackrel{p \in U_L}{=} 0$ implies $pq \in U_L$.

Lemma and Notation 4.8. Remember that $L' := L_{\mathbb{R}[\underline{X}]_{2d+1}}$. Let us denote as B_L the transformation matrix of the following bilinear form with respect to the standard monomial basis:

$$\mathbb{R}[\underline{X}]_{d+1} \times \mathbb{R}[\underline{X}]_d \longrightarrow \mathbb{R}, (p, q) \longmapsto L(pq)$$

Then it holds $\text{rank } M_{L'} = \text{rank } B_L$ and for every such L linear form we can define its respective modified moment matrix as:

$$\widetilde{M}_L := \left(\begin{array}{c|c} M_{L'} & M_{L'}W_L \\ \hline W_L^T M_{L'} & W_L^T M_{L'}W_L \end{array} \right)$$

where W_L is a matrix such that $M_{L'}W_L = C_L$, where C_L is the submatrix of B_L remaining from eliminating the columns corresponding to the matrix $M_{L'}$. \widetilde{M}_L is well defined since it does not depend from the election of W_L and it is positive semidefinite.

Proof. Notice that $M_{L'}$ is the transformation matrix of the linear map:

$$\varphi := \mathbb{R}[\underline{X}]_d \longrightarrow \mathbb{R}[\underline{X}]_d^*, p \mapsto (a \mapsto L(pq))$$

with respect to the standard monomial basis and in the same way B_L is the transformation matrix of the linear map:

$$\psi := \mathbb{R}[\underline{X}]_{d+1} \longrightarrow \mathbb{R}[\underline{X}]_d^*, p \mapsto (q \mapsto L(pq))$$

with respect to the standard monomial basis. Note that to prove $\text{rank } M_{L'} = \text{rank } B_L$ it is the same than to prove $\varphi(\mathbb{R}[\underline{X}]_d) = \psi(\mathbb{R}[\underline{X}]_{d+1})$. It is obvious that $\varphi(\mathbb{R}[\underline{X}]_d) \subseteq \psi(\mathbb{R}[\underline{X}]_{d+1})$. For the other inclusion we take $\Lambda \in \psi(\mathbb{R}[\underline{X}]_{d+1})$ then there exists $p \in \mathbb{R}[\underline{X}]_{d+1}$ such that $\Lambda(q) = \psi(p)(q) = L(pq)$ for all $q \in \mathbb{R}[\underline{X}]_d$. We look for a $g \in \mathbb{R}[\underline{X}]_d$ such that $\Lambda(q) = L(gq)$ for all $q \in \mathbb{R}[\underline{X}]_d$, because then $\varphi(h)(q) = L(gq) = \Lambda(q)$ for all $q \in \mathbb{R}[\underline{X}]_d$ implying $\varphi(g) = \Lambda$ and then we could conclude $\Lambda \in \varphi(\mathbb{R}[\underline{X}]_d)$. In other words, we want to show that there exists $g \in \mathbb{R}[\underline{X}]_d$ such that $L(pq) = L(gq)$ for every $q \in \mathbb{R}[\underline{X}]_d$. With more different words, our aim is to find $g \in \mathbb{R}[\underline{X}]_d$ such that:

$$\langle \bar{p}, \bar{q} \rangle_L = \langle \bar{g}, \bar{q} \rangle_L \text{ for every } q \in \mathbb{R}[\underline{X}]_d$$

For this we define the following linear form:

$$\Lambda_p := \frac{\mathbb{R}[\underline{X}]_d}{U_L \cap \mathbb{R}[\underline{X}]_d} \longrightarrow \mathbb{R}, \bar{q} \mapsto L(pq) \text{ for every } q \in \mathbb{R}[\underline{X}]_d$$

$\Lambda_p \in \left(\frac{\mathbb{R}[\underline{X}]_d}{U_L \cap \mathbb{R}[\underline{X}]_d}\right)^*$. Since $\frac{\mathbb{R}[\underline{X}]_d}{U_L}$ is a finite dimensional euclidean vector space is in particular a Hilbert space and then by the Fréchet-Riesz Representation Theorem there exists $\bar{g} \in \frac{\mathbb{R}[\underline{X}]_d}{U_L \cap \mathbb{R}[\underline{X}]_d}$ with $g \in \mathbb{R}[\underline{X}]_d$, such that:

$$\Lambda_p(\bar{q}) = \langle \bar{q}, \bar{g} \rangle_L \text{ for every } q \in \mathbb{R}[\underline{X}]_d.$$

Therefore $L(pq) = \Lambda_p(\bar{q}) = \langle \bar{q}, \bar{g} \rangle_L = L(gq)$ for every $q \in \mathbb{R}[\underline{X}]_d$. Then $\text{rank } M_{L'} = \text{rank } B_L$ and therefore there exists W_L (may not be unique) such that $M_{L'}W_L = C_L$. Now, we claim that the modified moment matrix \widetilde{M}_L , does not depend from the choice of the matrix W_L with the property $M_{L'}W = C_L$. Indeed, assume there are matrices W_1, W_2 such that $M_{L'}W_1 = M_{L'}W_2$. Let us denote:

$$W_1 := (P_1, \dots, P_{r_{d+1}}), \quad W_2 := (Q_1, \dots, Q_{r_{d+1}})$$

where $P_1, \dots, P_{r_{d+1}}$ and $Q_1, \dots, Q_{r_{d+1}}$ are the respective column vectors of the matrices W_1 and W_2 and define $p_i := \text{poly}(Q_i)$ and $q_i := \text{poly}(Q_i)$ for $i \in \{1, \dots, r_{d+1}\}$. Then we have the following matrix equality:

$$M_{L'}(P_1, \dots, P_{r_{d+1}}) = M_{L'}(Q_1, \dots, Q_{r_{d+1}})$$

Let $i \in \{1, \dots, r_{d+1}\}$ then:

$$M_{L'}P_i = M_{L'}Q_i \iff M_{L'}(P_i - Q_i) = 0 \iff (P_i - Q_i)^T M_{L'}(P_i - Q_i) \stackrel{3.7}{\underset{(22)}{\iff}} p_i - q_i \in U_{L'}$$

This implies $L'(p_i p_j) = L'(q_i q_j)$ and again due to 3.7 we get that:

$$P_i^T M_{L'} P_j = Q_i^T M_{L'} Q_j \text{ for all } i, j \in \{1, \dots, r\}$$

Then we have got that:

$$\begin{aligned} W_1^T M_{L'} W_1 &= \begin{pmatrix} P_1^T M_{L'} P_1 & \dots & P_1^T M_{L'} P_r \\ \vdots & \ddots & \vdots \\ P_r^T M_{L'} P_1 & \dots & P_r^T M_{L'} P_r \end{pmatrix} = \\ &= \begin{pmatrix} Q_1^T M_{L'} Q_1 & \dots & Q_1^T M_{L'} Q_r \\ \vdots & \ddots & \vdots \\ Q_r^T M_{L'} Q_1 & \dots & Q_r^T M_{L'} Q_r \end{pmatrix} = W_2^T M_{L'} W_2 \end{aligned}$$

Therefore $W_1^T M_{L'} W_1 = W_2^T M_{L'} W_2$, and we can conclude that \widetilde{M}_L is well defined. Moreover since $M_{L'}$ is a positive semidefinite matrix, then there exists a matrix $C \in \mathbb{R}^{s_d \times s_d}$ such that $M_{L'} = CC^T$ due to 2.5. Then we have the following factorization:

$$\widetilde{M}_L = \left(\begin{array}{c|c} CC^T & CC^T W \\ \hline W^T CC^T & W^T CC^T W \end{array} \right) = \left(\begin{array}{c|c} C & 0 \\ \hline W^T C & 0 \end{array} \right) \left(\begin{array}{c|c} C & 0 \\ \hline W^T C & 0 \end{array} \right)^T$$

then taking $P := \left(\begin{array}{c|c} C & 0 \\ \hline W^T C & 0 \end{array} \right)$ we get that $\widetilde{M}_L = PP^T$, which due to 2.5, proves \widetilde{M}_L is positive semidefinite. \square

5. GAUSSIAN QUADRATURE RULE

In this section we will prove the existence of a quadrature rule representation for the positive semidefinite linear form L on a set that contains $\mathbb{R}[\underline{X}]_{2d+1}$ by providing that the truncated GNS multiplication operators commute. We will also demonstrate that this condition is strictly more general than the very well known condition of being flat, condition that for its part ensure the existence of a quadrature rule representation for L on the whole space in contrast with the quadrature rule in a space that contains $\mathbb{R}[\underline{X}]_{2d+1}$ that we get in case the truncated GNS multiplication operators commute.

Proposition 5.1. The vector spaces T_L and $\frac{\mathbb{R}[\underline{X}]_d}{U_L \cap \mathbb{R}[\underline{X}]_d}$ are canonically isomorphic.

Proof. Let us consider the following linear map between euclidean vector spaces:

$$(29) \quad \sigma_L : T_L \longrightarrow V_{L'} : \bar{p}^L \longmapsto \bar{p}^{L'} \text{ for every } p \in \mathbb{R}[\underline{X}]_d$$

where remember we denoted $L' := L|_{\mathbb{R}[\underline{X}]_{2d}}$. It is well defined since for every $\bar{p}^L, \bar{q}^L \in T_L$ such that $\bar{p}^L = \bar{q}^L$ we can assume without loss of generality that $p, q \in \mathbb{R}[\underline{X}]_d$, and therefore:

$$\bar{p}^L = \bar{q}^L \iff L((p - q)^2) = 0 \iff L'((p - q)^2) = 0 \iff \bar{p}^{L'} = \bar{q}^{L'} \iff \sigma_0(\bar{p}^L) = \sigma_0(\bar{q}^L)$$

σ_0 is also a linear isometry, since for every $p, q \in \mathbb{R}[\underline{X}]_d$ we have:

$$\langle \bar{p}^L, \bar{q}^L \rangle_L = L(pq) = L'(pq) = \langle \bar{p}^{L'}, \bar{q}^{L'} \rangle_{L'} = \langle \sigma_L(\bar{p}^L), \sigma_L(\bar{q}^L) \rangle_{L'}$$

Then σ_L is immediately injective. On other side, σ_L is surjective since for every $\bar{p}^{L'} \in V_{L'}$ with $p \in \mathbb{R}[\underline{X}]_d$, it holds that $\sigma_L(\bar{p}^L) = \bar{p}^{L'}$. Thence σ_L is an isomorphism between vector spaces. \square

Notation 5.2. For a linear form $\ell \in \mathbb{R}[\underline{X}]_{2d}^*$ such that $\ell(\sum \mathbb{R}[\underline{X}]_{d+1}^2) \subseteq \mathbb{R}_{\geq 0}$ we will denote by σ_ℓ the following isomorphism of euclidean vector spaces already defined in (29):

$$(30) \quad \sigma_\ell : T_\ell \mapsto V_{\ell'}, \bar{p}^\ell \mapsto \bar{p}^{\ell'}, \text{ for } p \in \mathbb{R}[\underline{X}]_d$$

Remark 5.3. For $v_1, \dots, v_r \in \mathbb{R}[\underline{X}]_d$, we have $\bar{v}_1^L, \dots, \bar{v}_r^L$ is an orthonormal basis of T_L if and only if $\bar{v}_1^{L'}, \dots, \bar{v}_r^{L'}$ is an orthonormal basis of $V_{L'}$.

The following Theorem and Lemma, are probably very well known and we will use them to prove Proposition (5.8). The proofs can be seen for example in [2] and [13].

Theorem 5.4. And ideal $I \subseteq \mathbb{R}[\underline{X}]$ is zero dimensional (i.e. $|V_{\mathbb{C}}(I)| < \infty$) if and only if the vector space $\mathbb{R}[\underline{X}]/I$ is finite dimensional. Moreover $|V_{\mathbb{C}}(I)| \leq \dim(\mathbb{R}[\underline{X}]/I)$, with equality if and only if the ideal I is radical.

Proof. Theorem 2.6 page 15 in [13]. \square

Definition 5.5. Let $I \subseteq \mathbb{R}[\underline{X}]$ be an ideal. I is said to be radical when $I = \mathcal{I}(V_{\mathbb{C}}(I))$.

Lemma 5.6. Let $I \subseteq \mathbb{R}[\underline{X}]$ be an ideal. I is radical if and only if

$$(31) \quad \text{For all } g \in \mathbb{R}[\underline{X}] \text{ such that } g^2 \in I \implies g \in I$$

Proof. There is a proof in [13, Lemma 2.2]. \square

Proposition 5.7. Let $\Lambda \in \mathbb{R}[\underline{X}]^*$ such that $\Lambda(\sum \mathbb{R}[\underline{X}]^2) \subseteq \mathbb{R}_{\geq 0}$. Then U_Λ is a radical ideal.

Proof. In 4.7 we saw that U_Λ is an ideal, let us prove that it is real radical ideal. Let $g \in \mathbb{R}[\underline{X}]$ such that $g^2 \in U_\Lambda$. In particular $\Lambda(g^2 1) = 0$ and this implies $g \in U_\Lambda$. \square

Proposition 5.8. Let $\Lambda = \sum_{i=1}^N \lambda_i \text{ev}_{a_i} \in \mathbb{R}[\underline{X}]^*$, with $N \in \mathbb{N}$, $\lambda_1 > 0, \dots, \lambda_N > 0$, and $a_1, \dots, a_N \in \mathbb{R}^n$ then:

$$\dim\left(\frac{\mathbb{R}[\underline{X}]}{U_\Lambda}\right) = |\{a_1, \dots, a_N\}|$$

Proof. We have the following equalities:

$$\begin{aligned} U_\Lambda &= \{p \in \mathbb{R}[\underline{X}] \mid \sum_{i=1}^N \lambda_i p^2(a_i) = 0\} = \{p \in \mathbb{R}[\underline{X}] \mid p^2(a_i) = 0 \text{ for all } i \in \{1, \dots, n\}\} \\ &= \{p \in \mathbb{R}[\underline{X}] \mid p(a_i) = 0 \text{ for all } i \in \{1, \dots, n\}\} = I(\{a_1, \dots, a_N\}) \end{aligned}$$

and since $\{a_1, \dots, a_N\} \subseteq \mathbb{R}^n$ is an algebraic set, by the ideal-variety correspondence (see [2]), it holds:

$$V_{\mathbb{C}}(I(\{a_1, \dots, a_N\})) = \{a_1, \dots, a_N\}$$

what is the same as $V_{\mathbb{C}}(U_\Lambda) = \{a_1, \dots, a_N\}$. Notice that by Theorem (5.4) is enough to prove that U_Λ is radical to finish the proof. In fact by Proposition 5.7 U_Λ is radical. Applying Theorem 5.4 we have the result. \square

Let us review some known bounds on the number of nodes of quadrature rules for L on $\mathbb{R}[\underline{X}]_{2d+2}$ and on $\mathbb{R}[\underline{X}]_{2d+1}$ (see [4] and [18]).

Proposition 5.9. Then number of nodes N , of a quadrature rule for L satisfies:

$$\text{rank } M_L \leq N \leq |V_{\mathbb{C}}(U_L)|$$

Proof. Let $L = \sum_{i=1}^N \lambda_i \text{ev}_{a_i} \in \mathbb{R}[\underline{X}]_{2d+2}^*$ for $a_1, \dots, a_N \in \mathbb{R}^n$ pairwise different points and $\lambda_1, \dots, \lambda_N > 0$ weights and define $\Lambda := \sum_{i=1}^N \lambda_i \text{ev}_{a_i} \in \mathbb{R}[\underline{X}]^*$. Let us consider the following canonical map:

$$\frac{\mathbb{R}[\underline{X}]_{d+1}}{U_L} \hookrightarrow \frac{\mathbb{R}[\underline{X}]}{U_\Lambda}$$

By Proposition 5.8 we have that:

$$\text{rank } M_L = \dim\left(\frac{\mathbb{R}[\underline{X}]_{d+1}}{U_L}\right) \leq \dim\left(\frac{\mathbb{R}[\underline{X}]}{U_\Lambda}\right) = N$$

On the other side, it holds that $\{a_1, \dots, a_N\} \subseteq V_C(U_L)$, since for all $p \in U_L$ we have $L(p^2) = 0$ and then $p(a_i) = 0$ for all $i \in \{1, \dots, N\}$. this implies $N \leq |V_C(U_L)|$. \square

Proposition 5.10. The number of nodes N , of a quadrature rule for L on $\mathbb{R}[\underline{X}]_{2d+1}$ satisfies:

$$N \geq \dim(T_L)$$

Proof. Assume that L has a quadrature rule on $\mathbb{R}[\underline{X}]_{2d+1}$ such that:

$$L(p) = \sum_{i=1}^N \lambda_i p(a_i)$$

for every $p \in \mathbb{R}[\underline{X}]_{2d+1}$, where we can assume without loss of generality that the points $a_1, \dots, a_N \in \mathbb{R}^n$ are pairwise different and $\lambda_1, \dots, \lambda_N > 0$ with $N < \infty$ for $N \in \mathbb{N}$. Let us set $\Lambda := \sum_{i=1}^N \lambda_i \text{ev}_{a_i} \in \mathbb{R}[\underline{X}]^*$. Then, the following linear map between euclidean vector spaces is an isometry:

$$(32) \quad \sigma_1 : T_L \longrightarrow \frac{\mathbb{R}[\underline{X}]}{U_\Lambda}, \bar{p}^L \mapsto \bar{p}^\Lambda \text{ for every } p \in \mathbb{R}[\underline{X}]_d$$

It is easy to see that is well defined since $U_L \subseteq U_\Lambda$. It holds also that σ_1 is a linear isometry since, for all $p, q \in \mathbb{R}[\underline{X}]_d$:

$$\langle \bar{p}^L, \bar{q}^L \rangle_L = L(pq) = \Lambda(pq) = \langle \bar{p}^\Lambda, \bar{q}^\Lambda \rangle_\Lambda = \langle \sigma_1(\bar{p}^L), \sigma_1(\bar{q}^L) \rangle_\Lambda$$

Since, σ_1 is a linear isometry is immediately injective, and then:

$$\dim(T_L) \leq \dim\left(\frac{\mathbb{R}[\underline{X}]}{U_\Lambda}\right)$$

And now we can apply the Proposition 5.8, to conclude the proof. \square

Definition 5.11. A quadrature rule for L on $\mathbb{R}[\underline{X}]_{2d+1}$ with minimal number of nodes, that is to say with $\dim(T_L)$ nodes is called a Gaussian quadrature rule.

Lemma 5.12. Assume that the truncated multiplication operators commute. Then for all $p \in \mathbb{R}[\underline{X}]_{d+1}$ we have the following equality:

$$(33) \quad p(M_{L,1}, \dots, M_{L,n})(\bar{1}^L) = \Pi_L(\bar{p}^L)$$

Proof. Let $p = \underline{X}^\alpha$ for $\alpha \in \mathbb{N}^n$ with $|\alpha| \leq d+1$. We continue the proof by induction on $|\alpha|$:

- For $|\alpha| = 0$, we have that $\underline{X}^\alpha = 1$ then:

$$1(M_{L,1}, \dots, M_{L,n})(\bar{1}^L) = \text{Id}_{V_L}(\bar{1}^L) = \bar{1}^L = \Pi_L(\bar{1}^L)$$

- Let assume the statement is true for $|\alpha| = d$. Let us show it is also true for $|\alpha| = d+1$. Let $p = X_i q$ for some $i \in \{1, \dots, n\}$ and $q = \underline{X}^\beta$ with $|\beta| = d$, then $\Pi_L(\bar{q}^L) = \bar{q}^L$ since $\bar{q}^L \in T_L$, and we have:

$$\begin{aligned} p(M_{L,1}, \dots, M_{L,n})(\bar{1}^L) &= (M_{L,i} \circ q(M_{L,1}, \dots, M_{L,n}))(\bar{1}^L) = \\ &M_{L,i}(q(M_{L,1}, \dots, M_{L,n})(\bar{1}^L)) = M_{L,i}(\bar{q}^L) = \Pi_L(\bar{X}_i \bar{q}^L) = \Pi_L(\bar{p}^L) \end{aligned}$$

since we have proved (33) for monomials then by the linearity of the orthogonal projection (33) is also true for polynomials. \square

Theorem 5.13. Assume the truncated multiplication operators of L commute, and consider the set:

$$(34) \quad G_L := \left\{ \sum_{i=1}^s p_i q_i \mid s \in \mathbb{N}, p_i \in \mathbb{R}[\underline{X}]_{d+1} \text{ and } q_i \in \mathbb{R}[\underline{X}]_d + U_L \right\}$$

then there exists a quadrature rule for L on G_L with $\dim(T_L)$ many nodes.

Proof. Since the truncated multiplication operators of L commute by the Remainder 3.10 there exists an orthonormal basis $v := \{v_1, \dots, v_N\}$ of T_L consisting of common eigenvectors of the GNS truncated multiplication operators of L . That is to say, there exist $a_1, \dots, a_N \in \mathbb{R}^n$ such that:

$$M_{L,i} v_j = a_{j,i} v_j \text{ for all } i \in \{1, \dots, n\} \text{ and } j \in \{1, \dots, N\}$$

where $N := \dim(T_L)$. Since it always holds $\bar{1}^L \in T_L$ since $d \in \mathbb{N}$, we can write:

$$(35) \quad \bar{1}^L = b_1 v_1 + \cdots + b_N v_N$$

for some $b_1, \dots, b_N \in \mathbb{R}$. Let us define $\lambda_i := b_i^2$ for all $i \in \{1, \dots, N\}$. Let $g = pq$ such that $p \in \mathbb{R}[\underline{X}]_{d+1}$ and $q \in \mathbb{R}[\underline{X}]_d + U_L$, then using Lemma 5.12 we have the two equalities:

$$(36) \quad \Pi_L(\bar{p}^L) = p(M_{L,1}, \dots, M_{L,n})(\bar{1}^L) \text{ and } \bar{q}^L = q(M_{L,1}, \dots, M_{L,n})(\bar{1}^L)$$

Using this equalities (36), using that the orthogonal projection Π_L is selfadjoint, using that $\{v_1, \dots, v_N\}$ is an orthonormal basis of T_L consisting of common eigenvectors of the GNS truncated multiplication operators of L and also using the equation (35), with the same idea as we got the reformulation of the problem in (20) we have:

$$\begin{aligned} L(g) &= L(pq) = \langle \bar{p}^L, \bar{q}^L \rangle_L \stackrel{\bar{q}^L \in T_L}{=} \langle \bar{p}^L, \Pi_L(\bar{q}^L) \rangle_L = \\ &= \langle \Pi_L(\bar{p}^L), \bar{q}^L \rangle_L = \sum_{j=1}^N b_j^2 p(a_j) q(a_j) = \sum_{j=1}^N \lambda_j p(a_j) q(a_j) \end{aligned}$$

Then by linearity it holds that $L(p) = \sum_{i=1}^N \lambda_i p(a_i)$ for all $p \in G_L$. It remains to prove that the nodes of the quadrature rule for L that we got, $a_1, \dots, a_N \in \mathbb{R}^n$ are pairwise different, but this is true since $N = \dim T_L$ is the minimal possible number of nodes for a quadrature rule on $\mathbb{R}[\underline{X}]_{2d+1}$ as we proved in 5.10. \square

Remark 5.14. Since $\mathbb{R}[\underline{X}]_{2d+1} \subseteq G_L$, in the conditions of Theorem 5.13 we got in particular a Gaussian quadrature rule for the linear form L .

Corollary 5.15. Let $n = 1$, i.e. $L \in \mathbb{R}[X]_{2d+2}^*$ with $L(\sum \mathbb{R}[X]^2) \geq 0$. Then L has a quadrature rule on G_L (34).

Proof. L has one truncated GNS multiplication operator, therefore the hypothesis of Theorem 5.13 holds and there is a quadrature rule on G_L for L . \square

Proposition 5.16. The following assertions are equivalent:

- (i) $\mathbb{R}[\underline{X}]_{d+1} = \mathbb{R}[\underline{X}]_d + U_L$
- (ii) $T_L = V_L$
- (iii) For all $\alpha \in \mathbb{N}_0^n$ with $|\alpha| = d+1$, there exists $p \in \mathbb{R}[\underline{X}]_d$ such that $\underline{X}^\alpha - p \in U_L$
- (iv) The canonical map:

$$(37) \quad V_{L'} = \mathbb{R}[\underline{X}]_d / U_{L'} \hookrightarrow \mathbb{R}[\underline{X}]_{d+1} / U_L = V_L$$

is an isomorphism.

- (v) $\dim(V_{L'}) = \dim(V_L)$
- (vi) The moment matrices $(L(\underline{X}^{\alpha+\beta}))_{|\alpha|, |\beta| \leq d}$ and $(L(\underline{X}^{\alpha+\beta}))_{|\alpha|, |\beta| \leq d+1}$ have the same rank.
- (vii) $M_L = \widetilde{M}_L$.

Proof. Note that the map (37) it is well defined since $\mathbb{R}[\underline{X}]_d \cap U_L = U_{L'}$. And one can see immediately that:

$$(i) \iff (ii) \iff (iii) \iff (iv) \iff (v).$$

Let us show $(v) \iff (vi)$: $(L(\underline{X}^{\alpha+\beta}))_{|\alpha|, |\beta| \leq d+1}$ is the transformation matrix (or the associated matrix) of the bilinear form:

$$\mathbb{R}[\underline{X}]_{d+1} \times \mathbb{R}[\underline{X}]_{d+1} \longrightarrow \mathbb{R}, (p, q) \mapsto L(pq)$$

with respect to the the standard monomial basis, and therefore it is also the transformation matrix (or the associated matrix) of the linear map:

$$(38) \quad \mathbb{R}[\underline{X}]_{d+1} \longrightarrow \mathbb{R}[\underline{X}]_{d+1}^*, p \mapsto (q \mapsto L(pq))$$

with respect to the corresponding dual basis of the standard monomial basis. The kernel of this linear map (38) is U_L , in consequence:

$$\text{rank}((L(\underline{X}^{\alpha+\beta}))_{|\alpha|,|\beta|\leq d+1}) = \dim \mathbb{R}[\underline{X}]_{d+1} - U_L = \dim V_L$$

reasoning in the same way:

$$\text{rank}((L(\underline{X}^{\alpha+\beta}))_{|\alpha|,|\beta|\leq d}) = \dim V_{L'}$$

Finally (vi) \iff (vii):

$$\begin{aligned} & \text{rank}((L(\underline{X}^{\alpha+\beta}))_{|\alpha|,|\beta|\leq d}) \iff \text{rank}(M_{L'}) = \text{rank}(M_L) \\ & \iff \left(\begin{array}{c|c} M_{L'} & M_{L'}W_L \\ \hline W_L^T M_{L'} & W_L^T M_{L'}W_L \end{array} \right) = \left(\begin{array}{c|c} M_{L'} & M_{L'}W_L \\ \hline W_L^T M_{L'} & C_L \end{array} \right) \\ & \iff W_L M_{L'} W_L = C_L \iff \widetilde{M}_L = M_L \end{aligned}$$

□

Definition 5.17. We say the linear form L is flat if the conditions (i), (ii), (iii), (iv), (v), (vi) and (vii) in (5.16) are satisfied.

Proposition 5.18. Suppose L is flat then the truncated GNS operators of L commute

Proof. Assume L is flat, and let $i, j \in \{1, \dots, n\}$ and $p \in \mathbb{R}[\underline{X}]_d$. We want to prove:

$$M_{L,i} \circ M_{L,j}(\overline{p}^L) = M_{L,j} \circ M_{L,i}(\overline{p}^L)$$

Let us write $X_i p = p_1 + q_1$ and $X_j p = p_2 + q_2$ with $p_1, p_2 \in \mathbb{R}[\underline{X}]_d$ and $q_1, q_2 \in U_L$. Then $M_{L,j}(\overline{p}^L) = \Pi_L(\overline{X_j p}^L) = \Pi_L(\overline{p_2 + q_2}^L) = \Pi_L(\overline{p_2}^L) + \Pi_L(\overline{q_2}^L) = \Pi_L(\overline{p_2}^L) = \overline{p_2}^L$. In the same way we get $M_{L,i}(\overline{p}^L) = \overline{p_1}^L$. Therefore:

$$\begin{aligned} M_{L,i} \circ M_{L,j}(\overline{p}^L) &= M_{L,j} \circ M_{L,i}(\overline{p}^L) \iff M_{L,i}(\overline{p_2}^L) = M_{L,j}(\overline{p_1}^L) \\ &\iff \Pi_L(\overline{X_i p_2}^L) = \Pi_L(\overline{X_j p_1}^L) \end{aligned}$$

In other words, define $\overline{g}^L := \Pi_L(X_i p_2 - X_j p_1) \in T_L$ for some $g \in \mathbb{R}[\underline{X}]_d$, then it is enough to show $g \in U_L$. Indeed:

$$\begin{aligned} L(g^2) &= \langle \Pi_L(\overline{X_i p_2 - X_j p_1}), \overline{g}^L \rangle_L = \langle \overline{X_i p_2 - X_j p_1}, \Pi_L(\overline{g}^L) \rangle_L = \langle \overline{X_i p_2 - X_j p_1}, \overline{g} \rangle_L \\ &= L((X_i p_2 - X_j p_1)g) = L((X_i g)p_2) - L((X_j g)p_1) = \langle \overline{X_i g}^L, \overline{p_2}^L \rangle_L - \langle \overline{X_j g}^L, \overline{p_1}^L \rangle_L \\ &= \frac{\overline{X_j p_2}^L = \overline{p_2}^L}{\overline{X_i p_1}^L = \overline{p_1}^L} \langle \overline{X_i g}^L, \overline{X_j p_1}^L \rangle_L - \langle \overline{X_j g}^L, \overline{X_i p_1}^L \rangle_L = L(X_i g X_j p) - L(X_j g X_i p) = 0 \end{aligned}$$

□

Here we show some examples which shows that the reverse of Proposition 5.18 does not hold.

Example 5.19. The truncated GNS multiplication operators of the following linear form:

$$L : \mathbb{R}[X_1, X_2]_4 \rightarrow \mathbb{R}, p \mapsto \frac{1}{4}(p(0, 0) + p(1, 0) + p(-1, 0) + p(0, 1))$$

commute but L is not flat. Indeed, if we do the truncated GNS-construction we have:

$$M_L = \begin{array}{c} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \end{array} \left(\begin{array}{ccc|cc} 1 & X_1 & X_2 & X_1^2 & X_1 X_2 & X_2^2 \\ \hline 1 & 0 & \frac{1}{4} & \frac{1}{2} & 0 & \frac{1}{4} \\ 0 & \frac{1}{2} & 0 & 0 & 0 & 0 \\ \frac{1}{4} & 0 & \frac{1}{4} & 0 & 0 & \frac{1}{4} \\ \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ \frac{1}{4} & 0 & \frac{1}{4} & 0 & 0 & \frac{1}{4} \end{array} \right) = \left(\begin{array}{c|c} A_L & B_L \\ \hline B_L^T & C_L \end{array} \right)$$

is the associated moment matrix of the linear form L and a basis of the truncated GNS-kernel of L is $\langle X_1 X_2, X_2^2 - X_2 \rangle$. That is, the rank of M_L is 4. And since in the kernel there is no polynomials of degree less or equal to 1, we get that the unique element in the kernel of L' is 0, then the truncated GNS space is $\frac{\mathbb{R}[X_1, X_2]_1}{U_{L'}} \cong \mathbb{R}[X_1, X_2]_1$, which implies the dimension of the GNS-truncated space is 3 and therefore L is not flat by (vi)

in 5.16. We can also verify that L is not flat by computing \widetilde{M}_L . Indeed, in this case A_M is invertible and W_M is uniquely defined by $W_M = A_M^{-1}B_M$, then \widetilde{M}_L reads:

$$\widetilde{M}_L = \left(\begin{array}{ccc|ccc} 1 & 0 & \frac{1}{4} & \frac{1}{2} & 0 & \frac{1}{4} \\ 0 & \frac{1}{2} & 0 & 0 & 0 & 0 \\ \frac{1}{4} & 0 & \frac{1}{4} & 0 & 0 & \frac{1}{4} \\ \hline \frac{1}{2} & 0 & 0 & \frac{1}{3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ \frac{1}{4} & 0 & \frac{1}{4} & 0 & 0 & \frac{1}{4} \end{array} \right)$$

Since $\widetilde{M}_L \neq M_L$ then L is not flat by (vii) in 5.16.

Let us compute the truncated GNS multiplication operators of L . First note that:

$$T_L \cong \frac{\mathbb{R}[X_1, X_2]_1}{U_{L'}} = \langle \overline{1}^{L'}, \overline{X_1}^{-L'}, \overline{X_2}^{-L'} \rangle$$

Therefore by Remark 5.3 the truncated GNS space of L is:

$$T_L = \langle \overline{1}^L, \overline{X_1}^L, \overline{X_2}^L \rangle$$

With the Gram-Schmidt orthonormalization process we get the following orthonormal basis with respect to the GNS product of L :

$$\underline{v} := \{ \overline{1}^L, \overline{\sqrt{2}X_1}^L, -\frac{\sqrt{3}}{3} + \frac{4\sqrt{3}}{3}\overline{X_2}^L \}$$

The matrices of the GNS-multiplication operators with respect to this orthonormal basis are:

$$A_1 := M(M_{L, X_1}, \underline{v}) = \begin{pmatrix} 0 & \frac{\sqrt{2}}{2} & 0 \\ \frac{\sqrt{2}}{2} & 0 & -\frac{\sqrt{6}}{6} \\ 0 & -\frac{\sqrt{6}}{6} & 0 \end{pmatrix}$$

$$A_2 := M(M_{L, X_2}, \underline{v}) = \begin{pmatrix} \frac{1}{4} & 0 & \frac{\sqrt{3}}{4} \\ 0 & 0 & 0 \\ \frac{\sqrt{3}}{4} & 0 & \frac{3}{4} \end{pmatrix}$$

It is easy to check that the truncated GNS multiplication operators of L commute, that is $M_{L, X_1} \circ M_{L, X_2} - M_{L, X_2} \circ M_{L, X_1} = 0$. Now since M_{L, X_1} and M_{L, X_2} commute we can do the simultaneous diagonalization on both of them, in order to find an orthonormal basis of the GNS truncation of L consisting of common eigenvectors of M_{L, X_1} and M_{L, X_2} . To do this we follow the same idea as in [15, Algorithm 4.1, Step 1] and compute for a matrix:

$$A = r_1 A_1 + r_2 A_2 \text{ where } r_1^2 + r_2^2 = 1$$

a matrix P orthogonal such that $P^T A P$ is a diagonal matrix. In this case, we get for:

$$P = \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{6}}{4} & -\frac{\sqrt{6}}{4} \\ 0 & \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{3}}{2} & \frac{\sqrt{2}}{4} & \frac{\sqrt{2}}{4} \end{pmatrix}$$

$$P^T A_1 P = \begin{pmatrix} 0 & 0 & 0 \\ 0 & -\frac{\sqrt{6}}{3} & 0 \\ 0 & 0 & \frac{\sqrt{6}}{3} \end{pmatrix} \text{ and } P^T A_2 P = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Looking over the proof of 5.13 we can obtain the weights $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}_{>0}$ through the following operations:

$$P^T \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \\ -\frac{\sqrt{6}}{4} \\ -\frac{\sqrt{6}}{4} \end{pmatrix}$$

then $\lambda_1 = (\frac{1}{2})^2$ and $\lambda_2 = \lambda_3 = (-\frac{\sqrt{6}}{4})^2$. Therefore we get the following decomposition:

$$\widetilde{M}_L = \frac{1}{4}V_2(0, 1)V_2(0, 1)^T + \frac{3}{8}V_2(-\frac{\sqrt{6}}{3}, 0)V_2(-\frac{\sqrt{6}}{3}, 0)^T + \frac{3}{8}V_2(\frac{\sqrt{6}}{3}, 0)V_2(\frac{\sqrt{6}}{3}, 0)^T.$$

Example 5.21. Let us consider the following generalized Hankel matrix in two variables of order 2 taken from [6, Example 1.13]:

$$M = \begin{array}{c} \\ \\ \\ \\ \\ \\ \\ \end{array} \begin{array}{c} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \end{array} \left(\begin{array}{ccc|ccc} 1 & X_1 & X_2 & X_1^2 & X_1 X_2 & X_2^2 \\ 1 & 1 & 2 & 2 & 0 & 3 \\ 1 & 2 & 0 & 4 & 0 & 0 \\ 1 & 0 & 3 & 0 & 0 & 9 \\ \hline 2 & 4 & 0 & 9 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 3 & 0 & 9 & 0 & 0 & 28 \end{array} \right)$$

This matrix does not have a quadrature rule representation with the minimal number of nodes, as has been proved in [6, Example 1.13], however it admits a Gaussian quadrature rule. Indeed, we can compute with the truncated GNS construction that:

$$\widetilde{M} = \frac{1}{6}V_2(0,0)V_2(0,0)^T + \frac{1}{3}V_2(0,3)V_2(0,3)^T + \frac{1}{2}V_2(2,0)V_2(2,0)^T.$$

The following corollary is a very well known result of Curto and Fialkow (see [4, corollary 5.14]) in terms of quadrature rules instead of nonnegative measures. In [1] there is a proof about the correspondence between quadrature rules and nonnegative measures. This result of Curto and Fialkow uses tools of functional analysis like the Spectral theorem and the Riesz representation theorem. Monique Laurent gave also a more elementary proof (see [12, corollary 1.4]) that uses a corollary of the Hilbert Nullstellensatz and elementary linear algebra. The main contribution of this proof is that it does not need to find a flat extension of the linear form since the truncated GNS multiplication operators commute and we can apply directly the Theorem 3.10, and despite of it uses the Hilbert Nullstellensatz in the proof of Theorem 5.4, we do not need to apply the Hilbert Nullstellensatz to show that the nodes are in \mathbb{R}^n , since the nodes are real because its coordinates are the eigenvalues of a real symmetric matrix.

Corollary 5.22. Suppose L is flat then L has a quadrature rule with $\text{rank}(M_L)$ many nodes (the minimal number of nodes).

Proof. If L is flat by Proposition 5.18 the truncated GNS multiplication operators of L commute and applying 5.13 then L has a quadrature rule on G_L (34), with $\dim(T_L) \stackrel{L \text{ is flat}}{=} \dim(V_L) = \text{rank}(M_L)$ many nodes. Since L is flat $\mathbb{R}[\underline{X}]_{d+1} = \mathbb{R}[\underline{X}]_d + U_L$ and therefore one can easily see that $G_L = \mathbb{R}[\underline{X}]_{2d+2}$. As a conclusion we get a quadrature rule for L with $\text{rank}(M_L)$ many nodes. \square

6. MAIN THEOREM

In this section we will demonstrate that the commutativity of the truncated GNS multiplication operators of L is equivalent to the matrix $W_L^T A_L W_L$ being Hankel.

Main Theorem 6.1. The following assertions are equivalent:

- (1) The truncated multiplication operators $M_{L,1}, \dots, M_{L,n}$ pairwise commute.
- (2) There exists $\hat{L} \in \mathbb{R}[\underline{X}]_{2d+2}^*$ such that $L = \hat{L}$ on $\mathbb{R}[\underline{X}]_{2d+1}$ and \hat{L} is flat.

Proof. 1 \Rightarrow 2. By the theorem 5.13 there exist $a_1, \dots, a_N \in \mathbb{R}^n$ pairwise different nodes and $\lambda_1 > 0, \dots, \lambda_N > 0$ weights, where $N := \dim(T_L)$ such that: $L(p) = \sum_{i=1}^N \lambda_i p(a_i)$ for all $p \in G_L$, where G_L was defined in (34). Let us define, $\hat{L} := \sum_{i=1}^N \lambda_i \text{ev}_{a_i} \in \mathbb{R}[\underline{X}]_{2d+2}^*$. We have shown in theorem 5.13 that $\hat{L} = L$ on $\mathbb{R}[\underline{X}]_{2d+1}$, $U_L \subset U_{\hat{L}}$ and obviously $\tilde{L}(\sum \mathbb{R}[\underline{X}]_{d+1}^2) \subseteq \mathbb{R}_{\geq 0}$, so it remains to show that \tilde{L} is flat, that is to say:

$$\dim V_{\hat{L}} = \dim T_{\hat{L}}$$

or equivalently using 5.1, it remains to show:

$$\dim\left(\frac{\mathbb{R}[\underline{X}]_{d+1}}{U_{\hat{L}}}\right) = \dim\left(\frac{\mathbb{R}[\underline{X}]_d}{U_{\hat{L}} \cap \mathbb{R}[\underline{X}]_d}\right)$$

Since $U_{\hat{L}} \cap \mathbb{R}[\underline{X}]_d = U_L \cap \mathbb{R}[\underline{X}]_d$ and using again proposition 5.1, we have the following:

$$\dim\left(\frac{\mathbb{R}[\underline{X}]_d}{U_{\hat{L}} \cap \mathbb{R}[\underline{X}]_d}\right) = \dim\left(\frac{\mathbb{R}[\underline{X}]_d}{U_L \cap \mathbb{R}[\underline{X}]_d}\right) \stackrel{5.1}{=} \dim(T_L) = N$$

therefore, in the following we will prove $\dim\left(\frac{\mathbb{R}[\underline{X}]_{d+1}}{U_{\hat{L}}}\right) = N$. For this, let us consider the following linear map, between euclidean vector spaces:

$$(40) \quad \frac{\mathbb{R}[\underline{X}]_{d+1}}{U_{\hat{L}}} \hookrightarrow \frac{\mathbb{R}[\underline{X}]}{U_{\Lambda}}, \bar{p}^{\hat{L}} \mapsto \bar{p}^{\Lambda}$$

where $\Lambda := \sum_{i=1}^N \lambda_i \text{ev}_{a_i} \in \mathbb{R}[\underline{X}]^*$. Notice that the canonical map (40) is well defined since $U_{\hat{L}} = U_{\Lambda} \cap \mathbb{R}[\underline{X}]_d$ and therefore it is injective. Then

$$\dim\left(\frac{\mathbb{R}[\underline{X}]_{d+1}}{U_{\hat{L}}}\right) \leq \dim\left(\frac{\mathbb{R}[\underline{X}]}{U_{\Lambda}}\right) \stackrel{5.8}{=} N$$

It remains to show $N \leq \dim\left(\frac{\mathbb{R}[\underline{X}]_{d+1}}{U_{\hat{L}}}\right)$. But this is true, since:

$$N = \dim(T_L) \stackrel{5.1}{=} \dim\left(\frac{\mathbb{R}[\underline{X}]_d}{U_L \cap \mathbb{R}[\underline{X}]_d}\right) = \dim\left(\frac{\mathbb{R}[\underline{X}]_d}{U_{\hat{L}} \cap \mathbb{R}[\underline{X}]_d}\right) \leq \dim\left(\frac{\mathbb{R}[\underline{X}]_{d+1}}{U_{\hat{L}}}\right)$$

2 \Rightarrow 1. Since \hat{L} is flat, then by 5.18 we know that the truncated GNS multiplication operators of \hat{L} pairwise commute. Then by applying again 5.13 there exists $a_1, \dots, a_N \in \mathbb{R}^n$, pairwise different nodes, and $\lambda_1 > 0, \dots, \lambda_N > 0$ weights, with $N = \dim(T_{\hat{L}})$ such that if we set $\Lambda := \sum_{i=1}^N \lambda_i \text{ev}_{a_i} \in \mathbb{R}[\underline{X}]^*$, we get $\Lambda(p) = \hat{L}(p) = L(p)$ for all $p \in \mathbb{R}[\underline{X}]_{2d+1}$, and $U_L \subseteq U_{\hat{L}} \subseteq U_{\Lambda}$. Indeed notice that $U_L \subseteq U_{\hat{L}}$ since for $p \in U_L$, $L(pq) = \hat{L}(pq) = 0$ for all $q \in \mathbb{R}[\underline{X}]_d$, and since \hat{L} is flat this implies $p \in U_{\hat{L}}$. Obviously $M_{\Lambda, i}$ pairwise commute for all $i \in \{1, \dots, n\}$, since they are the original GNS operators modulo U_{Λ} defined in 4.7. In order to prove that $M_{L, i}$ pairwise commute for all $i \in \{1, \dots, n\}$, let us first consider the linear isometry σ_1 (32) of the proposition 5.10. Since σ_1 is a linear isometry is immediately injective, and then $\dim(T_L) \leq \dim\left(\frac{\mathbb{R}[\underline{X}]}{U_{\Lambda}}\right)$. Therefore we have the following inequalities:

$$N = \dim(T_{\hat{L}}) = \dim\left(\frac{\mathbb{R}[\underline{X}]_d}{U_{\hat{L}} \cap \mathbb{R}[\underline{X}]_d}\right) = \dim\left(\frac{\mathbb{R}[\underline{X}]_d}{U_L \cap \mathbb{R}[\underline{X}]_d}\right) = \dim(T_L) \leq \dim\left(\frac{\mathbb{R}[\underline{X}]}{U_{\Lambda}}\right) \stackrel{5.8}{=} N$$

then $\dim(T_L) = \dim\left(\frac{\mathbb{R}[\underline{X}]}{U_{\Lambda}}\right)$. Then σ_1 , in this case, is in particular surjective and in conclusion is an isomorphism. With this result we be able to prove that the following diagram is commutative, for all $i \in \{1, \dots, n\}$:

$$(41) \quad \begin{array}{ccc} T_L & \xrightarrow{M_{L, i}} & T_L \\ \downarrow \sigma_1 & & \uparrow \sigma_1^{-1} \\ \frac{\mathbb{R}[\underline{X}]}{U_{\Lambda}} & \xrightarrow{M_{\Lambda, i}} & \frac{\mathbb{R}[\underline{X}]}{U_{\Lambda}} \end{array}$$

That is to say $M_{L, i} = \sigma_1^{-1} \circ M_{\Lambda, i} \circ \sigma_1$. To show this let $p, q \in \mathbb{R}[\underline{X}]_d$, then we have:

$$\begin{aligned} \langle M_{L, i}(\bar{p}^L), \bar{q}^L \rangle_L &= \langle \Pi_L(\overline{X_i p^L}), \bar{q}^L \rangle_L \stackrel{\Pi_L \circ \Pi_{\underline{L}} = \Pi_L}{=} \langle \overline{X_i p^L}, \bar{q}^L \rangle_L = L(X_i p q) \stackrel{\Lambda = L \text{ on } \mathbb{R}[\underline{X}]_{2d+1}}{=} \\ \Lambda(X_i p q) &= \langle \overline{X_i p^{\Lambda}}, \bar{q}^{\Lambda} \rangle_{\Lambda} = \langle \sigma_1 \circ \sigma_1^{-1}(\overline{X_i p^{\Lambda}}), \bar{q}^{\Lambda} \rangle_{\Lambda} = \langle \sigma_1^{-1}(\overline{X_i p^{\Lambda}}), \sigma_1^{-1}(\bar{q}^{\Lambda}) \rangle_L = \\ &= \langle \sigma_1^{-1} \circ M_{\Lambda, i}(\bar{p}^{\Lambda}), \bar{q}^{\Lambda} \rangle_L = \langle \sigma_1^{-1} \circ M_{\Lambda, i} \circ \sigma_1(\bar{p}^L), \bar{q}^L \rangle_L \end{aligned}$$

Finally we can conclude that the truncated GNS multiplication operators of L pairwise commute, using the commutativity of the GNS multiplication operators of Λ . Indeed:

$$(42) \quad \begin{aligned} M_{L,i} \circ M_{L,j} &= \sigma_1^{-1} \circ M_{\Lambda,i} \circ \sigma_1 \circ \sigma_1^{-1} \circ M_{\Lambda,j} \circ \sigma_1 = \sigma_1^{-1} \circ M_{\Lambda,i} \circ M_{\Lambda,j} \circ \sigma_1 = \\ &= \sigma_1^{-1} \circ M_{\Lambda,j} \circ M_{\Lambda,i} \circ \sigma_1 = \sigma_1^{-1} \circ M_{\Lambda,j} \circ \sigma_1 \circ \sigma_1^{-1} \circ M_{\Lambda,i} \circ \sigma_1 = M_{L,j} \circ M_{L,i} \end{aligned}$$

□

Theorem 6.2. The following assertions are equivalent:

- (1) $M_{L,1}, \dots, M_{L,n}$ pairwise commute.
- (2) \widetilde{M}_L is a Generalized Hankel matrix

Proof. (1) \Rightarrow (2) Assume $M_{L,1}, \dots, M_{L,n}$ pairwise commute. Then by Theorem 6.1 there exists a linear form $\hat{L} \in \mathbb{R}[\underline{X}]_{2d+2}^*$ such that: $\hat{L}(\mathbb{R}[\underline{X}]_{d+1}^2) \subseteq \mathbb{R}_{\geq 0}$, $L = \hat{L}$ on $\mathbb{R}[\underline{X}]_{2d+1}$ and \hat{L} is flat, what implies by 5.14 that \hat{L} has a quadrature rule representation and therefore $M_{\hat{L}} \succeq 0$ and $M_{\hat{L}}$ is a Generalized Hankel matrix. It is enough to show that $C_{\hat{L}} = W_L^T A_L W_L$. The last follows from the fact that \hat{L} is flat and $L = \hat{L}$ on $\mathbb{R}[\underline{X}]_{2d+1}$ since:

$$\text{rank}(M_{\hat{L}}) = \text{rank} \left(\begin{array}{c|c} A_L & A_L W_L \\ \hline W_L^T A_L & C_{\hat{L}} \end{array} \right) \stackrel{\hat{L} \text{ is flat}}{=} \text{rank}(A) \iff C_{\hat{L}} = W_L^T A_L W_L$$

(2) \Rightarrow (1) Suppose \widetilde{M}_L is a Generalized Hankel matrix, and denote $\hat{L} := L_{\widetilde{M}_L} \in \mathbb{R}[\underline{X}]_{2d+2}^*$. Since \widetilde{M}_L is the moment matrix of the linear form $\hat{L} \in \mathbb{R}[\underline{X}]_{2d+2}^*$ then \hat{L} is flat and by Theorem 5.18 the truncated GNS multiplication operators of \hat{L} commute. Now, to prove the truncated GNS operators commute let us define $\sigma := \sigma_{\hat{L}}^{-1} \circ \sigma_L$ an isomorphism of euclidean vector spaces. We will prove that the following diagram is commutative:

$$\begin{array}{ccc} T_L & \xrightarrow{M_{L,i}} & T_L \\ \downarrow \sigma & & \uparrow \sigma^{-1} \\ T_{\hat{L}} & \xrightarrow{M_{\hat{L},i}} & T_{\hat{L}} \end{array}$$

The diagram is commutative if and only if $M_{L,i} = \sigma \circ M_{\hat{L},i} \circ \sigma^{-1}$. To prove this equality let us take $p, q \in \mathbb{R}[\underline{X}]_d$. Then:

$$\begin{aligned} \langle M_{L,i}(\bar{p}^L), \bar{q}^L \rangle_L &= \langle \Pi_L(\overline{X_i p^L}), \bar{q}^L \rangle_L \stackrel{\Pi_L \circ \Pi_L = \Pi_L}{=} \langle \overline{X_i p^L}, \bar{q}^L \rangle_L = L(X_i p q) \stackrel{\hat{L} = L \text{ on } \mathbb{R}[\underline{X}]_{2d+1}}{=} \\ &= \hat{L}(X_i p q) = \langle \overline{X_i p^{\hat{L}}}, \bar{q}^{\hat{L}} \rangle_{\hat{L}} = \langle \overline{X_i p^{\hat{L}}}, \Pi_{\hat{L}}(\bar{q}^{\hat{L}}) \rangle_{\hat{L}} = \langle \Pi_{\hat{L}}(\overline{X_i p^{\hat{L}}}), \bar{q}^{\hat{L}} \rangle_{\hat{L}} = \\ &= \langle \sigma \circ \sigma^{-1}(\Pi_{\hat{L}}(\overline{X_i p^{\hat{L}}}), \bar{q}^{\hat{L}} \rangle_{\hat{L}} = \langle \sigma^{-1}(\Pi_{\hat{L}}(\overline{X_i p^{\hat{L}}}), \sigma^{-1}(\bar{q}^{\hat{L}}) \rangle_L = \langle \sigma^{-1} \circ M_{\hat{L},i}(\bar{p}^{\hat{L}}), \bar{q}^L \rangle_L = \\ &= \langle \sigma^{-1} \circ M_{\hat{L},i} \circ \sigma(\bar{p}^L), \bar{q}^L \rangle_L \end{aligned}$$

Finally we can conclude the truncated GNS multiplication operators of L commute using the commutativity of the truncated GNS multiplication operators of \hat{L} in the identical way we already did in the previous Theorem in (42). □

Corollary 6.3. Suppose L is flat, then \widetilde{M} is a generalized Hankel matrix.

Proof. If L is flat then by Theorem 5.18 the truncated GNS multiplication operators of L commute, and therefore by Theorem 6.2 we get that this is equivalent to \widetilde{M}_L being a generalized Hankel matrix. □

The following result uses the Theorem 6.1 together with ideas from [18] and give us a generalization of a classical Theorem from Mysovskikh [16], Dunkl and Xu [7, Theorem 3.8.7] and Putinar [18, pages 189-190]. They proved the equivalence between the existence of a minimal Gaussian quadrature rule with the commutativity of the truncated GNS multiplication operators for a positive definite linear form on $\mathbb{R}[\underline{X}]$. The generalization here comes from the fact that the result holds also if the linear form is defined on $\mathbb{R}[\underline{X}]_{2d+2}$

for $d \in \mathbb{N}_0$ and it is positive semidefinite i.e. we do not assume $U_L = \{0\}$. We also provide a third equivalent condition in the result which is $W_L^T A_L W_L$ is a generalized Hankel matrix, a fact which seems no to have been noticed so far.

Corollary 6.4. The following assertions are equivalent:

- (1) The linear form L admits a Gaussian quadrature rule.
- (2) The truncated GNS multiplication operators of L commute.
- (3) \widetilde{M}_L is a generalized Hankel matrix.

Proof. (1) \Rightarrow (2). Assume that L admits a Gaussian quadrature rule, that is to say $L(p) = \sum_{i=1}^N \lambda_i p(a_i)$ for all $p \in \mathbb{R}[\underline{X}]_{2d+1}$ where $N := \dim(T_L)$, the points a_1, \dots, a_N are pairwise different and $\lambda_1 > 0, \dots, \lambda_N > 0$. Let us set $\Lambda := \sum_{i=1}^N \lambda_i \text{ev}_{a_i} \in \mathbb{R}[\underline{X}]^*$. Using 5.8 we have the following:

$$(43) \quad \dim\left(\frac{\mathbb{R}[\underline{X}]}{U_\Lambda}\right) = N = \dim T_L$$

Let us consider again the linear isometry σ_1 , already defined in (32):

$$\sigma_1 : T_L \longrightarrow \frac{\mathbb{R}[\underline{X}]}{U_\Lambda}, \bar{p}^L \mapsto \bar{p}^\Lambda$$

As we proved in 5.10 is well defined and is an isometry, what implies σ_1 is injective, and considering that in this case it holds (43), σ_1 is moreover an isomorphism. We continue as in the implication 2 \Rightarrow 1 of the proof of Theorem 6.1, showing that the diagram (41) is commutative, what together with the fact that $M_{\Lambda,i}$ always commute for all $i \in \{1, \dots, n\}$ implies that the truncated GNS multiplication operators of L commute.

(2) \Rightarrow (1). This part was already proved in the Remark 5.14 as a consequence of the Theorem 5.13.

(2) \iff (3) It is the Theorem 6.2. □

The following result of Möller will give us a better lower bound in the number of nodes of a quadrature rule on $\mathbb{R}[\underline{X}]_{2d+1}$ than the very well-known bound given in Proposition 5.10. This bound, was already found for positive linear forms by Möller in 1975 and by Putinar in 1997 ([17],[18]). This result will show that the bound it is also true for positive semidefinite linear forms and it uses the same ideas as in [18]. We include the proof for the convenience of the reader. This bound will help us in polynomial optimization problems in which we know the number of global minimizers in advance, to discard optimality if this bound is bigger than the number of global minimizers, see Example 6.7 below.

Theorem 6.5. The number of nodes N of a Gaussian quadrature rule for L satisfies:

$$(44) \quad N \geq \dim(T_L) + \frac{1}{2} \max_{1 \leq j, k \leq n} (\text{rank}[M_{L,j}, M_{L,k}])$$

Proof. Assume L has a quadrature rule with N nodes, that is to say, there exist $\lambda_1 > 0, \dots, \lambda_N > 0$ weights and a_1, \dots, a_N in \mathbb{R}^n pairwise different nodes, such that $L(p) = \sum_{i=1}^N \lambda_i p(a_i)$ for all $p \in \mathbb{R}[\underline{X}]_{2d+1}$. Let us set $\Lambda := \sum_{i=1}^N \lambda_i \text{ev}_{a_i} \in \mathbb{R}[\underline{X}]^*$. By using the proposition 5.8 we have that:

$$\dim\left(\frac{\mathbb{R}[\underline{X}]}{U_\Lambda}\right) = N < \infty.$$

Then we can choose an orthonormal basis of $\frac{\mathbb{R}[\underline{X}]}{U_\Lambda}$. Let us denote such a basis by $\beta_\Lambda := \{\overline{\beta_1}^\Lambda, \dots, \overline{\beta_N}^\Lambda\}$ for $\beta_1, \dots, \beta_N \in \mathbb{R}[\underline{X}]$ pairwise different. Then we have that the transformation matrix of the multiplication operators $M_{\Lambda,i}$ with respect to this orthonormal basis is:

$$(\Lambda(X_i \beta_k \beta_j))_{1 \leq k, j \leq N}$$

The set $A := \{\bar{p}^\Lambda \mid p \in \mathbb{R}[\underline{X}]_d\}$ is a subspace of $\frac{\mathbb{R}[\underline{X}]}{U_\Lambda}$ so we can assume without loss of generality that $\beta_1, \dots, \beta_r \in \mathbb{R}[\underline{X}]_d$ where $r := \dim A$ generate a basis of A . Then since $L = \Lambda$ on $\mathbb{R}[\underline{X}]_{2d+1}$, we obtain:

$$(45) \quad (M(M_{\Lambda,i}, \beta_\Lambda)) := (\Lambda(X_i \beta_k \beta_j))_{1 \leq k, j \leq N} = \left(\frac{(\Lambda(X_i \beta_k \beta_j))_{1 \leq k, j \leq r}}{B_i^t} \mid \frac{B_i}{C_i} \right)$$

$(M(M_{\Lambda,i}, \beta_{\Lambda}))$ is the transformation matrix of the i -th truncated GNS multiplication operator of Λ with respect to the basis β_{Λ} and where $B_i \in \mathbb{R}^{r \times N-r}$ and $C_i \in \mathbb{R}^{N-r \times N-r}$ are symmetric matrices. We will show that $\beta_L := \{\overline{\beta_1^L}, \dots, \overline{\beta_r^L}\}$ is an orthonormal basis of T_L .

Taking σ_1 the isometry defined in (32), we get:

$$\sigma_1(T_L) = \{\sigma_1(\overline{p^L}) \mid \overline{p^L} \in T_L\} = \{\sigma_1(\overline{p^L}) \mid p \in \mathbb{R}[\underline{X}]_d\} = \{\overline{p^{\Lambda}} \mid p \in \mathbb{R}[\underline{X}]_d\} = A$$

Hence, we get:

$$\sigma_1(T_L) = \langle \overline{\beta_1^{\Lambda}}, \dots, \overline{\beta_r^{\Lambda}} \rangle$$

And since we have chosen $\beta_i \in \mathbb{R}[\underline{X}]_d$ for all $i \in \{1, \dots, r\}$, then we have $\sigma_1(\overline{\beta_i^L}) = \overline{\beta_i^{\Lambda}}$. Therefore $\beta_L := \{\overline{\beta_1^L}, \dots, \overline{\beta_r^L}\}$ generate a basis of T_L . It remains to show that β_L is orthonormal. To see that β_L is orthonormal we use again the fact that σ_1 is an isometry and that $\sigma_1(T_L) = A$. Indeed for $1 \leq i, j \leq r$:

$$\delta_{ij} = \Lambda(\beta_i \beta_j) = \langle \overline{\beta_j^{\Lambda}}, \overline{\beta_i^{\Lambda}} \rangle_{\Lambda} = \langle \sigma_1^{-1}(\overline{\beta_j^{\Lambda}}), \sigma_1^{-1}(\overline{\beta_i^{\Lambda}}) \rangle_L = \langle \overline{\beta_j^L}, \overline{\beta_i^L} \rangle_L$$

Therefore we have shown that:

$$M(M_{\Lambda,i}, \beta_{\Lambda}) = \left(\begin{array}{c|c} M(M_{L,i}, \beta_L) & B_i \\ \hline B_i^T & C_i \end{array} \right)$$

where we use the notation:

$$(M(M_{L,i}, \beta_L)) := (L(X_i \beta_k \beta_j))_{1 \leq k, j \leq N}$$

to refer us to the transformation matrix of the i -th truncated GNS multiplication operators of L with respect to the basis β_L . Using the fact that the matrices $M(M_{\Lambda,i}, \beta_{\Lambda})$ commute, we have the following equality:

$$M(M_{L,j}, \beta_L)M(M_{L,i}, \beta_L) - M(M_{L,i}, \beta_L)M(M_{L,j}, \beta_L) = B_i B_j^T - B_j B_i^T$$

Therefore the following it holds:

$$\text{rank}(B_i B_j^T - B_j B_i^T) \leq 2 \text{rank}(B_i B_j^T) \leq 2 \text{rank}(B_i) \leq 2 \min\{r, N-r\} \leq 2(N-r)$$

and then:

$$\text{rank}[M(M_{L,j}, \beta_L), M(M_{L,i}, \beta_L)] \leq 2(N-r)$$

Since we have already proved $r = \dim T_L$. And then we can conclude:

$$N \geq \dim(T_L) + \frac{1}{2} \max_{1 \leq j, k \leq n} (\text{rank}[M_{L,j}, M_{L,k}])$$

□

Remark 6.6. Note that we can use the previous Theorem 6.5 to show in a different way (1) \Rightarrow (2) in the Corollary 6.4. Indeed, let us suppose that L has a Gaussian quadrature rule that is to say with $N = \dim T_L$ nodes. Using the inequality (44) we get that $\text{rank}[M_{L,j}, M_{L,k}] = 0$ for $j, k \in \{1, \dots, n\}$ therefore the truncated GNS multiplication operators of L commute.

Example 6.7. Let us consider the following polynomial optimization problem taken from [11]:

$$\begin{aligned} & \text{minimize} && f(x) = x_1^2 x_2^2 (x^2 + y^2 - 1) \\ & \text{subject to} && x_1, x_2 \in \mathbb{R} \end{aligned}$$

By Calculus we know that the minimizers of f occur in the real points common to the partial derivatives of f (the real gradient variety) and we can easily check that this derivatives intersect in 4 real points: $(\pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}) \in \mathbb{R}^2$. Therefore we know in advance that (P) has at most 4 minimizers. On other side, an optimal solution of the moment relaxation of order 8 (P_8), that is $\mathbf{M} := M_{8,1}(y)$ read as:

$$\begin{pmatrix}
1.00 & 0.00 & 0.00 & 62.12 & -0.00 & 62.12 & 0.00 & 0.00 & 0.00 & 0.00 \\
0.00 & 62.12 & -0.00 & 0.00 & 0.00 & 0.00 & 9666.23 & -0.00 & 8.33 & -0.00 \\
0.00 & -0.00 & 62.12 & 0.00 & 0.00 & 0.00 & -0.00 & 8.33 & -0.00 & 9666.23 \\
62.12 & 0.00 & 0.00 & 9666.23 & -0.00 & 8.33 & 0.00 & -0.00 & 0.00 & 0.00 \\
-0.00 & 0.00 & 0.00 & -0.00 & 8.33 & -0.00 & -0.00 & 0.00 & 0.00 & -0.00 \\
62.12 & 0.00 & 0.00 & 8.33 & -0.00 & 9666.23 & 0.00 & 0.00 & -0.00 & 0.00 \\
0.00 & 9666.23 & -0.00 & 0.00 & -0.00 & 0.00 & 3150633.17 & -0.00 & 2.27 & 0.00 \\
0.00 & -0.00 & 8.33 & -0.00 & 0.00 & 0.00 & -0.00 & 2.27 & 0.00 & 2.27 \\
0.00 & 8.33 & -0.00 & 0.00 & 0.00 & -0.00 & 2.27 & 0.00 & 2.27 & -0.00 \\
0.00 & -0.00 & 9666.23 & 0.00 & -0.00 & 0.00 & 0.00 & 2.27 & -0.00 & 3150630.69 \\
9666.23 & 0.00 & -0.00 & 3150633.17 & -0.00 & 2.27 & 0.42 & -0.00 & -0.00 & 0.00 \\
-0.00 & -0.00 & 0.00 & -0.00 & 2.27 & 0.00 & -0.00 & -0.00 & 0.00 & 0.00 \\
8.33 & 0.00 & 0.00 & 2.27 & 0.00 & 2.27 & -0.00 & 0.00 & 0.00 & -0.00 \\
-0.00 & 0.00 & -0.00 & 0.00 & 2.27 & -0.00 & 0.00 & 0.00 & -0.00 & -0.00 \\
9666.23 & -0.00 & 0.00 & 2.27 & -0.00 & 3150630.69 & 0.00 & -0.00 & -0.00 & 0.33 \\
& & & 9666.23 & -0.00 & 8.33 & -0.00 & 9666.23 & & \\
& & & 0.00 & -0.00 & 0.00 & 0.00 & -0.00 & & \\
& & & -0.00 & 0.00 & 0.00 & 0.00 & -0.00 & & \\
& & & 3150633.17 & -0.00 & 2.27 & 0.00 & 2.27 & & \\
& & & -0.00 & 2.27 & 0.00 & 2.27 & -0.00 & & \\
& & & 2.27 & 0.00 & 2.27 & -0.00 & 3150630.69 & & \\
& & & 0.42 & -0.00 & -0.00 & 0.00 & 0.00 & & \\
& & & -0.00 & -0.00 & 0.00 & 0.00 & -0.00 & & \\
& & & -0.00 & 0.00 & 0.00 & -0.00 & -0.00 & & \\
& & & 0.00 & 0.00 & -0.00 & -0.00 & -0.00 & & \\
& & & 2466755083.36 & -43.48 & 169698627.89 & -6.08 & 134568970.57 & & \\
& & & -43.48 & 169698627.89 & -6.08 & 134568970.57 & 15.08 & & \\
& & & 169698627.89 & -6.08 & 134568970.57 & 15.08 & 169698562.66 & & \\
& & & -6.08 & 134568970.57 & 15.08 & 169698562.66 & 25.61 & & \\
& & & 134568970.57 & 15.08 & 169698562.66 & 25.61 & 2466752654.76 & &
\end{pmatrix}$$

and the rank of the commutator of the truncated GNS multiplication operators is:

$$\text{rank}[M_{\mathbf{M}, X_1}, M_{\mathbf{M}, X_1}] =
\begin{pmatrix}
0 & 0.00 & 0.00 & -0.00 & -0.00 & 0.00 & 0.00 & -0.00 & 0.00 & -0.00 \\
-0.00 & 0 & -0.00 & 0.00 & 0.00 & 0.00 & -0.00 & 0.00 & 0.00 & -0.00 \\
-0.00 & 0.00 & 0 & 0.00 & 0.00 & 0.00 & 0.00 & -0.00 & 0.00 & 0.00 \\
0.00 & -0.00 & -0.00 & 0 & -0.00 & -0.00 & -0.00 & -0.00 & 0.00 & -0.00 \\
0.00 & -0.00 & -0.00 & 0.00 & 0 & 0.00 & 0.00 & 0.00 & 0.00 & 0.00 \\
-0.00 & -0.00 & -0.00 & 0.00 & -0.00 & 0 & -0.00 & -0.00 & 0.00 & -0.00 \\
-0.00 & 0.00 & -0.00 & 0.00 & -0.00 & 0.00 & 0 & -313.91 & -0.00 & 115.50 \\
0.00 & -0.00 & 0.00 & 0.00 & -0.00 & 0.00 & 313.91 & 0 & 0.18 & 0.00 \\
-0.00 & -0.00 & -0.00 & -0.00 & -0.00 & -0.00 & 0.00 & -0.18 & 0 & 0.05 \\
0.00 & 0.00 & -0.00 & 0.00 & -0.00 & 0.00 & -115.50 & -0.00 & -0.05 & 0
\end{pmatrix} = 4$$

If \mathbf{M} had a quadrature rule on $\mathbb{R}[X_1, X_2]_7$ with N nodes, since $f \in \mathbb{R}[X_1, X_2]_7$ by 3.9 (iii) the N nodes of the quadrature rule would be global minimizers of f and $P^* = P_8^*$, and according to Theorem 6.5:

$$N \geq \dim(T_L) + \frac{1}{2} \max_{1 \leq j, k \leq n} (\text{rank}[M_{L,j}, M_{L,k}]) = 10 + \frac{1}{2}4 = 12$$

Therefore the polynomial f would have at least 12 global minimizers and this is a contradiction with the fact that f has at most 4 global minimizers. Notice that then \mathbf{M} does not have a quadrature rule on $\mathbb{R}[X]_7$, and in particular it does not have a quadrature rule.

7. ALGORITHM FOR EXTRACTING MINIMIZERS IN POLYNOMIAL OPTIMIZATION PROBLEMS

As an application of all the previous results in this section we find a stopping criterion for the moment relaxation hierarchy, in other words, we find a condition on the optimal solution of (P_d) L , such that $L(f) = P_d^* = P^*$. In this case we also find potential global minimizers. In [9] Henrion and Lasserre the stopping criterion was L to be flat and in this algorithm the stopping criterion is $W_L^T M_L W_L$ being Hankel, and as we have already seen in 6.3 this condition is more general. It is important to point out that despite this condition is more general than being flat we can not ensure optimality until we check that the candidate to minimizers are inside to the basic closed semialgebraic set S , condition that it is always possible to ensure if the set S is a set described with linear polynomials, if the set S is \mathbb{R}^n or we have flat extension of some degree on the optimal solution, that is to say $\text{rank } M_d(y) = \text{rank } M_s(y)$ for sufficient small s , see [13, Theorem 6.18] or [5, Theorem 1.6] for a proof. At the end of this paper we summarize all this results in an algorithm with examples and also we illustrate polynomial optimization problems where this new stopping criterion allow us to conclude optimality even in case where the optimal solution is not flat as we already advance in 2.9 and in 2.11.

Theorem 7.1. Let $f, p_1, \dots, p_m \in \mathbb{R}[\underline{X}]_{2d}$ and L be an optimal solution of (P_{2d}) . Suppose that $W_L^T A_L W_L$ is a generalized Hankel matrix. Then L has a quadrature rule on G_L . Moreover, suppose the nodes of the quadrature rule lie on S and $f \in \mathbb{R}[\underline{X}]_{2d-1}$, then $L(f) = P^*$ and the nodes are global minimizers.

Proof. Since $W_L^T A_L W_L$ is Hankel by Corollary 6.2 and 5.13 there exists nodes $a_1, \dots, a_N \in \mathbb{R}^n$ and weights $\lambda_1 > 0, \dots, \lambda_N > 0$, where $N := \dim T_L$ such that:

$$L(p) = \sum_{i=1}^N \lambda_i p(a_i) \text{ for all } p \in G_L$$

Moreover if the nodes of this quadrature rule are contained in S by 3.9 (iii) $P^* = P_d^* = f(a_i)$ for $i \in \{1, \dots, N\}$. \square

The following Lemma was already proved in [12, lemma 2.7]. We will use it to prove the Corollary 7.3.

Lemma 7.2. Let $L = \sum_{i=1}^N \lambda_i \text{ev}_{a_i} \in \mathbb{R}[\underline{X}]_{2d}^*$ for a_1, \dots, a_N pairwise different points and $\lambda_1 > 0, \dots, \lambda_N > 0$ such that L is flat. Then there exist interpolation polynomials $q_1, \dots, q_N \in \mathbb{R}[\underline{X}]_{2d}$ at the points a_1, \dots, a_N of degree at most $d-1$.

Proof. Let us consider the isometry map (32) already defined in 5.10:

$$\sigma_1 : T_L \longrightarrow \frac{\mathbb{R}[\underline{X}]}{U_\Lambda}, \bar{p}^L \mapsto \bar{p}^\Lambda, \text{ for } p \in \mathbb{R}[\underline{X}]_{d-1}$$

It is moreover an isomorphism of euclidean vector spaces since $\dim(\frac{\mathbb{R}[\underline{X}]}{U_\Lambda}) = N$ by Proposition 5.8. It is very well known that there exists interpolation polynomial $h_1, \dots, h_N \in \mathbb{R}[\underline{X}]$ at the points a_1, \dots, a_N , such that $h_i(a_j) = \delta_{i,j}$ for $i, j \in \{1, \dots, n\}$. Define $\bar{q}_j^L := \sigma^{-1}(\bar{p}^\Lambda)$ for $q_j \in \mathbb{R}[\underline{X}]_{d-1}$. Then for $j \in \{1, \dots, N\}$:

$$0 \leq \sum_{i=1}^N \lambda_i q_j^2(a_i) = L(q_j^2) = \langle \bar{q}_j^L, \bar{q}_j^L \rangle_L = \langle \bar{h}_j^\Lambda, \bar{h}_j^\Lambda \rangle_\Lambda = \Lambda(h_j^2) = \lambda_j$$

and therefore $q_j(a_i) = \delta_{i,j}$ for $i, j \in \{1, \dots, N\}$. \square

Corollary 7.3. Let $p_1, \dots, p_m \in \mathbb{R}[\underline{X}]_1$ and L be an optimal solution of (P_{2d}) with $f \in \mathbb{R}[\underline{X}]_{2d-1}$. Suppose that $W_L^T A_L W_L$ is a Hankel matrix. Then L has a quadrature rule representation on G_L , $L(f) = P^*$ and the nodes are minimizers of (P) .

Proof. From Theorem 7.1 there exists nodes $a_1, \dots, a_N \in \mathbb{R}^n$ and weights $\lambda_1 > 0, \dots, \lambda_N > 0$, where $N := \dim T_L$ such that:

$$L(p) = \sum_{i=1}^N \lambda_i p(a_i) \text{ for all } p \in G_L$$

To conclude the Corollary by Theorem 7.1 it is enough to show that the nodes a_1, \dots, a_N are contained in S . In Theorem 6.1 we proved that $\hat{L} := \sum_{i=1}^N \lambda_i \text{ev}_{a_i} \in \mathbb{R}[\underline{X}]_{2d}^*$ is flat. Then by the Lemma 7.2 there are interpolation polynomials q_1, \dots, q_N at the points a_1, \dots, a_N having at most degree $d-1$. Since $\deg(q_i^2 p_j) \leq 2d-1$ then $q_i^2 p_j \in T_{2d}(p_1, \dots, p_m)$ and therefore:

$$0 \leq L(q_i^2 p_j) = \hat{L}(q_i^2 p_j) = \lambda_i p_j(a_i)$$

This equality proves that $p_j(a_i) \geq 0$ for $j \in \{1, \dots, m\}$ and $i \in \{1, \dots, N\}$ so we can conclude $\{a_1, \dots, a_N\} \subseteq S$. \square

Remark 7.4. The above results: Theorem 7.1 and Corollary 7.3 can be written in terms of an optimal solution of a Moment relaxation of even degree by taking as an optimal solution its restriction to one degree less.

Algorithm 1: Algorithm for extracting minimizers of (P)

Input: A polynomial optimization problem (P) (2).

Output: The minimum \mathbf{P}^* and minimizers $\mathbf{a}_1, \dots, \mathbf{a}_r \subseteq \mathbf{S}$ of (P).

- 1 $\mathbf{k} := \max\{\deg f, \deg p_1, \dots, \deg p_m\}$
- 2 Compute an optimal solution $\mathbf{M} := M_{\lfloor \frac{\mathbf{k}}{2} \rfloor}(\mathbf{y})$ of the moment relaxation $(\mathbf{P}_{\mathbf{k}})$ and also compute $\mathbf{W}_{\mathbf{M}}$ matrix such that:

$$\mathbf{M} = \left(\begin{array}{c|c} A_{\mathbf{M}} & A_{\mathbf{M}}\mathbf{W}_{\mathbf{M}} \\ \hline \mathbf{W}_{\mathbf{M}}^T A_{\mathbf{M}} & C_{\mathbf{M}} \end{array} \right)$$

- 3 **if** $\mathbf{W}_{\mathbf{M}}^T A_{\mathbf{M}} \mathbf{W}_{\mathbf{M}}$ is a Hankel matrix **then**
 - 4 | go to 7
 - 5 **else**
 - 6 | $\mathbf{k} := \mathbf{k} + 1$ and go to 2.
 - 7 **if** (\mathbf{k} even and $f \in \mathbb{R}[\mathbf{X}]_{\mathbf{k}-1}$) **or** (\mathbf{k} odd and $f \in \mathbb{R}[\mathbf{X}]_{\mathbf{k}-2}$) **then**
 - 8 | go to 14
 - 9 **else**
 - 10 | **if** $C_{\mathbf{M}} = \mathbf{W}_{\mathbf{M}} A_{\mathbf{M}} \mathbf{W}_{\mathbf{M}}$ **then**
 - 11 | | go to 14
 - 12 | **else**
 - 13 | | $\mathbf{k} := \mathbf{k} + 1$ go to 2
 - 14 Compute the truncated multiplication operators of \mathbf{M} : $\mathbf{A}_{1,\mathbf{M}}, \dots, \mathbf{A}_{n,\mathbf{M}}$ and go to 15.
 - 15 Compute an orthonormal basis $\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$ of $\mathbf{T}_{\mathbf{M}}$ of common eigenvectors of the truncated multiplication operators such that $\mathbf{A}_{i,\mathbf{M}}\mathbf{v}_j = \mathbf{a}_{j,i}\mathbf{v}_j$ and go to 16.
 - 16 **if** $\mathbf{a}_1, \dots, \mathbf{a}_n \in \mathbf{S}$ **then**
 - 17 | go to 20
 - 18 **else**
 - 19 | $\mathbf{k} := \mathbf{k} + 1$ and go to 2
 - 20 We can conclude that the points $\{\mathbf{a}_1, \dots, \mathbf{a}_r\} \subseteq \mathbf{S}$ are minimizers of (P), and $\mathbf{P}^* = \mathbf{f}(\mathbf{a}_i)$ for all $i \in \{1, \dots, r\}$
-

8. SOFTWARE AND EXAMPLES

To find an optimal solution of the Moment relaxation and for the big calculations we have used the following softwares:

- YALMIP: developed by J. Löfberg. It is a toolbox for Modeling and Optimization in MATLAB. Published in the Journal Proceedings of the CACSD Conference in 2004. For more information see: <http://yalmip.github.io/>.
- SEDUMI: developed by J. F. Sturm. It is a toolbox for optimization over symmetric cones. Published in the Journal Optimization Methods and Software in 1999. For more information see: <http://sedumi.ie.lehigh.edu/>.
- MPT: the Multi-parametric Toolbox is an open source, Matlab-based toolbox for parametric optimization, computational geometry and model predictive control. For more information see: <http://people.ee.ethz.ch/~mpt/3/>.
- MATLAB and Statistics Toolbox Release 2016a, The MathWorks, Inc., Natick, Massachusetts, United States.

$$W_{\mathbf{M}} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

and

$$c_{\mathbf{M}} = \begin{matrix} X_1^3 & X_1^2 X_2 & X_1^2 X_3 & X_1 X_2^2 & X_1 X_2 X_3 & X_1 X_3^2 & X_2^3 & X_2^2 X_3 & X_2 X_3^2 & X_3^3 \\ X_1^3 & 5.2880 & 0.9994 & 0.9994 & 2.4826 & 0.9989 & 2.4744 & 0.9988 & 1.0004 & 1.0020 & 1.0001 \\ X_1^2 X_2 & 0.9994 & 2.4826 & 0.9989 & 0.9988 & 1.0004 & 1.0020 & 2.4832 & 1.0010 & 1.6671 & 1.0007 \\ X_1^2 X_3 & 0.9994 & 0.9989 & 2.4744 & 1.0004 & 1.0020 & 1.0001 & 1.0010 & 1.6671 & 1.0007 & 2.4638 \\ X_1 X_2^2 & 2.4826 & 0.9988 & 1.0004 & 2.4832 & 1.0010 & 1.6671 & 0.9983 & 1.0007 & 1.0015 & 1.0001 \\ X_1 X_2 X_3 & 0.9989 & 1.0004 & 1.0020 & 1.0010 & 1.6671 & 1.0007 & 1.0007 & 1.0015 & 1.0001 & 1.0016 \\ X_1 X_3^2 & 2.4744 & 1.0020 & 1.0001 & 1.6671 & 1.0007 & 2.4638 & 1.0015 & 1.0001 & 1.0016 & 0.9912 \\ X_2^3 & 0.9988 & 2.4832 & 1.0010 & 0.9983 & 1.0007 & 1.0015 & 5.2883 & 1.0071 & 2.4669 & 1.0072 \\ X_2^2 X_3 & 1.0004 & 1.0010 & 1.6671 & 1.0007 & 1.0015 & 1.0001 & 1.0071 & 2.4669 & 1.0072 & 2.4579 \\ X_2 X_3^2 & 1.0020 & 1.6671 & 1.0007 & 1.0015 & 1.0001 & 1.0016 & 2.4669 & 1.0072 & 2.4579 & 1.0040 \\ X_3^3 & 1.0001 & 1.0007 & 2.4638 & 1.0001 & 1.0016 & 0.9912 & 1.0072 & 2.4579 & 1.0040 & 14.6604 \end{matrix}$$

We calculate that:

$$W_{\mathbf{M}}^T A_{\mathbf{M}} C_{\mathbf{M}} = \begin{pmatrix} 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 \\ 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 \\ 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 \\ 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 \\ 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 \\ 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 \\ 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 \\ 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 \\ 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 \\ 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 \end{pmatrix}$$

is a generalized Hankel matrix and $f \in \mathbb{R}[X_1, X_2, X_3]_5$. By Theorem 7.3 we have optimality with optimal value $P^* = P_6^* = 2.3527 \cdot 10^{-8} \approx 0$. Finally we get that the matrices of the truncated GNS operators with respect to the orthonormal basis $v := \langle 1 \rangle_{\mathbf{M}}$ are:

$$M(M_{\mathbf{M}, X_1}, v) = (1), M(M_{\mathbf{M}, X_2}, v) = (1) \text{ and } M(M_{\mathbf{M}, X_3}, v) = (1)$$

The operators are in diagonal form so we have already an orthonormal basis of $T_{\mathbf{M}}$ of common eigenvectors of the truncated GNS operators of \mathbf{M} $v := \langle 1 \rangle_{\mathbf{M}}$, then a global minimizer is $(1, 1, 1) \in \mathbb{R}^n$, and:

$$\widetilde{\mathbf{M}} = V_3(1, 1, 1)V_3^T(1, 1, 1).$$

Example 8.2. Let us consider the following polynomial optimization problem, defined on a non convex closed semialgebraic set, taken from [8, problem 4.6] :

$$\begin{aligned} & \text{minimize} && f(x) = -x_1 - x_2 \\ & \text{subject to} && x_2 \leq 2x_1^4 - 8x_1^3 + 8x_1^2 + 2 \\ & && x_2 \leq 4x_1^4 - 32x_1^3 + 88x_1^2 - 96x_1 + 36 \\ & && 0 \leq x_1 \leq 3 \\ & && 0 \leq x_2 \leq 4 \end{aligned}$$

We initialize $\mathbf{k} = 4$. An optimal solution of (P_4) reads as:

(47)

$$\mathbf{M} := M_{4,1}(y) = \begin{matrix} & 1 & X_1 & X_2 & X_1^2 & X_1 X_2 & X_2^2 \\ 1 & 1.0000 & 3.0000 & 4.0000 & 9.0000 & 12.0000 & 16.0000 \\ X_1 & 3.0000 & 9.0000 & 12.0000 & 27.0000 & 36.0000 & 48.0000 \\ X_2 & 4.0000 & 12.0000 & 16.0000 & 36.0000 & 48.0000 & 64.0000 \\ X_1^2 & 9.0000 & 27.0000 & 36.0000 & 107.6075 & 109.0814 & 176.3211 \\ X_1 X_2 & 12.0000 & 36.0000 & 48.0000 & 109.0814 & 176.3211 & 194.9661 \\ X_2^2 & 16.0000 & 48.0000 & 64.0000 & 176.3211 & 194.9661 & 368.5439 \end{matrix}$$

and

$$(48) \quad \widetilde{\mathbf{M}} = \begin{array}{c} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \end{array} \begin{array}{c} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \end{array} \begin{array}{c} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \end{array} \left(\begin{array}{ccc|ccc} 1.0000 & 3.0000 & 4.0000 & 9.0000 & 12.0000 & 16.0000 \\ 3.0000 & 9.0000 & 12.0000 & 27.0000 & 36.0000 & 48.0000 \\ 4.0000 & 12.0000 & 16.0000 & 36.0000 & 48.0000 & 64.0000 \\ \hline 9.0000 & 27.0000 & 36.0000 & 81.0000 & 108.0000 & 144.0000 \\ 12.0000 & 36.0000 & 48.0000 & 108.0000 & 144.0000 & 192.0000 \\ 16.0000 & 48.0000 & 64.0000 & 144.0000 & 192.0000 & 256.0000 \end{array} \right)$$

taking for example:

$$W_{\mathbf{M}} = \begin{pmatrix} 9 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 3 & 4 \end{pmatrix}$$

$\widetilde{\mathbf{M}}$ is a generalized Hankel matrix. The matrix of the truncated GNS multiplaction operators with respect to the orthonormal basis $v = \langle \widetilde{\mathbf{1}}^{\mathbf{M}} \rangle$ are:

$$M(M_{\mathbf{M},1}, v) = \begin{pmatrix} 3 \end{pmatrix} \text{ and } M(M_{\mathbf{M},1}, v) = \begin{pmatrix} 4 \end{pmatrix}$$

Hence the candidate to minimizer is $(3, 4)$, however it does not lie in S , then $(3, 4)$ cannot be a minimizer and $f(3, 4) = -7$ cannot be the minimum. Then we try with a relaxation of order $\mathbf{k} = 5$. An optimal solution of the moment relaxation (P_5) is the following:

$$(49) \quad \mathbf{M} := M_{5,1}(y) = \begin{array}{c} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \end{array} \begin{array}{c} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \end{array} \begin{array}{c} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \end{array} \left(\begin{array}{ccc|ccc} 1.00 & 2.67 & 4.00 & 8.00 & 10.67 & 16.00 \\ 2.67 & 8.00 & 10.67 & 24.00 & 32.00 & 42.67 \\ 4.00 & 10.67 & 16.00 & 32.00 & 42.67 & 64.00 \\ \hline 8.00 & 24.00 & 32.00 & 72.00 & 96.00 & 128.00 \\ 10.67 & 32.00 & 42.67 & 96.00 & 128.00 & 170.67 \\ 16.00 & 42.67 & 64.00 & 128.00 & 170.67 & 256.00 \end{array} \right)$$

In this case $C_{\mathbf{M}} = W_{\mathbf{M}}^T A_{\mathbf{M}} W_{\mathbf{M}}$, therefore \mathbf{M} is flat and in particular the operators commute by 5.18. After the simultaneous diagonalization of the truncated GNS operators we get that the candidate to minimizers are $(0, 4) \notin S$ and $(3, 4) \notin S$. Hence we try with a relaxation of order $\mathbf{k} = 6$. An optimal solution of the moment relaxation (P_6) reads as:

$$(50) \quad \mathbf{M} := M_{6,1}(y) = \begin{array}{c} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \\ X_1^3 \\ X_1^2 X_2 \\ X_1 X_2^2 \\ X_2^3 \end{array} \begin{array}{c} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \\ X_1^3 \\ X_1^2 X_2 \\ X_1 X_2^2 \\ X_2^3 \end{array} \begin{array}{c} 1 \\ X_1 \\ X_2 \\ X_1^2 \\ X_1 X_2 \\ X_2^2 \\ X_1^3 \\ X_1^2 X_2 \\ X_1 X_2^2 \\ X_2^3 \end{array} \left(\begin{array}{cccc|cccc|cc} 1.00 & 2.67 & 4.00 & 8.00 & 10.67 & 16.00 & 24.00 & 32.00 & 42.67 & 64.00 \\ 2.67 & 8.00 & 10.67 & 24.00 & 32.00 & 42.67 & 72.00 & 96.00 & 128.00 & 170.67 \\ 4.00 & 10.67 & 16.00 & 32.00 & 42.67 & 64.00 & 96.00 & 128.00 & 170.67 & 256.00 \\ \hline 8.00 & 24.00 & 32.00 & 72.00 & 96.00 & 128.00 & 216.00 & 288.00 & 384.00 & 512.00 \\ 10.67 & 32.00 & 42.67 & 96.00 & 128.00 & 170.67 & 288.00 & 384.00 & 512.00 & 682.66 \\ 16.00 & 42.67 & 64.00 & 128.00 & 170.67 & 256.00 & 384.00 & 512.00 & 682.66 & 1024.00 \\ \hline 24.00 & 72.00 & 96.00 & 216.00 & 288.00 & 384.00 & 204299.70 & 870.25 & 19035.69 & 1583.15 \\ 32.00 & 96.00 & 128.00 & 288.00 & 384.00 & 512.00 & 870.25 & 19035.69 & 1583.15 & 18023.54 \\ 42.67 & 128.00 & 170.67 & 384.00 & 512.00 & 682.66 & 19035.69 & 1583.15 & 18023.54 & 2822.34 \\ 64.00 & 170.67 & 256.00 & 512.00 & 682.66 & 1024.00 & 1583.15 & 18023.54 & 2822.34 & 58336.42 \end{array} \right)$$

and

$$W_{\mathbf{M}}^T A_{\mathbf{M}} W_{\mathbf{M}} = \begin{array}{c} X_1^3 \\ X_1^2 X_2 \\ X_1 X_2^2 \\ X_2^3 \end{array} \begin{array}{c} X_1^3 \\ X_1^2 X_2 \\ X_1 X_2^2 \\ X_2^3 \end{array} \begin{array}{c} X_1^3 \\ X_1^2 X_2 \\ X_1 X_2^2 \\ X_2^3 \end{array} \left(\begin{array}{cccc} 648.00 & 863.99 & 1151.99 & 1535.99 \\ 863.99 & 1151.99 & 1535.99 & 2047.99 \\ 1151.99 & 1535.99 & 2047.99 & 2730.65 \\ 1535.99 & 2047.99 & 2730.65 & 4095.99 \end{array} \right)$$

is a Hankel matrix. However we get the same candidate to minimizers as in the previous relaxation which does not belong to S . Finally we increase to $\mathbf{k} = 7$, and we get after rounding, the following optimal solution of (P_7) :

Again we follow the same idea as in [15, algorithm 4.1 Step 1] to apply simultaneous diagonalization to the matrices A_1 and A_2 . For this we find the orthogonal matrix P that diagonalize a matrix of the following form:

$$A = r_1 A_1 + r_2 A_2 \text{ where } r_1^2 + r_2^2 = 1$$

For

$$P = \begin{pmatrix} 0.7589 & 0.5572 & 0.3371 \\ -0.6512 & 0.6493 & 0.3929 \\ 0.0000 & -0.5177 & 0.8556 \end{pmatrix}$$

we get the following diagonal matrices:

$$P^T A_1 P = \begin{pmatrix} 1.0000 & 0.0000 & -0.0000 \\ 0.0000 & 2.0000 & 0.0000 \\ -0.0000 & 0.0000 & 2.0000 \end{pmatrix}, P^T A_2 P = \begin{pmatrix} 2.0000 & -0.0000 & 0.0000 \\ -0.0000 & 2.0000 & -0.0000 \\ 0.0000 & -0.0000 & 3.0000 \end{pmatrix}$$

and with the operation:

$$P^T \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0.7589 \\ 0.5572 \\ 0.3371 \end{pmatrix}$$

we get the square roots of the weights of the quadrature formula. Then we have the following decomposition:

$$\mathbf{M} = \widetilde{\mathbf{M}} = 0.5759 V_2(1, 2) V_2^T(1, 2) + 0.3105 V_2(2, 2) V_2^T(2, 2) + 0.1137 V_2(2, 3) V_2^T(2, 3)$$

In this case the points $(1, 2)$, $(2, 2)$, and $(2, 3)$ lie on S , as we already know since it holds the condition of the Theorem 1.6 in [5], and therefore they are global minimizers of (P) , and the minimum is $P^* = P_4^* = -2$.

ACKNOWLEDGEMENTS

This is part of the ongoing research for the authors PhD thesis. I want to thank Markus Schweighofer for supervising my thesis and many fruitful discussions.

REFERENCES

- [1] Christian Bayer and Josef Teichmann: The proof of Tchakaloff's Theorem. Proceedings of the American Mathematical Society Volume **134**, Number 10. October 2006. Pages 3035-3040, May 4, 2006.
- [2] D. A. Cox, J. Little, D. O'Shea: Ideals, Varieties and Algorithms. An introduction to Computational Algebraic Geometry and Commutative Algebra. Fourth Edition. Springer. Undergraduate text in Mathematics.
- [3] R. E. Curto, L. A. Fialkow: Recursiveness, positivity and truncated moment problems. Houston Journal of Mathematics. Volum **17**, No.4, 1991.
- [4] R. E. Curto and L. A. Fialkow: Solution of the truncated complex moment problem for flat data, Memoirs of the American Mathematical Society **119** (568), 1996.
- [5] R. E. Curto, L. A. Fialkow: The truncated complex K-moment problem. Transactions of the mathematical society Volume **352**, Number 6, Pages 2825-2855, February 28, 2000.
- [6] R. E. Curto and L. A. Fialkow: Solution of the singular quartic moment problem, J. Operator Theory, Volume **48**, pages 315-354, 2002.
- [7] C. F. Dunkl and Y. Xu: Orthogonal Polynomials of several variables. Second Edition. Encyclopedia of Mathematics and Its Applications 2014.
- [8] C. A. Floudas and P.M Pardalos, A collection of Test Problems for Constrained Global Optimization Algorithms, Lecture Notes in Computer Sciences Sci:455, Springer-Verlag, Berlin, 1990.
- [9] D. Henrion and J. B. Lasserre, Gloptipoly: Global optimization over polynomials with MATLAB and SEDUMI. ACM. Transactions math Soft. **29**:165-194. 2003.
- [10] E. de Klerk, M. Laurent, Zhao Sun: Convergence analysis for Lasserre's measure-based hierarchy of upper bounds for polynomial optimization, last revised 8 Sep 2015. arXiv:1411.6867v4.
- [11] J. B. Lasserre: Global optimization with polynomials an the problems of moments, SIAM J.Optim. **11**, No. **3**, 796-817 ,2001
- [12] M. Laurent: Revisiting two theorems of Curto and Fialkow on moment matrices. Proceedings of the American Mathematical Society. Volume 133. Number **10** pages 2965-2976. May 9, 2009.
- [13] M. Laurent: Sums of squares, moment matrices and optimization over polynomials, February 6, 2010
- [14] M. Marshall: Positive Polynomials and sums of squares. Amer. Math. Soc, Mathematical Surveys and Monographs, vol:**146**,2000
- [15] M. Murota, Y. Kanno, M. Kojima, S. Kojima: A numerical Algorithm for Block-Diagonal Decomposition of Matrix*-Algebras, Part I: Proposed Approach and Application to semidefinite programming. Jpn.J.Ind.Appl.Math. Volume **27**, pages 125-160, June 2010.
- [16] I.P. Mysovskikh, Interpolatory Cubature Formulas, Nauka, Moscow, 1981 (in Russian). Interpolatorische Kubaturformel, Institut für Geometrie und Praktische Mathematik der RWTH Aachen, 1992, Berich No.74 (in German)

- [17] H. M. Möller, Kubaturformeln mit minimaler Knotenzahl. *Numerische Mathematik* **25** 1976, 185-200. (in German)
- [18] M. Putinar: A Dilation theory Approach to Cubature Formulas. *Expo. Math* **15** 183-192 Heidelberg, 1997
- [19] C. Riener, M. Schweighofer: Optimization approaches to quadrature: new characterizations of Gaussian quadrature on the line and quadrature with few nodes on plane algebraic curves, on the plane and in higher dimensions, arXiv preprint: 1607.08404, July 2016.
- [20] M. Schweighofer: Polynomiale Optimierung unpublished Lecture Notes. Universität Konstanz, Summer Semester 2012 and 2015.
- [21] M. Schweighofer: Optimization of polynomials on compact semialgebraic sets, *SIAM J.Optim* **15** No. 3, 805-825, 2005.
- [22] M. Schweighofer and I. Klep: An Exact Duality Theory for Semidefinite Programming Based on Sums of Squares. *Mathematics of Operations Research* **38**, 569-590 2013
- [23] J. L. Smul'jan, An operator Hellinger integral (Russian), *Mat Sb* **91** 1959; 381-430.