



INSTITUT DE FRANCE
Académie des sciences

Comptes Rendus

Mathématique


Mickaël Matusinski and Simon Müller

Quantifier elimination for quasi-real closed fields

Volume 359, issue 3 (2021), p. 291-295.

<https://doi.org/10.5802/crmath.169>

© Académie des sciences, Paris and the authors, 2021.
Some rights reserved.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



Les Comptes Rendus. Mathématique sont membres du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org



Mathematical logic / *Logique mathématique*

Quantifier elimination for quasi-real closed fields

Mickaël Matusinski^a and Simon Müller^{*, b}

^a Univ. Bordeaux, IMB, UMR 5251, F-33400 Talence, France

^b Universität Konstanz, 78467 Konstanz, Germany

E-mails: mickael.matusinski@math.u-bordeaux.fr (M. Matusinski),
simon.2.mueller@uni-konstanz.de (S. Müller)

Abstract. We prove quantifier elimination for the theory of quasi-real closed fields with a compatible valuation. This unifies the same known results for algebraically closed valued fields and real closed valued fields.

Résumé. Nous prouvons l'élimination des quantificateurs pour la théorie des corps quasi-réels clos munis d'une valuation compatible. Cela reprend et unifie les mêmes résultats connus pour les corps algébriquement clos et les corps réels clos.

2020 Mathematics Subject Classification. 03C10, 03C64, 12J10, 12J15, 12L12.

Manuscript received 26th May 2020, accepted 13th December 2020.

1. Introduction

Ordered fields and valued fields share several similar features, most notably via the notion of (order-)compatible valuations (see e.g. [9, 10, 13] and below). In the present note we illustrate this by subsuming the following major model theoretic results: the (first order) theories of algebraically closed fields with a non-trivial valuation, respectively real closed fields with a *compatible* non-trivial valuation, both admit quantifier elimination (cf. [11], respectively [3]). We prove that the same applies if we consider the theory of the union of these two classes in a common language, the one of quasi-ordered fields. Quasi-ordered fields were introduced in [7] and provide a uniform axiomatisation of ordered and valued fields. We derive a notion of quasi-real closed field, which subsumes the classes of algebraically closed fields and real closed fields. Note that real closed fields were developed by Artin and Schreier in [1] as maximal ordered fields, in analogy to the notion of algebraic closure of a field.

* Corresponding author.

2. Preliminaries on quasi-ordered fields

Throughout this note, all orderings and quasi-orderings are always assumed to be total. An *ordered field* is a field K equipped with an ordering \leq (i.e. a binary, reflexive, transitive and anti-symmetric relation) such that the following axioms are satisfied for all $x, y, z \in K$:

$$(O1) \quad x \leq y \wedge 0 \leq z \Rightarrow xz \leq yz,$$

$$(O2) \quad x \leq y \Rightarrow x + z \leq y + z.$$

Orderings are in one-to-one correspondence with so-called *positive cones*, i.e. subsets $P \subseteq K$ such that $PP \subseteq P$, $P + P \subseteq P$, $P \cup (-P) = K$ and $-1 \notin P$: $0 \leq x \Leftrightarrow x \in P$.

A valuation v on a field K (in the sense of Krull [8]) is known to endow K with a *quasi-ordering*, i.e. a binary, reflexive, transitive relation \preceq , as follows:

$$a \preceq b \iff v(b) \leq v(a).$$

If \preceq is a quasi-ordering, we write $x \simeq y$ as shorthand for $x \preceq y \preceq x$. Note that if \preceq is an ordering, then \simeq is just equality, whereas if \preceq is induced by a valuation v as above, then $x \simeq y$ if and only if $v(x) = v(y)$. Moreover, we write $x < y$ as shorthand for $x \preceq y$ and $x \neq y$.

Pushing the analogy of ordered and valued fields further, Fakhruddin introduced the notion of *quasi-ordered field*, that is a field K equipped with a quasi-ordering \preceq such that the following axioms are satisfied for all $x, y, z \in K$:

$$(Q0) \quad x \simeq 0 \Rightarrow x = 0,$$

$$(Q1) \quad x \preceq y \wedge 0 \preceq z \Rightarrow xz \preceq yz,$$

$$(Q2) \quad x \preceq y \wedge z \not\simeq y \Rightarrow x + z \preceq y + z.$$

Evidently, quasi-ordered fields arise from ordered fields by replacing anti-symmetry with the weaker axiom (Q0), and by weakening the compatibility with addition. The condition $z \not\simeq y$ in (Q2) relates to the fact that $v(y) \neq v(z)$ implies $v(y+z) = \min\{v(y), v(z)\}$ for any valuation v on K .

Theorem 1 (Fakhruddin's dichotomy [7, Theorem 2.1]). *Let K be a field and \preceq a binary relation on K . Then (K, \preceq) is a quasi-ordered field if and only if it is either an ordered field or there is a valuation v on K such that $x \preceq y \Leftrightarrow v(y) \leq v(x)$ for all $x, y \in K$.*

Note that \preceq is induced by a valuation v on K if and only if $0 < -1$. In that case v is unique up to equivalence of valuations, and we denote the quasi-ordering \preceq also by \preceq_v .

Remark 2. In [5], a *locality* on a field is, by definition, either an ordering or a valuation. Hence, by Fakhruddin's dichotomy, localities and quasi-orderings are, in fact, equivalent notions.

Baer and Krull exhibited another strong relation between orderings and valuations on fields [2, 8]: any ordering \leq on K induces a valuation on K , the so-called *natural valuation* of (K, \leq) . It is the valuation on K whose valuation ring is the convex hull of \mathbb{Z} in K , i.e. the smallest convex subring of K w.r.t. \leq . An ordering \leq and a valuation v on K are called *compatible*, if K_v is convex. We also say that v , respectively K_v , is convex. In [9], the authors generalised this notion to quasi-orderings: given a quasi-ordered field (K, \preceq) , a valuation v on K – or equivalently a quasi-ordering \preceq_v – is said to be *compatible with \preceq* , if the valuation ring K_v is \preceq -convex, or equivalently, if $0 \leq x \leq y \Rightarrow x \preceq_v y$ for all $x, y \in K$. Note that if \preceq is induced by a valuation w , then \preceq_v is compatible with \preceq_w if and only if v is a coarsening of w . For a further discussion on this subject we refer the interested reader to [9] and [10].

We conclude this section by recalling some basic facts about real closed fields. An ordered field (K, \leq) is said to be *real closed* if one of the following equivalent conditions is satisfied:

(RC1) \leq does not extend to any proper algebraic extension of K .

(RC2) (K, \leq) satisfies the following conditions:

- (i) any $x \in K$ with $0 \leq x$ has a square root in K ,

(ii) any polynomial $f \in K[X]$ of odd degree has a root in K .

Any ordered field (K, \leq) admits a *real closure*, i.e. an algebraic field extension which is real closed. It is unique up to isomorphism of ordered fields (cf. [1, Satz 8]).

3. Quasi-real closed fields

In the present section we introduce quasi-real closed fields, that way unifying algebraically closed fields and real closed fields. To this end, we impose (RC1) on quasi-ordered fields.

Definition 3. A quasi-ordered field (K, \leq) is called quasi-real closed, if \leq does not extend to any proper algebraic extension of K .

Lemma 4. Let K be a field. The following are equivalent:

- (1) K is algebraically closed,
- (2) (K, \leq_v) is quasi-real closed for some valuation v on K ,
- (3) (K, \leq_v) is quasi-real closed for any valuation v on K .

Proof. If K is algebraically closed, then K admits no proper algebraic extension, whence (K, v) is quasi-real closed for any valuation v on K . Consequently, (1) implies (3). Moreover, since any field admits a valuation, clearly (3) implies (2). Finally, suppose that (2) holds. Note that v extends to any field extension L of K by an immediate consequence of Chevalley's Extension Theorem [6, Theorems 3.1.1 and 3.1.2]. Since (K, v) is quasi-real closed, this means that K does not admit any proper algebraic extension, i.e. that K is algebraically closed. Thus, (2) implies (1). \square

Theorem 5. The following are equivalent for any quasi-ordered field (K, \leq) :

- (a) (K, \leq) is quasi-real closed.
- (b) (QRC1) either \leq is an ordering and (K, \leq) is a real closed field, or \leq is induced by a valuation and K is an algebraically closed field.
- (c) (QRC2) (K, \leq) satisfies the following conditions:
 - (i) any $x \in K$ with $0 \leq x$ has a square root in K .
 - (ii) any polynomial $f \in K[X]$ of odd degree has a root in K .

Proof. The equivalence of (a) and (b) is an immediate consequence of Fakhruddin's dichotomy (Theorem 1) and (RC1) (if \leq is an ordering), respectively Lemma 4 (if \leq is induced by a valuation).

It remains to show that (b) and (c) are equivalent. If \leq is an ordering, then this follows from (RC2). So let \leq be induced by some valuation. If (K, \leq) is quasi-real closed, then K is algebraically closed according to Lemma 4, whence (c) is fulfilled. Conversely, suppose that (c) holds. Since $K = \{0 \leq x\}$, we obtain by (i) that all elements in K have a square root in K . Along with condition (ii) this implies that K is algebraically closed according to [12, Theorem 2]. Hence, (K, \leq) is quasi-real closed again by Lemma 4. \square

Definition 6. Let (K, \leq) be a quasi-ordered field. A quasi-ordered field (L, \leq') is called a quasi-real closure of (K, \leq) , if

- (1) (L, \leq') is a quasi-real closed field,
- (2) $L|K$ is an algebraic field extension,
- (3) \leq' is an extension of \leq , i.e. $K \cap \leq' = \leq$.

Proposition 7. Any quasi-ordered field (K, \leq) admits a quasi-real closure. It is unique up to isomorphism of quasi-ordered fields.

Proof. Let us consider an algebraic closure \tilde{K} of K and the set

$$\{(L, \leq') : (L, \leq') \text{ is a quasi-ordered field, } K \subseteq L \subseteq \tilde{K}, K \cap \leq' = \leq\},$$

partially ordered by

$$(L_1, \leq'_1) \leq (L_2, \leq'_2) \iff L_1 \subseteq L_2 \text{ and } L_1 \cap \leq'_2 = \leq'_1.$$

By Zorn's lemma, there is a maximal quasi-ordered algebraic field extension (L, \leq') of (K, \leq) . Hence, (L, \leq') is quasi-real closed and a quasi-real closure of (K, \leq) .

The proof of the uniqueness relies on Fakhruddin's dichotomy and (QRC1). So let (K_1, \leq_1) and (K_2, \leq_2) be quasi-real closures of (K, \leq) . If \leq is an ordering, then (K_1, \leq_1) and (K_2, \leq_2) are real closures of (K, \leq) , whence they are order-isomorphic according to [1, Satz 8].

Likewise, if \leq is induced by some valuation on K , then K_1 and K_2 are algebraic closures of K . Hence, they are isomorphic as fields. Let $\leq_1 = \leq_{v_1}$ and $\leq_2 = \leq_{v_2}$. Then the corresponding valuation rings K_{v_1} and K_{v_2} are conjugated by an element of $\text{Gal}(\bar{K}|K)$ (cf. [6, Conjugation Theorem 3.2.15]). By composition, we obtain an isomorphism between (K_1, \leq_1) and (K_2, \leq_2) . \square

4. The theory of quasi-real closed fields and quantifier elimination

We conclude this note by proving that the theory of quasi-real closed fields with a non-trivial compatible valuation admits quantifier elimination. Moreover, we deduce that this theory is the model companion of the theory of quasi-ordered fields with a non-trivial compatible valuation. To this end, we exploit Fakhruddin's dichotomy and the fact that the theories of real closed valued fields and algebraically closed valued fields both admit quantifier elimination.

Theorem 8 (cf. [4, Theorems 4.4.2 and 4.5.1]).

- (1) *The theory of algebraically closed fields with a non-trivial valuation v admits quantifier elimination in the language of fields adjoined with \leq_v .*
- (2) *The theory of real closed fields with a non-trivial compatible valuation v admits quantifier elimination in the language of ordered fields adjoined with \leq_v .*

The characterization (QRC2) from Theorem 5 allows us to formalise quasi-real closed fields in a first order language. As language of quasi-ordered fields with a compatible valuation v , we fix $\mathcal{L} = \{+, \cdot, -, \leq, \leq_v, 0, 1\}$. Moreover, we denote by Σ_{QRC} the following set of \mathcal{L} -sentences:

- (1) the axioms stating that $(K, +, \cdot, -, 0, 1)$ is a field.
- (2) the axioms stating that (K, \leq) is a quasi-real closed field.
- (3) the axioms stating that (K, \leq_v) is a quasi-ordered field with v non trivial.
- (4) the following \mathcal{L} -sentences that determine the relationship of \leq and \leq_v :
 - (i) $\phi_1 \equiv 0 < -1 \rightarrow (\forall x, y: 0 \leq x \leq y \leftrightarrow x \leq_v y)$
 - (ii) $\phi_2 \equiv -1 < 0 \rightarrow (\forall x, y: 0 \leq x \leq y \rightarrow x \leq_v y)$

The \mathcal{L} -sentences ϕ_1 and ϕ_2 , respectively, state that $\leq = \leq_v$ if \leq is induced by a valuation, and that v is \leq -compatible if \leq is an ordering. That way, we have unified the languages of real closed valued fields and algebraically closed valued fields that are used in Theorem 8. We refer to the theory of Σ_{QRC} as the *theory of quasi-real closed fields with a non-trivial compatible valuation*. We also consider the *theory of quasi-ordered fields with a non-trivial compatible valuation*, denoted by Σ_{QO} , which consists of (1), (3), (4), and the axioms (Q0), (Q1) and (Q2) (see Section 2).

Theorem 9. *The theory of Σ_{QRC} admits quantifier elimination.*

Proof. Any model of Σ_{QRC} is either an algebraically closed field with a non-trivial valuation or a real closed field with a non-trivial compatible valuation (Theorem 5). Therefore, the result follows from the fact that each of these two classes admits quantifier elimination (Theorem 8). \square

Definition 10. *Let $\Sigma \subseteq \text{Sent}(\mathcal{L})$ for a given language \mathcal{L} . A set $\Sigma^* \subseteq \text{Sent}(\mathcal{L})$ is called a model companion of Σ , if*

- (1) *every model of Σ^* is a model of Σ ,*

- (2) every model of Σ can be extended to a model of Σ^* ,
 (3) Σ^* is model complete.

Corollary 11 (cf. [4, Corollaries 4.4.3 and 4.5.4]). *The theory of Σ_{QRC} is model complete. It is the model companion of the theory of Σ_{QO} .*

Proof. Property 10(1) is obviously satisfied. Model completeness follows immediately from the quantifier elimination that we obtained in Theorem 9. It remains to show that condition (2) of Definition 10 holds in our setting.

So let (K, \leq, \leq_ν) be some quasi-ordered field with a non-trivial compatible valuation, and let (L, \leq') be a quasi-real closure of (K, \leq) (Proposition 7). Then (L, \leq') is a quasi-real closed field. Moreover, $L|K$ is algebraic, whence [5, Theorem 20.1.1 (a)] yields a unique extension \leq'_ν of \leq_ν from K to L such that ν is compatible with \leq' . If \leq is a valuation (i.e. $\leq_\nu = \leq$), then the uniqueness tells us that also $\leq'_\nu = \leq'$. Furthermore, ν is non-trivial since ν is non-trivial. Hence, (L, \leq', \leq'_ν) is an extension of (K, \leq, \leq_ν) and a model of the theory of quasi-real closed fields with a non-trivial compatible valuation. \square

Remark 12. The theory of Σ_{QRC} is not complete. For example the \mathcal{L} -sentence $\exists x: x^2 + 1 = 0$ is false for any real closed field, but true for any algebraically closed field. Alternatively, this follows from the fact that for algebraically closed fields the characteristic is not fixed.

Open Question. Can we replace the \mathcal{L} -sentences ϕ_1 and ϕ_2 in Σ_{QRC} with the single \mathcal{L} -sentence $\phi \equiv \forall x, y: x \leq y \rightarrow x \leq_\nu y$? This is equivalent to the question, whether algebraically closed fields admit quantifier elimination in the language of fields adjoined by \leq_ν and \leq_w , where ν and w are valuations on K such that ν is coarser than w .

Acknowledgment

The authors wish to thank Salma Kuhlmann for initiating and supporting their collaboration, and also for helpful discussion on the subject. Without her, this paper would not have come to exist.

References

- [1] E. Artin, O. Schreier, "Algebraische Konstruktion reeller Körper", *Abh. Math. Semin. Univ. Hamb.* **5** (1927), no. 1, p. 85-99.
- [2] R. Baer, "Über nicht-archimedisch geordnete Körper. (Beiträge zur Algebra 5.).", 1927, 3-13 pages.
- [3] G. Cherlin, M. A. Dickmann, "Real closed rings II", *Ann. Pure Appl. Logic* **25** (1983), no. 3, p. 213-231.
- [4] C. N. Delzell, A. Prestel, *Mathematical logic and model theory. A brief introduction*, Universitext, Springer, 2011, x+193 pages.
- [5] I. Efrat, *Valuations, orderings, and Milnor K-theory*, Mathematical Surveys and Monographs, vol. 124, American Mathematical Society, 2006, xiv+288 pages.
- [6] A. J. Engler, A. Prestel, *Valued fields*, Springer Monographs in Mathematics, Springer, 2005, x+205 pages.
- [7] S. M. Fakhruddin, "Quasi-ordered fields", *J. Pure Appl. Algebra* **45** (1987), no. 3, p. 207-210.
- [8] W. Krull, "Allgemeine Bewertungstheorie", *J. Reine Angew. Math.* **167** (1932), p. 160-196.
- [9] S. Kuhlmann, M. Matusinski, F. Point, "The Valuation Difference Rank of a Quasi-Ordered Difference Field", in *Groups, Modules, and Model Theory – Surveys and Recent Developments : In Memory of Rüdiger Göbel* (M. Droste, L. Fuchs, B. Goldsmith, L. Strüngmann, eds.), Springer, 2017, p. 399-414.
- [10] S. Kuhlmann, S. Müller, "Compatibility of Quasi-Orderings and Valuations: A Baer–Krull Theorem for Quasi-Ordered Rings", *Order* **36** (2019), no. 2, p. 249-269.
- [11] A. Robinson, *Complete theories*, Studies in Logic and the Foundations of Mathematics, North-Holland, 1956.
- [12] J. Shipman, "Improving the fundamental theorem of algebra", *Math. Intell.* **29** (2007), no. 4, p. 9-14.
- [13] O. Zariski, P. Samuel, *Commutative algebra. Vol. II*, Graduate Texts in Mathematics, vol. 29, Springer, 1975, Reprint of the 1960 edition, x+414 pages.