

# Tannins and Their Complex Interaction with Different Organic Nitrogen Compounds and Enzymes: Old Paradigms versus Recent Advances

Bartosz Adamczyk,<sup>\*[a]</sup> Judy Simon,<sup>\*[b]</sup> Veikko Kitunen,<sup>[c]</sup> Sylwia Adamczyk,<sup>[c]</sup> and Aino Smolander<sup>[c]</sup>

Tannins, an abundant group of plant secondary compounds, raise interest in different fields of science, owing to their unique chemical characteristics. In chemical ecology, tannins play a crucial role in plant defense against pathogens, herbivores, and changing environmental conditions. In the food industry and in medicine, tannins are important because of their proven positive effect on human health and disease treatment. Such wide interests fueled studies on tannin chemistry, espe-

cially on their flagship ability to precipitate proteins. In this Review, we expand the basic knowledge on tannin chemistry to the newest insights from the field. We focus especially on tannin reactions with different non-protein organic N compounds, as well as the complex interactions of tannins with enzymes, resulting in either an increase or decrease in enzyme activity.

## 1. Introduction

Plants produce a wide variety of compounds to sustain and support growth, development, and reproduction, including secondary metabolites that are not essential for plant growth and, in contrast to primary metabolites, typically bear complex structures. The precise composition and chemical complexity of secondary metabolites became known only with significant improvements with regard to analytical techniques in the middle of the 20th century; particularly the development of chromatography.<sup>[1]</sup> Extractable plant secondary compounds constitute up to 30% of the dry weight of terrestrial plants, especially in forest ecosystems,<sup>[2]</sup> with their main role as defense against pathogens and herbivores,<sup>[3]</sup> but also as allelopathic agents,<sup>[4]</sup> antioxidants protecting leaves from UV radiation, and

excess of light<sup>[5]</sup> as well as regulators of nutrient and carbon cycling.<sup>[6]</sup>


Among the vast amount of plant secondary compounds, tannins quantitatively dominate, representing the fourth most abundant group of compounds in vascular plant tissue after cellulose, hemicelluloses, and lignin.<sup>[7]</sup> Plants can contain up to 20% of their dry weight in tannins,<sup>[7,8]</sup> the amount, however, changes in response to environmental conditions.<sup>[7]</sup> In turn, the effect of environmental stress, such as drought, on the production of tannins is complex and depends on further factors, for example the ontogenetic stage at which the drought stress occurs.<sup>[9]</sup> Additionally, many studies found high tannin concentrations in plants occurring in habitats with low soil fertility and low pH.<sup>[10]</sup> Moreover, it was shown that warming and altered precipitation can affect the chemistry of tannins by increasing their reactivity.<sup>[11]</sup> Chemically, tannins are often divided into two main groups: hydrolysable tannins (HTs) and condensed tannins (CTs) (Figure 1). Hydrolysable tannins can be separated into gallotannins and ellagitannins built up from gallic acid or hexahydroxydiphenic acid esters, respectively, linked to a sugar moiety (Figures 1 A and 1 B). Condensed tannins (proanthocyanidins) are polymers of three-ring flavonols joined through C–C bonds<sup>[12]</sup> (Figure 1 D). Monomers of CTs are divided into procyanidins and prodelfinidins (Figure 1 C). The newest findings point to a specific chloroplast-derived organelle called tannosome as the location of tannin production at the cellular level,<sup>[13,14]</sup> from which tannins are transported to vacuoles. Overall, the chemical structure of tannins is plant species-specific and shows a very high variability with probably no two species bearing the same tannin pattern,<sup>[15]</sup> thus, studying tannin chemistry can be very challenging. However, the problem of methodological development is not presented here.

[a] Dr. B. Adamczyk  
Department of Food and Environmental Sciences  
University of Helsinki  
PO Box 66, 00791 Helsinki (Finland)  
E-mail: Bartosz.Adamczyk@helsinki.fi

[b] Dr. J. Simon  
Ecology, Department of Biology  
University of Konstanz  
Universitätsstrasse 10, 78457 Konstanz (Germany)  
E-mail: judy.simon@uni-konstanz.de

[c] Dr. V. Kitunen, Dr. S. Adamczyk, Dr. A. Smolander  
Natural Resources Institute Finland  
PL 2, 00791 Helsinki (Finland)

 The ORCID identification number(s) for the author(s) of this article can be found under <https://doi.org/10.1002/open.201700113>.

 © 2017 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

In this Review, we focus on the recent novel insights into the chemistry of tannins, their interactions with other chemicals, and their influence on enzymatic activity. We challenge tannin chemistry paradigms with the newest findings to obtain a more holistic view on these plant secondary compounds. In Section 2, we evaluate the basic knowledge on tannin–protein interactions, adding the newest findings in the field. In Section 3, we challenge the traditional view on tannin chemistry, that is, that tannins are simply enzyme inhibitors. In Section 4, we expand the reactions of tannins to non-protein N compounds, underlining the remarkable versatility of tannin chemistry.


## 2. Interaction of Tannins with Proteins

According to Bate-Smith and Swain,<sup>[16]</sup> tannins are “water-soluble phenolic compounds, with a molecular mass between 0.5 and 3 kDa, able to precipitate proteins and alkaloids”. These tannin mass limitations (0.5–3 kDa) have been extended with time, as also lower and higher molecular mass polyphenolics are able to precipitate proteins;<sup>[17]</sup> however, the ability to form complexes with proteins is still a unique characteristic of tan-

Bartosz Adamczyk was born in Lodz (Poland) in 1979. He received a Master's degree in 2003 (Master of Biology) and PhD degree in 2009 (Doctor of Biology), both from the University of Lodz (Poland). After defending his PhD, he started as a Post-doc at The Finnish Forest Research Institute (Finland). In 2013, he obtained the title of docent (habilitation) from the University of Helsinki (Finland) and started to work there in 2015. His main interests span chemistry of plant secondary compounds, their role in boreal forest ecosystem, plant biochemistry and mitigation of climate change.



Judy Simon leads the Plant Interactions Ecophysiology Group at the University of Konstanz (Germany). After her studies in biology (RWTH Aachen, Germany), biogeography, soil science and geology (Saarland University, Germany), she conducted her PhD research at the University of Melbourne (Australia). She then worked as a Post-doctoral Fellow at the University of Freiburg (Germany), earning her Habilitation (postdoctoral qualification) in 2013. Since 2014, she conducts her research at the University of Konstanz on the influence of global change on plant interactions with regard to resource allocation strategies (i.e. different N acquisition strategies, N allocation to growth vs. defense) in woody species in boreal, temperate and tropical forest ecosystems.



**Figure 1.** Structure of tannins. A) simple gallotannin, B) simple ellagitannin, C) basic unit of condensed tannins, and D) condensed tannin trimer. Modified from Ref. [6].

nins,<sup>[16]</sup> for example, used already in ancient times to produce leather from animal skin. The reaction between tannins and proteins involves two stages: first the binding and second the aggregation, resulting in the formation of the precipitate.<sup>[18–20]</sup> Although earlier work on tannin–protein complexes pointed to non-covalent bonding and insoluble precipitates, more recent studies add covalent bonding and soluble complexes as a possible result of interactions between tannins and proteins.<sup>[16]</sup>

The formation of tannin–protein complexes depends on numerous factors dominated by tannin and protein chemistry (e.g. proline content), concentration, protein isoelectric point, pH, and ionic strength of the solution and also presence of other compounds in the solution.<sup>[21–25]</sup> The importance of the molecular complexity of tannins for a reaction with proteins was underlined by Haslam,<sup>[18]</sup> who first used structurally well-defined polyphenols and found that the most crucial features of tannins are phenolic sites crosslinked with proteins. Furthermore, proteins, which are especially prone to reactions with tannins, are proline-rich proteins (PRPs) found in mammalian saliva.<sup>[26]</sup> These interactions between PRPs and tannins protect dietary nitrogen from polyphenols, but also play a role in taste sensation known as astringency, a feeling of loss of lubrication and dryness.<sup>[19,27]</sup>

According to a well-known paradigm in tannin chemistry, precipitation of different proteins by tannins strictly depends on the protein isoelectric point ( $pI$ ).<sup>[21]</sup> At a pH close to the isoelectric point, proteins aggregate more eagerly because they carry no net electrical charge.<sup>[21]</sup> However, according to the newest findings, tannins can also form complexes with proteins at a pH far from their isoelectric point.<sup>[28,29]</sup> Bovine serum albumin (BSA) typically used in tannin–protein interaction studies with  $pI$  4.7 formed complexes with hydrolysable tan-


nins at neutral pH<sup>[28]</sup>, owing to tannin oxidative activity.<sup>[29]</sup> Although interactions between tannins and proteins have been intensively studied over the past 50 years, an in-depth understanding of all mechanisms regulating tannin–protein interactions is still lacking.

### 3. Specific Interaction of Tannins with Enzymes

As the majority of enzymes belong to proteins, it is widely believed that tannins decrease enzymatic activity as a result of enzyme complexation.<sup>[22,30–32]</sup> Although studies over the past decades have established tannins as potential inhibitors of enzymatic activity,<sup>[30,32,33]</sup> some studies found only a minor decrease in their activity.<sup>[34,35]</sup> Thus, our current understanding on their inhibiting role is still limited. Furthermore, the potential enhancement of enzyme activity by tannins has been overlooked for decades, with only very few exceptions. A study by Tagliacuzzi et al.<sup>[36]</sup> showed the ability of some phenolic compounds to enhance pepsin activity, which, however, was explained by phenolic-induced changes in the substrate protein.<sup>[37–39]</sup> Moreover, a highly heterogenic incubation study found that tannin-rich leaves in nylon-gauze bags in the rumen increased the activity of glutamate ammonia ligase, but no mechanical evidence was provided.<sup>[40]</sup> Only recently, evidence has been found that enzymatic activity is increased after the reaction with tannins present in low concentrations (Figure 2A).<sup>[41]</sup> This study showed that low concentrations of tannins increased the coiled structures of the enzymes, thereby boosting their catalytic activity.<sup>[41]</sup> High concentrations of tannins lead to opposite results by diminishing the catalytic activity (see Figure 2C), although even enzyme–tannin complexes exert some residual activity (Figure 2B).<sup>[41]</sup> The response of enzymes to tannins varied depending on the enzyme.<sup>[41]</sup> Overall, the interactions between tannins and enzymes follow the same rules as for tannins interacting with non-enzymatic proteins (see Section 2). However, enzymes vary in their affinity to tannins; thus, the potential influence of unknown tannins on a given enzyme is unpredictable. Recent findings in this section suggest that tannins are more than just inhibitors, but rather modifiers of enzyme activity, which should raise interest in different fields controlling enzymatic activity, such as food chemistry, medicine and industry.

### 4. Tannin Interactions with Organic Non-Protein N Compounds

According to the definition by Bate-Smith and Swain,<sup>[16]</sup> tannins form precipitates with proteins, but also with alkaloids. However, tannins also create complexes with metals,<sup>[16,43]</sup> and other compounds, i.e., tannic acid (TA), a common hydrolysable tannin, forms complexes with choline, an amine precursor of acetylcholine;<sup>[44]</sup> TA also adsorbs to chitosan.<sup>[45,46]</sup> It is widely assumed that tannins from the entire pool of organic N compounds precipitate only proteins/peptides.<sup>[42]</sup> However, only recently, it was shown that tannins can react with a wide set of different organic N compounds,<sup>[42]</sup> including arginine (from all amino acids), nitrogen bases, polyamines, chitin, and chito-



**Figure 2.** Influence of tannins on enzymatic activity of acid phosphatase: A) changes in enzymatic activity after addition of tannins in different concentrations, B) residual activity of enzymes after formation of complex with tannins, C) infrared spectra of enzyme secondary structure presented as a stacked plot of second derivative. Black lines represent enzyme without tannins and blue lines enzymes with low tannin concentrations. Red lines show enzymes with high tannin concentration. Spectra are smoothed by using eight points. Region of alpha-helix marked in yellow. Modified from Ref. [42].

san.<sup>[42]</sup> Similarly to tannin–protein reactions, the concentration, chemical structure, and pH of the solution seem to play a decisive role.<sup>[42]</sup> For example, the ability to form multiple hydrogen bonds<sup>[47]</sup> facilitates the formation of complexes with tannins. For proteinaceous amino acids, polyamines, and nitrogen bases, a higher reactivity towards tannins was found with higher molecular masses and more amine groups;<sup>[42]</sup> of all amino acids, arginine has the highest number of amine groups

**Table 1.** Reactivity of different organic N compounds towards tannins. Modified from Ref. [42].

Compound	Mw [Da]	N content [%]	Additional information (e.g. functional groups)	Reaction with tannins
<b>Amino acids</b>				
alanine	89	15.7	1 amine, 1 methyl, 1 carboxyl	–
arginine	174	32.1	4 amine, 1 carboxyl	+
asparagine	132	21.2	1 amide, 1 amine, 1 carboxyl	–
aspartic acid	133	10.5	1 amine, 2 carboxyl	–
cysteine	121	11.5	1 amine, 1 carboxyl, 1 thiol	–
glutamic acid	147	9.5	1 amine, 2 carboxyl	–
glutamine	146	19.1	1 amide, 1 amine, 1 carboxyl	–
glycine	75	18.6	1 amine, 1 carboxyl	–
histidine	155	27.0	1 imidazol, 1 amine, 1 carboxyl	–
isoleucine	131	10.6	1 amine, 1 carboxyl, 1 methyl	–
leucine	131	10.6	1 amine, 1 carboxyl, 1 methyl	–
lysine	146	19.1	2 amine, 1 carboxyl	–
methionine	149	9.3	1 amine, 1 carboxyl, 1 thiol	–
phenyl-alanine	165	8.4	1 amine, 1 carboxyl, 1 phenyl	–
proline	115	12.1	1 carboxyl, 1 pyrrolidine	–
serine	105	13.3	1 amine, 1 carboxyl, 1 hydroxyl	–
threonine	119	11.7	1 amine, 1 carboxyl, 1 hydroxyl, 1 methyl	–
tryptophan	204	13.7	1 amine, 1 carboxyl, 1 indole	–
tyrosine	181	7.7	1 amine, 1 carboxyl, 1 phenyl, 1 hydroxyl	–
valine	117	11.9	1 amine, 1 carboxyl, 2 methyl	–
<b>Polyamines</b>				
putrescine	88	31.8	2 amine	+
spermidine	145	28.9	3 amine	+
spermine	202	27.7	4 amine	+
<b>N bases</b>				
adenine	135	51.8	1 amine, 4 N in heterocyclic ring	+
cytosine	111	37.8	1 amine, 1 ketone, 2 N in heterocyclic ring	+
guanine	151	46.3	1 amine, 1 ketone, 4 N in heterocyclic ring	+
uracil	112	25.0	1 methyl, 2 ketone, 2 N in heterocyclic ring	+
thymine	126	22.2	1 methyl, 2 ketone, 2 N in heterocyclic ring	+
<b>Aminosugars</b>				
chitin	(203)n	6.89	2 amide, 4 hydroxylic, 2 methyl	+
chitosan	(161)n	8.69	1 amine, 2 hydroxyl	+
N-acetyl-D-glucosamine	221	6.3	1 amide, 4 hydroxyl, 1 methyl	–

(4) and almost the highest molecular mass (174 Da); for polyamines, spermine has the highest molecular mass (202 Da) and amount of amine groups (4) (see Table 1). For nitrogen bases, the two having no amine groups exerted the weakest reactivity towards tannins.<sup>[42]</sup> Thus, these findings on tannin–non-protein interactions lead us to further emphasize the importance of tannin chemistry. Moreover, reactions with numerous N compounds call for a change in our way of thinking about tannins: they can react with non-protein organic N compounds similarly to their reaction with proteins.

## 5. Conclusions and Perspectives

Interactions between tannins and proteins have been studied for more than 50 years, because of their unique characteristics and potential use in food industry and pharmacology. However, with the new insights regarding regulation of enzymes by tannin concentration and the potential interaction with other non-protein N compounds, future studies are needed. Special attention should be paid to the use of well-purified and char-

acterized tannins, because the chemistry of polyphenols and the presence of other compounds in plant extracts may significantly affect tannin interactions with N compounds. Follow-up studies should aim to extrapolate these results to more complex, heterogenic, realistic systems. In conclusion, studies investigating the interactions between tannins and proteins, but also other organic compounds, are likely to attract significant attention due to the general interest in polyphenols with regard to human health and disease treatment, but also their role in the beverage and food industry.

## Conflict of Interest

The authors declare no conflict of interest.

**Keywords:** organic nitrogen compounds · tannin chemistry · tannin–enzyme interactions · tannin–non-protein interactions · tannin–protein interactions

- [1] J. P. Salminen, M. Karonen, J. Sinkkonen, *Chem. Eur. J.* **2011**, *17*, 2806–2816.
- [2] W. Horwath, *Carbon Cycling: The Dynamics and Formation of Organic Matter in Soil Microbiology, Ecology and Biochemistry*, 4th ed. (Ed.: E. A. Paul), Academic Press, Boston, **2015**, pp. 339–382.
- [3] A. A. Agrawal, M. G. Weber, *Ecol. Lett.* **2015**, *18*, 985–991.
- [4] R. Croteau, T. M. Kutchan, N. G. Lewis *Natural Products (Secondary Metabolites) in Biochemistry & Molecular Biology of Plants*, **2000**, pp. 1250–1318.
- [5] D. C. Close, C. McArthur, *Oikos* **2002**, *99*, 166–172.
- [6] A. Smolander, S. Kanerva, B. Adamczyk, V. Kitunen, *Plant Soil* **2012**, *350*, 1–26.
- [7] T. E. C. Kraus, R. Dahlgren, R. J. Zasoski, *Plant Soil* **2003**, *256*, 41–66.
- [8] B. Adamczyk, A. Ahvenainen, O.-M. Sietiö, S. Kanerva, A.-J. Kieloaho, A. Smolander, V. Kitunen, P. Saranpää, T. Laakso, P. Strakova, et al., *Soil Biol. Biochem.* **2016**, *103*, 394–404.
- [9] C. S. Malisch, J.-P. Salminen, R. Kölliker, M. T. Engström, D. Suter, B. Studer, A. Lüscher, *J. Agric. Food Chem.* **2016**, *64*, 9307–9316.
- [10] R. R. Northup, R. A. Dahlgren, J. G. McColl, *Biogeochemistry* **1998**, *42*, 189–220.
- [11] N. Tharayil, V. Suseela, D. J. Triebwasser, C. M. Preston, P. D. Gerard, J. S. Dukes, *New Phytol.* **2011**, *191*, 132–145.
- [12] G. P. Bolwell, *BioEssays* **1990**, *12*, 453.
- [13] J.-M. Brillouet, C. Romieu, B. Schoefs, K. Solymosi, V. Cheynier, H. Fulcrand, J.-L. Verdeil, G. Conéjéro, *Ann. Bot.* **2013**, *112*, 1003–1014.
- [14] J. M. Brillouet, C. Romieu, M. Lartaud, E. Jublanc, L. Torregrosa, C. Caze-vieille, *Protoplasma* **2014**, *251*, 1387–1393.
- [15] J. P. Salminen, M. Karonen, *Funct. Ecol.* **2011**, *25*, 325–338.
- [16] A. E. Hagerman *Fifty Years of Polyphenol–Protein Complexes in Recent Adv. Polyphen. Res.*, Vol. 3, **2012**, pp. 71–97, and references therein.
- [17] C. Li, R. Leverence, J. D. Trombley, S. Xu, J. Yang, Y. Tian, J. D. Reed, A. E. Hagerman, *J. Agric. Food Chem.* **2010**, *58*, 9033–9042.
- [18] E. Haslam, *Biochem. J.* **1974**, *139*, 285–288.
- [19] E. Haslam, *Phytochemistry* **2007**, *68*, 2713–2721.
- [20] H. Kawamoto, F. Nakatsubo, K. Murakami, *Phytochemistry* **1995**, *40*, 1503–1505.
- [21] A. E. Hagerman, L. G. Butler, *J. Agric. Food Chem.* **1978**, *26*, 809–812.
- [22] N. Fierer, J. P. Schimel, R. G. Cates, J. Zou, *Soil Biol. Biochem.* **2001**, *33*, 1827–1839.
- [23] V. de Freitas, E. Carvalho, N. Mateus, *Food Chem.* **2003**, *81*, 503–509.
- [24] J. P. McManus, K. G. Davis, J. E. Beart, S. H. Gaffney, T. H. Lilley, E. Haslam, *J. Chem. Soc. Perkin Trans. 2* **1985**, 1429.
- [25] J. P. McManus, K. G. Davis, T. H. Lilley, E. Haslam, *J. Chem. Soc. Chem. Commun.* **1981**, 309–310.
- [26] A. E. Hagerman, L. G. Butler, *J. Biol. Chem.* **1981**, *256*, 4494–4497.
- [27] N. J. Murray, M. P. Williamson, T. H. Lilley, E. Haslam, *Eur. J. Biochem.* **1994**, *219*, 923–935.
- [28] B. Adamczyk, J.-P. Salminen, A. Smolander, V. Kitunen, *Int. J. Food Sci. Technol.* **2012**, *47*, 875–878.
- [29] M. T. Engström, X. Sun, M. P. Suber, M. Li, J.-P. Salminen, A. E. Hagerman, *J. Agric. Food Chem.* **2016**, *64*, 8994–9003.
- [30] C. B. Hall, *Nature* **1966**, *212*, 717–718.
- [31] J. P. Schimel, R. G. Cates, R. Ruess, *Biogeochemistry* **1998**, *42*, 221–234.
- [32] D. J. Triebwasser, N. Tharayil, C. M. Preston, P. D. Gerard, *New Phytol.* **2012**, *196*, 1122–1132.
- [33] G. D. Joannis, R. L. Bradley, C. M. Preston, A. D. Munson, *New Phytol.* **2007**, *175*, 535–546.
- [34] D. H. Strumeyer, M. J. Malin, *Biochem. J.* **1970**, *118*, 899–900.
- [35] M. R. Juntheikki, R. Julkunen-Tiitto, *J. Chem. Ecol.* **2000**, *26*, 1151–1165.
- [36] D. Tagliacozzi, E. Verzelloni, A. Conte, *J. Agric. Food Chem.* **2005**, *53*, 8706–8713.
- [37] S. Mole, P. G. Waterman, *J. Chem. Ecol.* **1985**, *11*, 1323–1332.
- [38] Z. Tantoush, D. Apostolovic, B. Kravic, I. Prodic, L. Mihajlovic, D. Stanic-Vucinic, T. Cirkovic Velickovic, *J. Funct. Foods* **2012**, *4*, 650–660.
- [39] F. Shen, F. Niu, J. Li, Y. Su, Y. Liu, Y. Yang, *Food Res. Int.* **2014**, *59*, 100–107.
- [40] H. P. Makkar, B. Singh, R. K. Dawra, *Br. J. Nutr.* **1988**, *60*, 287–296.
- [41] B. Adamczyk, M. Karonen, S. Adamczyk, M. T. Engström, T. Laakso, P. Saranpää, V. Kitunen, A. Smolander, J. Simon, *Soil Biol. Biochem.* **2017**, *107*, 60–67.
- [42] B. Adamczyk, S. Adamczyk, A. Smolander, V. Kitunen, *Soil Biol. Biochem.* **2011**, *43*, 628–637.
- [43] L. Zhang, R. Liu, B. W. Gung, S. Tindall, J. M. Gonzalez, J. J. Halvorson, A. E. Hagerman, *J. Agric. Food Chem.* **2016**, *64*, 3025–3033.
- [44] M. Kalina, D. C. Pease, *J. Cell Biol.* **1977**, *74*, 726–741.
- [45] M.-Y. Chang, R.-S. Juang, *J. Colloid Interface Sci.* **2004**, *278*, 18–25.
- [46] J.-H. An, S. Dultz, *Appl. Clay Sci.* **2007**, *36*, 256–264.
- [47] C. L. Borders, J. A. Broadwater, P. A. Bekeny, J. E. Salmon, A. S. Lee, A. M. Eldridge, V. B. Pett, *Protein Sci.* **1994**, *3*, 541–548.

---

Received: June 8, 2017

Version of record online August 16, 2017