

# Exploring local dependence

Klaus Abberger, University of Konstanz, Germany

**Abstract:** This paper discusses two graphical methods for the investigation of local association of two continuous random variables. Often, scalar dependence measures, such as correlation, cannot reflect the complex dependence structure of two variables. However, dependence graphs have the potential to assess a far richer class of bivariate dependence structures. The two graphical methods discussed in this article are the chi-plot and the local dependence map. After the introduction of these methods they are applied to different data sets. These data sets contain simulated data and daily stock return series. With these examples the application possibilities of the two local dependence graphs are shown.

**Keywords:** Association, bivariate distribution, chi-plot, copula, correlation, kernel smoothing, local dependence, permutation test

## 1 Introduction

The dependence between a pair of continuous random variables is often more complex than a single scalar dependence measure can reflect. Therefore, a global summary statistic such as the correlation coefficient will not convey the dependence structure. Local measures of dependence can be used instead. In this paper two graphical methods for analyzing dependence locally are discussed and compared. These two methods handle the dependence quite different. One graphical tool is the chi-plot introduced by Fisher and Switzer (1985, 2001). They utilize the chi-measure to receive a plot which is approximately horizontal under independence. Another idea was introduced by Hollander and Wang (1987) and is further developed in a series of subsequent papers by Jones (1996,1998) and Jones and Koch (2002). This concept is called local dependence function and is based on the mixed partial

derivatives of the logarithm of the bivariate density function. It is possible to estimate the local dependence function from data using, for example, kernel methods. The resulting estimates can then be graphed in one of the usual ways, such as by contour plots. But these figures are often not meaningful and hard to interpret. It can be argued that the local dependence function convey information that is too detailed to be easily interpretable. This fact motivates Jones and Koch (2002) to make local dependence a more interpretable tool, by introducing so called dependence maps. Via local permutation testing, dependence maps simplify the estimated local dependence structure between variables by identifying regions of (significant) positive, (nonsignificant) zero and (significant) negative local dependence.

The organization of this paper is as follows: The next section introduces and discusses the local dependence concepts. Section 3 includes some examples for simulated data and Section 4 presents estimations for some daily stock return series.

## 2 Two local dependence graphs

The chi-plot defined by Fisher and Switzer (1985, 2001) is designed to provide a graph that has characteristic patterns depending on whether the variates are independent, have some degree of monotone relationship, or have more complex dependence structure. The chi-plot depends on the data only through the values of their ranks.

Let  $(x_1, y_1), \dots, (x_n, y_n)$  be a random sample from  $H$ , the joint (continuous) distribution function for a pair of random variables  $(X, Y)$ , and let  $1(A)$  be the indicator function of the event  $A$ . For each data point  $(x_i, y_i)$ , set

$$H_i = \sum_{j \neq i} 1(x_j \leq x_i, y_j \leq y_i)/(n-1), \quad (1)$$

$$F_i = \sum_{j \neq i} 1(x_j \leq x_i)/(n-1), \quad (2)$$

$$G_i = \sum_{j \neq i} 1(y_j \leq y_i)/(n-1), \quad (3)$$

and

$$S_i = \{(F_i - 1/2)(G_i - 1/2)\}. \quad (4)$$

Now calculate

$$\chi_i = (H_i - F_i G_i) / \{F_i(1 - F_i)G_i(1 - G_i)\}^{1/2} \quad (5)$$

and

$$\lambda_i = 4S_i \max\{(F_i - \frac{1}{2})^2, (G_i - \frac{1}{2})^2\}. \quad (6)$$

The chi-plot is a scatterplot of the pairs  $(\lambda_i, \chi_i)$ .

At each sample point,  $\chi_i$  is actually a correlation coefficient between dichotomized  $X$  values and dichotomized  $Y$  values. Therefore, all values of  $\chi_i$  lie in the interval  $[-1, 1]$ . If  $Y$  is a strictly increasing function of  $X$ , then  $\chi_i = 1$  for all sample cut points, and when  $Y$  is a strictly decreasing function of  $X$ , then  $\chi_i = -1$  for all sample cut points. The value  $\lambda_i$  is a measure of the distance of the data point  $(x_i, y_i)$  from the center of the data set. All values  $\lambda_i$  must lie in the interval  $[-1, 1]$ . When the data are a random bivariate sample from independent continuous marginals, then the values of the  $\lambda_i$  are individually uniform distributed. However, when  $X$  and  $Y$  are associated, then the values of  $\lambda_i$  may show clustering. In particular, if  $X$  and  $Y$  are positively correlated,  $\lambda_i$  will tend to be positive and vice versa for negative correlation.

$\chi_i$  can be seen also as an empirical measure of the “positive quadrant dependence” (PQD). The PQD is defined as follows (Joe, 1997):

$$P(X \leq a_1, Y \leq a_2) \geq P(X \leq a_1) \cdot P(Y \leq a_2), \text{ for all } a_1, a_2 \in R. \quad (7)$$

At this PQD is defined globally for all  $a_1, a_2 \in R$ . On the other hand,  $\chi_i$  measures this dependence locally and draws it against the distance of the data point to the data center. Joe (1997) defines various dependence concepts and shows the relations between them. Is  $(X; Y)$  for example (positively) “associated”, i.e.

$$E[g_1(X)g_2(Y)] \geq E[g_1(X)] \cdot E[g_2(Y)], \quad (8)$$

for all real valued functions  $g_1, g_2$  which are increasing and are such that the expectations exist, then is  $(X; Y)$  also PQD. Of course (8) contains the case of positive correlation.

As a second local dependence concept the “local dependence function” and the “local dependence map” are discussed. The local dependence function was introduced by Holland and Wang (1987). It is a generalization of

the cross-product ratio for the case of continuous random variables  $X$  and  $Y$ . If  $f$  is the bivariate density function, then the local dependence function  $\gamma$  is given by

$$\gamma(x, y) = \frac{\delta^2 \log f(x, y)}{\delta x \delta y}. \quad (9)$$

Jones (1996) motivated  $\gamma$  from the point of view of localizing the Pearson correlation coefficient  $\rho$ . The local dependence function is constant for the bivariate normal distribution, taking the value  $\rho/(1 - \rho^2)$  in the standard normal case. Jones (1998) examined for which general class of densities the local dependence function is constant. His considerations provide the following result: The local dependence function is constant if and only if  $Y|x$  has a linear exponential family distribution, for fixed dispersion parameter, with canonical parameter a linear function of  $x$ , i.e. the common generalized linear model situation.

The local dependence function can be nonparametrically estimated with the help of kernel methods. The local adaption of a bilinear form  $a + bx + cy + dxy$  to the log density using the kernel weighted local likelihood

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n K_{h_1}(X_i - x) K_{h_2}(Y_i - y) \log f(X_i, Y_i) \\ - \int \int K_{h_1}(X - x) K_{h_2}(Y - y) f(X, Y) dX dY \end{aligned} \quad (10)$$

is used here. This approach is treated in detail in Loader (1996). He delivers in addition with the S-Plus (resp. R) function *locfit()* a program for the calculation of estimates.

The resulting estimates can be graphed in one of the usual ways, such as by a contour plot. The results of the estimate can be hard to interpret sometimes. It can be argued that the estimated local dependence function convey information that is too detailed to be easily interpretable. For this reason Jones and Koch (2002) developed the local dependence map which is more simply to read. Via local permutation testing, dependence maps simplify the estimated local dependence structure between two variables by identifying regions of (significant) positive, (nonsignificant) zero and (significant) negative local dependence.

The construction of the local dependence map can be summarized as follows: The first step is to estimate the density  $f$  using the kernel estimator

defined in (10). The estimated density is used to identify points at which  $f$  appears to be small. These points are excluded from the dependence map because estimation of  $\gamma$ , which is a second derivative quantity is extremely unreliable there. Here and in later calculations the tricube kernel

$$K(u) = (1 - |u|^3)^3, |u| < 1, \quad (11)$$

which is the default weight function in *locfit()* is used. The bandwidths are chosen by a rule of thumb (Jones and Koch, 2002). They are calculated by

$$h_i = \frac{s_i}{n^{1/6}} \left\{ \frac{2\sqrt{\pi} \int K^2(u) du}{\int u^2 K(u) du} \right\}^{1/3} \frac{(1 - r^2)^{5/12}}{(1 + r^2/2)^{1/6}}, i = 1, 2. \quad (12)$$

where  $s_i$  is the sample standard deviation and  $r$  is the sample correlation.

Now, a regular grid is placed across the area of interest. At each grid point  $(x_i, y_i)$ ,  $\hat{\gamma}(x_i, y_i)$  can be calculated using *locfit()*. Jones and Koch (2002) suggest using the same bandwidth as above in (12). Though, for estimating  $\gamma$  itself, one would normally expect that  $h_i$  is inadequate and that larger bandwidths are necessary. Jones and Koch (2002) argue, that one can nevertheless use the above bandwidths. However,  $h_i$  is only used as lower bound and the bandwidth for estimating  $\gamma$  is adjusted by eye in this article. The next step is the execution of local permutation tests of the null hypothesis that  $\gamma(x_i, y_i) = 0$ . Therefore, the data within the kernel around  $(x_i, y_i)$  are used. Samples satisfying the null hypothesis can readily be generated if the  $Y$ 's are randomly permuted to break their ties to the  $X$ 's. This is done  $P$  times and  $\hat{\gamma}_p$  is computed for each permuted data set,  $p = 1, 2, \dots, P$ . Now, the local dependence map value is “+“ if the observed value is in the highest  $(\alpha/2)\%$  of simulated  $\hat{\gamma}_p$ 's, is defined to be “-“ if the observed value is in the lowest  $(\alpha/2)\%$  of simulated  $\hat{\gamma}_p$ 's, and set to be “0“ otherwise. In the examples still following we use  $P = 500$  and  $\alpha = 0.1$ .

To be able to judge the differences in the introduced methods, some examples with simulated data are presented in the next section.

### 3 Examples with simulated data

The various methods introduced above are applied to different data sets with a known underlying structure. Therefore their advantages and disadvantages can be easily recognized. First, a synthetic data set of size  $n = 500$ , where

$Y$  and  $X$  have a joint standard normal distribution with correlation  $\rho = 0.5$  is used. Nothing surprising can be recognized at the local dependence map. Positive dependence is indicated over almost the complete range looked at. The chi-plot shows a bent course. The chi-values are near to zero on the edges of the graph. For easier interpretation the mapping is extended by two horizontal lines at  $\pm c/n^{1/2}$ , where  $c$  is selected so that approximately 95% of the pairs  $(\lambda_i, \chi_i)$  lie between the lines.  $c$  can be calculated with Monte Carlo methods (Fisher and Switzer, 2001).

The bent course of the chi-plots reports the so-called “tail independence“ characteristic of the bivariate normal distribution, provided that  $|\rho| < 1$ . Tail dependence is often defined in terms of the copula of a joint distribution. For a complete treatment of copula see Joe (1997). For a bivariate distribution  $F(x, y)$  with  $j$ -th univariate margin  $F_j$ , the copula associated with  $F$  is a distribution function  $C : [0, 1]^2 \rightarrow [0, 1]$  that satisfies

$$F(x, y) = C(F_1(x), F_2(y)), x, y \in R. \quad (13)$$

So the copula is a distribution function of a random vector,  $\mathbf{U} = (U_1, U_2)$ , where each  $U_j \sim \text{uniform}(0, 1)$ . If a bivariate copula  $C$  is such that

$$\lim_{u \rightarrow 0} C(u, u)/u = \lambda_L \quad (14)$$

exists,  $C$  has lower tail dependence if  $\lambda_L \in (0, 1]$  and no lower tail dependence if  $\lambda_L = 0$ . Similarly the upper tail dependence can be defined (see Joe, 1997).

The chi-value is zero if  $P(X \leq u_1, Y \leq u_2) = P(X \leq u_1) \cdot P(Y \leq u_2)$ . This is equal to

$$C(u_1, u_2) = u_1 \cdot u_2 \quad (15)$$

and

$$\frac{C(u_1, u_2)}{u_1} = u_2 \text{ resp. } \frac{C(u_1, u_2)}{u_2} = u_1. \quad (16)$$

If the copula is lower tail independent, we know that  $\lim_{u \rightarrow 0} C(u, u)/u = 0$  and so the chi-values are also equal to zero. For the normal distribution it is known that it is not tail dependent (Embrechts et al., 2002) and the chi-plot shows exactly this.

The bivariate t-distribution provides an interesting contrast to the bivariate normal distribution, provided  $\rho > -1$  the copula of the bivariate t-distribution is tail dependent (see also Embrechts et al., 2002).




Figure 1: Scatterplot, chi-plot, density estimation and local dependence map for normal distributed data




Figure 2: Scatterplot, chi-plot, density estimation and local dependence map for t-distributed data




Figure 3: Scatterplot, chi-plot, density estimation and local dependence map for Cauchy distributed data

The data underlying the Figure 2 are from a t-distribution with three degrees of freedom. For the data generation the following qualities are used. At first bivariate standard normal distributed samples  $\mathbf{Z}$  with correlation 0.5 are produced. A  $\chi^2$  distributed variable  $U$  is then produced with  $df = 3$  degrees of freedom. With  $\mathbf{V} = (V_1, V_2)'$  and  $V_i = \sqrt{df}Z_i/\sqrt{U}$ ,  $i = 1, 2$ , the variable  $\mathbf{V}$  is then a sample from the t-distribution with three degrees of freedom and

$$cov(\mathbf{V}) = \frac{df}{df - 2}\Sigma, \quad (17)$$

$\Sigma$  denoting the variance-covariance matrix of the bivariate normal variables.

In comparison with the normal distributed data the chi-plot for the t-distributed data shows another course. The chi-plot does not incline on the right side of the graphic to the zero line anymore. This shows tail dependence on the upper right and lower left edge of the data. The local dependence map looks similar to the map for the normal distributed data. This changes in the next data example.

The data in Figure 3 are generated from a Cauchy distribution, in the same way as the t-distributed data. The degrees of freedom are set to one and the correlation of the bivariate standard normal is set to zero. The chi-plot in Figure 3 nevertheless shows dependence. Due to the definition of  $\lambda$  in (6), negative chi-values for negative  $\lambda$  mean negative dependence in the upper left and lower right corner of the data. On the other hand, the positive chi-values for positive  $\lambda$  indicate positive dependence in the lower left and upper right corner. Looking at the local dependence map, one recognizes also four fields where dependencies are indicated. This confirms Holland and Wang's (1987) observation that  $\gamma(x, y)$  for the bivariate Cauchy density "changes sign as  $(x, y)$  changes quadrant in the plane. Random variables  $X$  and  $Y$  are positively associated in the first and third quadrants and negatively associated in the second and fourth". Exactly these signs are also shown in the local dependence map. This can be interpreted as positive association between  $|X|$  and  $|Y|$ . The example shows, that the local dependence map can be used to indicate dependence when the direction of the association is different in different regions of the plane.

## 4 Applications to stock return series

The local dependence graphs are used now to examine the “autocorrelation“ of two daily stock market index return series. The series are the German DAX<sup>1</sup> index from January 1, 1990 to Dezember 30, 1998 and the S&P 500 from January 1, 1992 until July 19, 1999. The returns  $r_t$  are defined by

$$r_t = \log \frac{p_t}{p_{t-1}}, \quad (18)$$

with  $p_t$  the price of the index at time  $t$ . The Figures 4 and 5 contain scatterplots of  $r_t$  against  $r_{t-1}$  for the respective return series. For both indexes the chi-plots show dependencies. There are no dependencies in the center of the bivariate distributions but on the edges. So there seem to be tail dependencies in the lagged returns. Also in the local dependence maps spots with positive and negative dependence are recognized. The arrangement of the spots confirm the well known fact, that the lagged absolute returns are positively correlated. Both of the dependence graphs also show that the Bravais-Pearson correlation coefficient is not suitable to include the complex dependence structure in the lagged returns. The chi-plot shows in addition that when a model is developed for these data the tail dependence behavior has to be taken into account.

## 5 Summary

In the above sections two methods for local analysis of dependence were presented and illustrated at data examples. The first method, the chi-plot, is simple to calculate and is well suited to recognize dependence in the tails of the distribution. The second method, the local dependence map, is technically more demanding. It requires the estimation of mixed partial derivatives which is realized by nonparametric estimation methods. This also means that bandwidths must be chosen. After this the local dependence function must be converted with local permutation tests in the local dependence map. For this however regions can be indicated for the dependencies which are easy to interpret. Since regions with small densities are taken out of the calculations, the extreme tails of the distribution can not be analyzed. In summary, the two graphs are supplementary. The local dependence map delivers a mapping which is easy to interpret and the chi-plot permits representations of the dependencies up to the tails of the distribution.

---

<sup>1</sup>The data are provided by the Deutsche Finanzmarkt Datenbank Karlsruhe




Figure 4: Scatterplot, chi-plot, density estimation and local dependence map for lagged daily DAX returns




Figure 5: Scatterplot, chi-plot, density estimation and local dependence map for lagged daily S&P 500 returns

## 6 Literature

**Embrechts P., McNeil A., Straumann D. (2002):** Correlation and dependence in risk management: properties and pitfalls. In: Risk Management: Value at Risk and Beyond, ed.: M.A.H. Dempster, Cambridge University Press, Cambridge.

**Fisher N.I., Switzer P. (2001):** Graphical assessment of dependence: is a picture worth 100 tests? *The American Statistician*, 55, 233-239.

**Fisher N.I., Switzer P. (1985):** Chi-plots for assessing dependence. *Biometrika*, 72, 253-265.

**Hollander P.W., Wang Y.J. (1987):** Dependence function for bivariate continuous densities. *Communications in Statistics, Theory and Methods*, 16, 863-876.

**Joe H. (1997):** Multivariate models and dependence concepts. Chapman and Hall, New York.

**Jones M.C. (1998):** Constant local dependence. *Journal of Multivariate Analysis*, 64, 148-155.

**Jones M.C. (1996):** The local dependence function. *Biometrika*, 83, 899-904.

**Jones M.C., Koch I. (2002):** Dependence maps: local dependence in practice. Discussion paper.

**Loader C. (1999):** Local regression and likelihood. Springer Verlag, Heidelberg.