

## Molecular techniques and their limitations shape our view of the holobiont

Ira Cooke<sup>a,b,\*</sup>, Oliver Mead<sup>c,d</sup>, Casey Whalen<sup>a,b,e</sup>, Chloë Boote<sup>a,b,e</sup>, Aurelie Moya<sup>a,b,e</sup>, Hua Ying<sup>d</sup>, Steven Robbins<sup>f</sup>, Jan M. Strugnell<sup>b,g,h</sup>, Aaron Darling<sup>i</sup>, David Miller<sup>a,b,e</sup>, Christian R. Voolstra<sup>j</sup>, Maja Adamska<sup>c,d</sup>, Consortium of Australian Academy of Science Boden Research Conference Participants<sup>1</sup>

<sup>a</sup> Department of Molecular and Cell Biology, James Cook University, Townsville, QLD, 4811, Australia

<sup>b</sup> Centre for Tropical Bioinformatics and Molecular Biology, James Cook University, Townsville, QLD, 4811, Australia

<sup>c</sup> ARC Centre of Excellence for Coral Reef Studies, Australian National University, Canberra, ACT, 2601, Australia

<sup>d</sup> Research School of Biology, Australian National University, Canberra, ACT, 2601, Australia

<sup>e</sup> ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, QLD, 4811, Australia

<sup>f</sup> Australian Center for Ecogenomics, University of Queensland, St. Lucia, QLD, 4072, Australia

<sup>g</sup> Centre of Sustainable Tropical Fisheries and Aquaculture, James Cook University, Townsville, 4810, QLD, Australia

<sup>h</sup> Department of Ecology, Environment and Evolution, School of Life Sciences, La Trobe University, Melbourne, 3083, Australia

<sup>i</sup> The iThree institute, University of Technology Sydney, Ultimo, NSW, 2007, Australia

<sup>j</sup> Department of Biology, University of Konstanz, Konstanz, 78457, Germany

### ABSTRACT

#### Keywords:

Holobiont  
Imaging  
Metagenomic sequencing  
Evolutionary inference  
Model system  
Multi-omics

It is now recognised that the biology of almost any organism cannot be fully understood without recognising the existence and potential functional importance of associated microbes. Arguably, the emergence of this holistic viewpoint may never have occurred without the development of a crucial molecular technique, 16S rDNA amplicon sequencing, which allowed microbial communities to be easily profiled across a broad range of contexts. A diverse array of molecular techniques are now used to profile microbial communities, infer their evolutionary histories, visualise them in host tissues, and measure their molecular activity. In this review, we examine each of these categories of measurement and inference with a focus on the questions they make tractable, and the degree to which their capabilities and limitations shape our view of the holobiont.

### 1. Introduction

Most, if not all, animals and plants exist as part of complex multi-organismal assemblages (Bosch and McFall-Ngai, 2011; Bosch and Miller, 2016; Bang et al., 2018) comprised of the host and associated microorganisms. These associations, together called holobionts or metaorganisms (for a differentiation between both terms, see (Jaspers et al., 2019)) may include members that interact in a manner that contributes to the fitness of the whole as well as others without any identifiable functional impact. Adopting a holistic view of biology that acknowledges the existence of holobionts and metaorganisms raises fundamental questions specific to the multi-organismal condition. One such question is whether a core microbiome or a set of core microbiota exists (Qin et al., 2010; Hernandez-Agreda et al., 2017; Ziegler et al., 2019) and whether this should be defined by taxonomic or metabolic composition (Cho and Blaser, 2012). More mechanistic questions, such

as whether microbial composition is influenced by the host (Rawls et al., 2006; Fraune and Bosch, 2007; Franzenburg et al., 2013; Augustin et al., 2017) and the degree to which partner organisms are interdependent (Russell et al., 2013; Fraune et al., 2015) are being addressed through experiments in model metaorganisms such as *Hydra*, zebrafish and pea aphids. Finally, the close association and potential interactions between organisms gives rise to a suite of evolutionary questions, such as whether key microbial taxa have co-evolved with their hosts (Baumann et al., 1997; Pollock et al., 2018; O'Brien et al., 2019), how the microbiome has influenced speciation and emergence of novel traits in hosts (Brucker and Bordenstein, 2012), and how metabolic cooperation has shaped partner genomes (Russell et al., 2013).

These questions are challenging to answer, but advances in molecular techniques are rapidly making them tractable. Perhaps the most notable advance from the past two decades has been the development and widespread adoption of marker-gene targeted amplicon

\* Corresponding author at: Department of Molecular and Cell Biology, James Cook University, Townsville, QLD, 4811, Australia.

E-mail address: [ira.cooke@jcu.edu.au](mailto:ira.cooke@jcu.edu.au) (I. Cooke).

<sup>1</sup> Detailed in Acknowledgements.

sequencing. This has revolutionised detection, classification, and quantification of microorganisms, making it possible to rapidly and cheaply profile the microbiota of a holobiont. The adaptability of this technique has allowed it to be employed in answering questions related to the variability of microbial communities across a wide range of species, tissues, and environmental conditions. In contrast, many molecular techniques form part of a toolbox that is closely tied to a particular organism or model system. Several such model systems are being developed for use in metaorganism research with toolboxes that include the removal or manipulation of important microbes (Fraune and Bosch, 2007; Voolstra, 2013; Moran and Yun, 2015), the ability to alter gene expression or edit genomes (Dunn et al., 2007; Franzenburg et al., 2013; Ikmi et al., 2014) and the ability to monitor metabolic interactions between components (Hillyer et al., 2017; Rädicker et al., 2018). Experiments in these model systems provide insights that are otherwise intractable in non-model systems and which form a framework for understanding inter-organismal interactions. Extending this framework to understand the biology of non-model metaorganisms is now a key focus for research on an increasingly broad range of taxa including keystone marine species such as corals, sponges, and reef fishes, agricultural and aquacultural species, and plants.

Metaorganism research is now heavily dependent on molecular techniques and its direction is arguably shaped by their availability, limitations, and advancement. In this essay we explore the role of four core categories of molecular technique in shaping the questions we can address in metaorganism and holobiont research. For each, we describe the related set of questions for which it is most suitably applied (Table 1) and outline the challenges that must be overcome in order to obtain clear answers. Where possible, we also reflect on opportunities afforded by new technologies and observe that these sometimes cut across categories. While our review reflects current research trends, it is likely that as new techniques develop they will blur the lines between those that exist today.

## 2. Identification and quantification of microbial communities

The ability to identify, taxonomically classify and quantify the abundance of microorganisms or their gene products, is a foundational requirement of metaorganism research. It is the basic measurement that allows us to ask “what are the members?” and “what are they likely to be doing at the molecular level?”. This, in turn, leads to consideration of where to draw the boundaries of an individual association and how

holobiont membership varies with time, environment and host genotype.

Answering these questions requires tools that can comprehensively assay microbial communities from within different holobiont assemblages and their surrounding environments. Molecular techniques based on next-generation sequencing are potentially able to accomplish this because they require only that DNA or RNA be extracted from the sample. These techniques have undergone rapid development over the past few decades and have now largely replaced microbial culture for surveying microbial communities, although the latter remains important for phenotypic characterisation, reference genome sequencing and taxonomic classification (Yarza et al., 2014).

Current techniques for sequence-based microbial profiling can be broadly categorised into amplicon-based and whole metagenome methods. Amplicon-based approaches identify organisms based on the amplification of a marker gene sequence (Woese and Fox, 1977; Yarza et al., 2014) chosen to provide phylogenetic signal across a wide range of taxa, whereas whole metagenome approaches sequence in an untargeted manner across all of the DNA present. Alternatively, RNA sequencing (metatranscriptomics, (Abu-Ali et al., 2018)) or mass spectrometry based proteomics (metaproteomics, (Starr et al., 2018)) can be used, especially where metabolic activities rather than taxonomic composition are of primary interest (see Section 4).

Currently, all methods for surveying microbial composition, including traditional culture-based methods as well as amplicon and whole metagenome sequencing, are subject to biases and limitations which contribute significantly toward shaping our view of the holobiont. DNA sequencing based techniques generally provide less biased estimates of abundance than culture-based methods but still show strong biases due to differences in DNA extraction efficiency (Xue et al., 2018). Amplicon based methods introduce additional bias related to the use of PCR including the primer set (Klindworth et al., 2013), target region, GC content, and input DNA concentration (Kennedy et al., 2014; Brooks et al., 2015; Laursen et al., 2017; Rintala et al., 2017).

In addition to bias, another issue with sequencing-based microbial profiling techniques is that they produce compositional data, that is, abundance measurements as a percentage of a (potentially unknown) total. It is essential that this is accounted for when choosing statistical methods for data analysis (Lovén et al., 2012; Gloor et al., 2017). More fundamentally, the total abundance of microorganisms provides crucial context required to infer ecological scenarios (e.g. competitive exclusion, outgrowth or differential survival) that give rise to changes in

**Table 1**

Important and emerging molecular techniques in holobiont research categorised according to the type of measurement or inference they provide. (FACS: Fluorescence Activated Cell Sorting; qPCR: Quantitative Polymerase Chain Reaction; SIMS: Secondary Ion Mass Spectrometry; GC-MS Gas Chromatography Mass Spectrometry).

Measurement or Inference	Techniques	Related Question(s)
Identification and quantification of microbial communities	Relative microbial community profiling using amplicon or metagenome sequencing Quantification using qPCR, Fluorescent staining, Gold labeling, FACS counts	Is there a core microbiota?  How does the taxonomic composition and metabolic potential of the microbiome vary with factors of interest? What is the bacterial load/cargo of host organisms per unit of tissue/organ/compartment?
Evolutionary inference	Phylogenetics Comparative genomics	Has co-speciation occurred? Do genomic changes reflect metabolic interdependence? Have certain genes been subject to strong selection? Is there evidence of genetic drift in partner genomes?
Measure molecular activity	Profile gene expression with RNASeq or Mass Spectrometry Profile metabolite production with GC-MS Experimental manipulation of microbiota and/or gene expression	What are the benefits and costs of the association to different partners?  Does nutrient exchange occur between partners? Do partners exert molecular control over each other?
Map (visualise, identify, quantify) microbiome components in host tissues	Dissection FISH FISH-CLEM SIMS	Is there a defined spatial organization/structure of the microbiome? Is the distribution of microbes affected by the host and/or environmental conditions?

composition (relative abundance) (Props et al., 2017). Measurements that are sensitive to total microbe abundance are also important in the verification of xenobiotic models and in determining the difference in microbial load between host tissues versus the environment, an important indicator of host selectivity or filtering. Techniques for measuring abundance in absolute rather than relative terms such as qPCR based assays (Jian et al., 2018), flow cytometry (Props et al., 2017) and fluorescence in situ hybridisation (Daims et al., 2001) will help to fill this gap.

Another suite of issues arises from the fact that most amplicon-based studies to date have relied on short (100-300 base pair [bp]) reads. One consequence of this is that it is often difficult to correctly separate reads arising from related taxa, which is important as it underpins estimates of microbial diversity, and because variation in the abundance of microbes that differ at the species or strain level may be related to functionally or ecologically significant differences in the host (Moeller et al., 2016; Neave et al., 2017b; Pollock et al., 2018). For whole metagenome approaches this process (called binning) is especially challenging because reads can originate from different genomic contexts as well as different taxa. The fixed genomic context used by amplicon sequencing means that differences between reads should arise only due to taxonomic differences and sequencing artifacts (e.g. single-base read errors or chimeras). Until recently many studies were based on the concept of OTUs (Operational Taxonomic Units), which are effectively clusters of marker gene sequences at a 97% similarity cutoff. Newer approaches explicitly model the difference between sequencing errors and biological variation to infer ASVs (amplicon sequence variants) representing biologically distinct sequences within the original sample (Callahan et al., 2016).

It is also important that biologically distinct sequences within the sample are correctly assigned to an agreed taxonomy. This provides a framework for communication, facilitates comparison between experiments and may also allow inferences to be made on the basis of homology. The most complete taxonomies are currently based on curated efforts to reconcile phylogenetic analyses of full length amplicon sequences with taxonomic assignments based on cultured strains (Yarza et al., 2014; Yilmaz et al., 2014). In principle, taxonomic assignments based on amplicon sequencing should be well placed to benefit from the relative completeness of these databases but since many studies only sequence a small region rather than the complete marker gene they are still prone to misclassification errors (Yarza et al., 2014). A recent development that promises to address this issue is the use of long read amplicon sequencing combined with software that models the error modes of long read sequencing technologies (Callahan et al., 2018). Notably, even with such sophisticated approaches, the challenge remains as how to denote distinct 'species' based on sequence diversity cutoffs (Konstantinidis Konstantinos et al., 2006).

In principle, identification and quantification of microbes based on whole metagenome sequencing avoids many of the problems with the amplicon based approach. Not only is it less prone to biases due to PCR amplification, but it also sequences a far greater proportion of the genome that could be used for gene content analysis and more precise discrimination between taxa (Shakya et al., 2013; Neave et al., 2017a; Jain et al., 2018). In practice, this approach is currently limited by the availability of whole genome reference sequences (see <http://gtadb.ecogenomic.org/> for an example of efforts to tackle this problem), high sequencing costs due to the fact that host DNA often dominates holobiont-derived samples, and high bioinformatic costs due to the far greater complexity and volume of data that must be analysed. Many of the challenges with whole metagenome sequencing data arise from the fact that raw sequencing reads are difficult to separate according to their taxon of origin (Lindgreen et al., 2016), a process that is much more complex for whole metagenome than for amplicon data. This reduces the accuracy of diversity profiles and effectively prevents assembly of complete genome sequences from mixed microbial data. Several new technologies and associated computational approaches are

emerging as potential solutions to this issue.

A particularly promising approach is chromosome conformation capture (3C) and, specifically, the Hi-C method for preparing sequencing libraries (Liu and Darling, 2015). This technique captures information about the physical proximity of fragments of DNA, which can later be used to infer their co-location on a chromosome or within a cell (Liu and Darling, 2015), thereby improving the accuracy of metagenomic binning and assembly. Other new technologies, such as single-cell metagenomic sequencing (Xu and Zhao, 2018) and long read sequencing (Arumugam et al., 2018; Bertrand et al., 2018), offer alternative solutions to this issue (Nicholls et al., 2018). None of these new technologies are free from challenges (see (Liu and Darling, 2015) for a brief overview) but the rapid advances in this area suggest that it may soon be possible to accurately assemble microbial genomes directly from metagenomic sequencing data. This would not only allow for the accurate assessment of microbial communities in terms of metabolic capacity (via gene content), it would also provide the dense taxonomic sampling of bacterial genomes required to replace amplicon-based taxonomic classification with a whole genome-based system.

### 3. Evolutionary inference

Molecular sequence data from holobionts encodes rich information about the evolutionary forces that have shaped the association. Inferences based on these data can answer questions such as the extent to which co-evolution has occurred and for which taxa this is true. In addition, genes or other genomic features that have been shaped by co-evolution may be identified, which can provide insights into the mechanisms underpinning long standing interactions between taxa.

Molecular signatures indicative of co-evolution (O'Brien et al., 2019) include phylosymbiosis (Sanders et al., 2014; Brooks et al., 2016), codivergence and the existence of genomic changes such as metabolic complementarity (Russell et al., 2013; Poulsen et al., 2014). In phylosymbiosis, the host phylogeny is correlated with a divergence pattern based on the microbiome profile distances (e.g. from amplicon or whole genome sequencing). Phylosymbiosis is frequently (Sanders et al., 2014; Brooks et al., 2016; Pollock et al., 2018) but not universally (Chandler et al., 2011; Kelley and Dobler, 2011) observed in holobiont systems and is not necessarily an indicator of co-evolution since it can also arise purely due to differential dispersal and establishment (filtering) of microbes in response to phylogenetically or geographically correlated host traits (Sanders et al., 2014; Sieber et al., 2018). Stronger evidence of co-evolution can be obtained by observing codivergence which represents the congruence of phylogenies between host and microbe or between divergent groups of microbes. It is typically observed only for a subset of dominant or key organisms within the microbial community. In a recent study of the microbial composition of skeleton, mucus and tissue in scleractinian corals, codivergence was observed in only four out of hundreds of microbial genera (Pollock et al., 2018).

Comparative genomic analyses may reveal changes in genome size, gene content or rates of gene evolution that are indicative of co-evolution. In obligate endosymbionts such *Polynucleobacter* in ciliates (Boscaro et al., 2017) and the aphid symbiont, *Buchnera aphidicola* (van Ham et al., 2003) the close association has resulted in substantial gene loss and genome size reduction. Analysis of microbial and host gene content can also reveal metabolic complementarity whereby both partners contribute to a shared function such as the synthesis of key amino acids (Russell et al., 2013) or complete digestion of a primary energy source (Poulsen et al., 2014).

The expansion of genes required for inter-species interactions, changes in the evolutionary rate of these genes, and horizontal gene transfer (HGT) may all also be observed where co-evolution is taking place (Friesen et al., 2006; Husnik et al., 2013). The close association and potential for metabolic interaction between species in a metaorganism suggests that HGT might be prevalent in such systems

(Keeling and Palmer, 2008; Degnan, 2014) but the confident identification of instances of HGT is difficult. This is because the elimination of alternate hypotheses (e.g. widespread gene loss) requires a dense and high quality sampling of the genomes of related taxa, a condition that can rarely be fulfilled with current genomic databases.

The major bottleneck to the use of comparative genomics in the study of co-evolution comes from limitations in current genome and metagenome sequencing techniques. Sequencing of large eukaryotic genomes remains expensive and difficult, meaning that publicly available genomes are often sparsely distributed across high level taxa and, where genomes are available, they often contain assembly errors and are highly fragmented (Salzberg et al., 2012). Gene annotations in such draft genomes may also be fragmented, missing or falsely duplicated (Zhang et al., 2012; Denton et al., 2014). It is recognised that interpretation of de novo assembled genomes is challenging and multiple high quality genomes are required for fruitful comparative studies (Richards, 2018). Overall, these problems greatly complicate inferences based on gene content and arrangement which could otherwise be used to infer inter-organismal partnerships or horizontal gene transfer.

Several technologies are emerging that promise to improve our understanding of the mechanisms behind co-evolution. Long and linked-read sequencing (Ott et al., 2018; Wallberg et al., 2018) as well as Hi-C contact maps allow for substantial improvements in assembly contiguity and can account for haplotypic variation (Chin et al., 2016; Kronenberg et al., 2018; Ott et al., 2018), a major source of fragmentation and errors in short read assemblies (Kajitani et al., 2014; Goltsman et al., 2017). Furthermore, long read RNA sequencing has the potential to significantly improve genome annotation as it is capable of generating near perfect full length transcripts without an assembly step. These can be used as very high quality training data for gene prediction, greatly reducing the number of incomplete or incorrect gene models in draft genomes (Magrini et al., 2018).

Obtaining complete genome sequences for bacterial, archaeal and viral partners presents a separate set of challenges. Although these genomes are relatively compact and easily assembled when sequenced from pure cultures, such cultures are rarely available. Instead, sequencing is often done on complex community samples resulting in a mix of reads from many different organisms. As mentioned above (see “Identification and quantification of microbial communities”), there are several new technologies that facilitate the binning of metagenomic reads that should eventually allow for the assembly of complete genomes from uncultured microbial samples (Bishara et al., 2018).

#### 4. Measuring molecular activity

The ability to measure molecular activity and track metabolic exchanges is key to understanding the interactions between members of a metaorganism. Such measurements have the potential to provide insights into what costs and benefits are incurred, how these are distributed between members, and what signals are used to allow organisms to avoid conflict with or exert control over others. There is growing recognition that answering these questions is an essential requirement for understanding animal and plant health (Berendsen et al., 2012; Cho and Blaser, 2012). For example, a complex picture of interactions between the human gut microbiome and the brain (gut brain axis) is emerging, which has been linked to psychiatric disorders as well as Multiple Sclerosis and the inflammatory bowel diseases, Crohn’s disease and Ulcerative colitis (Collins et al., 2012).

Many of the best understood interactions between metaorganism members have been studied via experiments that manipulate the microbiota of a host (e.g. through removal and selective re-introduction). Notable examples include metabolic pathways shared between the bacterium, *Buchnera* and pea aphid hosts (Russell et al., 2013) to produce essential amino acids, fungal resistance in *Hydra* conferred by the presence of bacteria-bacteria interactions (Fraune et al., 2015) and the induction of intestinal Th17 cells in mice upon reintroduction of a

single species of bacteria (Ivanov et al., 2009). While such experiments are often crucial to confirming that an interaction occurs, they typically rely on univariate assays to demonstrate the outcome (e.g. amino acid production, fungal immunity and cell growth) and additional techniques are required in order to explore the underlying molecular mechanisms at play.


One approach is to manipulate the molecular activity of the microbiota or host through experiments that knockdown or over-express specific genes. The molecular mechanisms by which *Hydra* are able to influence their microbiome have been revealed through experiments that knockdown the expression of specific antimicrobial peptides (AMPs) (Franzenburg et al., 2013) or that knockdown the expression of genes that regulate AMP expression such as FoxO (Boehm et al., 2012; Mortzfeld et al., 2018). This approach is extremely powerful but it is time consuming and is therefore only useful where prior information is available to generate well developed hypotheses related to specific genes and their effects.

An alternative approach that can complement these experiments is to use high throughput technologies such as RNA sequencing, and Mass Spectrometry to measure the expression of thousands of genes, transcripts, proteins or metabolites. In principle, it is even possible to measure changes in the molecular repertoires for multiple taxa within a holobiont simultaneously (cf DualRNA Seq; (Westermann et al., 2017)). Collectively termed ‘Omics approaches, these techniques can be used to generate plausible hypotheses for mechanisms of interaction between members of a metaorganism (Mohamed et al., 2016; Oakley et al., 2016). Although such ‘Omics approaches are now widely used in molecular biology, their use in a metaorganism context poses additional challenges. Increased sensitivity and dynamic range may be required to overcome a dominance of host signal or to detect metabolic activity from low abundance microbes. Data interpretation is also difficult because detailed information about molecular pathways and their associated genes and metabolites is largely derived from experiments on a small set of classical model taxa (eg *E. coli*, Yeast, Mouse, *Drosophila*, *C. elegans*). It is therefore important that ‘Omics approaches are adopted in conjunction with efforts to expand fundamental knowledge of the molecular biology of model metaorganisms and, more broadly, for non-model taxa.

#### 5. Mapping microbiome components within host tissues

Holobionts are not homogenous associations but vary in composition and activity between host tissues. This has perhaps been most extensively studied in humans where body location dramatically affects microbiome composition (Human Microbiome Project Consortium, 2012; Donaldson et al., 2016; Tropini et al., 2017; Byrd et al., 2018), but such distinctions have also been observed for very simple animals such as *Hydra* (Augustin et al., 2017).

The roles of individual microbiome components within the metaorganism often prove enigmatic. However, functions of microbe-microbe and microbe-host interactions may be revealed by locating the physical sites where these interactions occur (Fig. 1). One way to approach this is through methods that reveal the location of specific microorganisms within host tissues such as Fluorescent In Situ Hybridization (FISH). Originally named “phylogenetic staining” (DeLong et al., 1989), FISH revolutionized the field by allowing detection and identification of microbes without the necessity of cultivation (reviewed by (Amann et al., 1995; Wagner et al., 2003)). This feature of FISH makes it particularly valuable for studies of non-model systems, including corals and sponges, where culture methods have not yet been developed. Despite difficulties associated with high levels of autofluorescence of coral tissues (Wada et al., 2016), FISH has successfully revealed an intimate physical association between *Endozoicomonas sp.*, considered a candidate mutualist within tissues of a coral host (Neave et al., 2017b; Pogoreutz et al., 2018). By detection of specific microorganisms within oocytes and larvae of marine sponges, FISH has



**Fig. 1.** Methods for mapping microbiome components within host tissues include techniques such as nanoscale secondary ion mass spectrometry (nanoSIMS) that allow subcellular imaging of metabolites, and techniques such as fluorescence *in situ* hybridisation (FISH) that map the location of specific microorganisms. When applied in combination, these techniques can be used to make inferences about metabolic activity in microbe-microbe or microbe-host interactions.

confirmed vertical transmission of symbionts as had been suggested by sequencing-based methods (Schmitt et al., 2008; Webster et al., 2010).

While originally a relatively simple technique, FISH, often used in combination with other techniques, has become an increasingly sophisticated set of methodologies allowing characterization, quantification and co-localization of diverse microbes. For example, Combinatorial Labeling and Spectral Imaging FISH (CLASI-FISH), which can identify and differentiate up to 15 microbial taxa simultaneously (Valm et al., 2011), has been used to describe a highly spatially structured, multi-genus assembly of microbes in human plaque and provided insight into the function of individual components of this complex consortium (Mark Welch et al., 2016). On the other hand, combining FISH with electron microscopy imaging (fluorescence *in situ* hybridisation-correlative light and electron microscopy; FISH-CLEM) permitted identification and characterization of Poribacteria, common but uncultivated symbionts of marine sponges (Jahn et al., 2016). Combining FISH with laser microdissection, followed by amplicon sequencing of the isolated samples (Klitgaard et al., 2005), has the capacity to provide a high-resolution map of the microbiome within host tissues. One such study revealed that the dermis, the deepest layer of human skin and previously considered sterile, contains a diverse microbiome permitting direct communication with the host tissue (Nakatsuji et al., 2013).

While FISH, by its nature, can only be used on fixed specimens, live imaging of microbial-host interactions is possible in those experimental model systems where symbiotic microbes are amenable to culture and genome manipulation, and the host tissues are transparent. By introducing genes encoding green and red fluorescent proteins into *Aeromonas* and *Vibrio* strains isolated from the zebrafish intestinal tract, Wiles and colleagues generated bacteria which could be visualized in a minimally-invasive way in zebrafish larvae (Wiles et al., 2016).

In this study, comparison of bacterial population dynamics in wild type and reduced gut motility mutant zebrafish demonstrated the importance of host (rather than simply direct bacterial competition) in structuring of the vertebrate gut bacterial community.

Methods that detect and identify metabolites with subcellular

spatial resolution are emerging as powerful tools for studying organismal interactions. Metabolic imaging at subcellular resolution with labelled or label-free methods permits the identification of a broad range of molecules as well as their localization. Live imaging (via introduced green fluorescent protein fluorescence) combined with mass spectrometry showed that an interaction between *Ralstonia solanacearum* and soil fungi may be responsible for the virulence, persistence and proliferation of the bacterial pathogen. Subsequent gene disruption experiments combined with confocal microscopy confirmed that the arrangement of the bacterium is primarily determined by a metabolite produced by the bacteria (ralsolamycin). Using mass spectrometry, the ralsolamycin was visualized at the interface between *R. solanacearum* colonies and approaching fungal hyphae (Spraker et al., 2016).

This direction of study promises to improve our understanding of the function and structural principles of inter-organismal relationships and to provide insight into an array of fundamental biology issues (e.g., cell-cell recognition, immunity, signaling, cell-cycle control). To this end, secondary ion mass spectrometry (SIMS), while originally mainly employed by material scientists, has developed into a technique capable of imaging tissues, single cells, and microbes revealing chemical species with sub-micrometer spatial resolution (Henss et al., 2013; Gamble and Anderton, 2016). One of these methods, NanoSIMS, provides nanometer-scale resolution able to resolve the location of specific molecules/metabolites within bacterial cells. On the other hand, ToF-SIMS instruments have a ‘coarser’ resolution (typically in the range of micrometers), but can be regarded as “molecular microscopes” that generate chemical maps either across an area or via depth profiling to allow for the three-dimensional reconstructions of cell and tissue structure and molecular composition (Gamble and Anderton, 2016). The high spatial resolution and ability to detect and quantify a wide range of compounds means that these techniques can trace metabolic activity to microbial aggregations or even to specifically labelled microorganisms (Alonso et al., 2012). This is particularly promising for the study of marine cnidarian invertebrates and their associated microalgae and bacteria (Neave et al., 2016), where nanoSIMS has traced key metabolic interactions between algae and bacteria (Raina et al., 2017) as well as

between microalgae and cnidarian hosts (Rädecker et al., 2018).

In addition to these emerging high-tech methods, it is important to note that simply dissecting host tissues can also help to understand the distribution of microbial aggregations. Here choice of model species can be an important pre-requisite. The small size of most coral polyps usually precludes dissection but the extended large polyps of the mushroom coral *Heliofungia* offer an opportunity to look at the biology and physiology of the polyp on a wholly different scale than in other corals. Just as the giant axons of the squid *Loligo* - at that time, a relatively unknown organism - enabled major advances in neurophysiology (e.g.(Young, 1938)), being creative in selecting target species may permit insights into the spatial organisation of microbes in organisms such as corals. *Heliofungia* are solitary polyps which disassociate from the substrate after development and grow to a maximum size of 50 cm, a diameter several orders of magnitude larger than the average polyp from other corals. This allows for the dissection of tissue layers (a method originally developed for anemone (Richier et al., 2006)), sampling of different localities on the polyp (ie distal vs. central, base of tentacle vs. tip of tentacle, etc.), treatment or exposure trials, and sampling of the oral/gastric system. This coral is a promising model on which applying the molecular methods described in this essay in the ultimate goal to map the functional profile of microbes across a coral polyp. Understanding where microbes aggregate, how they function and how this function varies with microhabitat will illuminate the relationship between host and microbe at the molecular level, thus allowing for a deeper understanding of the relationship at the scale of organism or population.

## 6. Conclusions

Advances in molecular techniques over the past few decades have allowed the diversity of microbes in a wide variety of environments and biological systems to be estimated. It is now clear that many aspects of host organism biology depend on interactions with associated microbes. In this essay we reviewed four categories of molecular techniques for studying such interactions and identified several key technological developments, some of which may blur the lines between these categories in the future. One such advance is the ability to sequence whole communities of micro-organisms and reliably track sequences back to their cell of origin. This will vastly improve our ability to assemble whole microbial genomes and may accelerate the shift from amplicon based to whole genome based methods for microbial community profiling. Widespread adoption of whole genome based methods could have far-reaching effects on our ability to understand metabolic interactions by facilitating an accurate assessment of gene content within microbial communities. It would also improve our ability to understand the evolution of holobionts by providing dense taxonomic sampling for comparative genomic analyses. Another promising advance is the development of molecular imaging techniques that allow complex information to be gathered in a spatially resolved fashion. These include techniques that measure metabolites at sub-cellular resolution (nanoSIMS) which, when combined with techniques that image the the spatial distribution of microbial communities within a host, could greatly improve our understanding of metabolic interactions between partner organisms.

## Acknowledgements

This review emerged from discussions on the role of molecular techniques in holobiont research that were held as part of an Australian Academy of Sciences Boden research conference on Magnetic Island in July 2018. We wish to thank organisations that funded this conference including the Australian Academy of Sciences, Great Barrier Reef Foundation, Ian Potter Foundation, the ARC CoE for coral reef studies, and the Collaborative Research Centre (CRC 1182, funded through the German Research Foundation DFG) "Origin and Function of

Metaorganisms".

## Consortium of Australian Academy of Science Boden Research Conference Participants (in alphabetical order):

Tracy Ainsworth (James Cook University, Townsville, Australia); Natalia R. Andrade (ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville QLD, Australia); A. Elizabeth Arnold (School of Plant Sciences and the Department of Ecology and Evolutionary Biology, The University of Arizona, USA); Eldon Ball (The Australian National University, Canberra, Australia); David Bourne (James Cook University, Townsville, Australia); Thomas CG Bosch (Zoological Institute, Kiel University, Kiel, Germany); Nicholas J. Butterfield (University of Cambridge, Cambridge, United Kingdom); Cheong Xin Chan (The University of Queensland, Brisbane, Australia); Peter F. Cowman (James Cook University, Townsville, Australia); Simon K. Davy (Victoria University of Wellington, Wellington, New Zealand); Katharina Fabricius (Australian Institute of Marine Science, Townsville, Australia); Sofia V. Fortunato (James Cook University, Townsville, Australia); Sebastian Fraune (Zoological Institute, Kiel University, Kiel, Germany); Alejandra Hernandez (James Cook University, Townsville, Australia); Mia Hoogenboom (James Cook University, Townsville, Australia); Cornelia Jaspers (GEOMAR – Helmholtz Centre for Ocean Research Kiel, Evolutionary Ecology of Marine Fishes, Kiel, Germany); Amin Mohamed (CSIRO, St. Lucia, Australia); Lucia Pita (GEOMAR Helmholtz Centre for Ocean Research, Kiel, Germany); Mark A. Ragan (The University of Queensland, Brisbane, Australia); Kazuhiro Sakamaki (Kyoto University, Kyoto, Japan); Verena Schoepf (The University of Western Australia, Perth, Australia); Torsten Seemann (The University of Melbourne, Melbourne, Australia); Chuya Shinzato (The University of Tokyo, Chiba, Japan); Jarosław Stolarski (Polish Academy of Sciences, Warsaw, Poland); Shunichi Takahashi (National Institute for Basic Biology, Okazaki, Japan); Sen-Lin Tang (National Taiwan University, Taipei, Taiwan); Nicole Webster (Australian Institute of Marine Science, Townsville, Australia); Brooke Whitelaw (James Cook University, Townsville, Australia).

## References

- Abu-Ali, G.S., Mehta, R.S., Lloyd-Price, J., Mallick, H., Branck, T., Ivey, K.L., Drew, D.A., DuLong, C., Rimm, E., Izard, J., Chan, A.T., Huttenhower, C., 2018. Metatranscriptome of human faecal microbial communities in a cohort of adult men. *Nat Microbiol* 3, 356–366.
- Alonso, C., Musat, N., Adam, B., Kuypers, M., Amann, R., 2012. HISH-SIMS analysis of bacterial uptake of algal-derived carbon in the Río de la Plata estuary. *Syst. Appl. Microbiol.* 35, 541–548.
- Amann, R.L., Ludwig, W., Schleifer, K.H., 1995. Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microbiol. Rev.* 59, 143–169.
- Arumugam, K., Bessarab, I., Liu, X., Natarajan, G., Drautz-Moses, D.I., Wuertz, S., Lauro, F.M., Law, Y.Y., Huson, D.H., Williams, R.B.H., 2018. Improving recovery of member genomes from enrichment reactor microbial communities using MinION based long-read metagenomics. *bioRxiv*. <https://doi.org/10.1101/465328>.
- Augustin, R., Schröder, K., Murillo Rincón, A.P., Fraune, S., Anton-Erxleben, F., Herbst, E.-M., Wittlieb, J., Schwentner, M., Grötzinger, J., Wassenaar, T.M., Bosch, T.C.G., 2017. A secreted antibacterial neuropeptide shapes the microbiome of *Hydra*. *Nat. Commun.* 8, 698.
- Bang, C., Dagan, T., Deines, P., Dubilier, N., Duschl, W.J., Fraune, S., Hentschel, U., Hirt, H., Hübler, N., Lachnit, T., Picazo, D., Pita, L., Pogoreutz, C., Rädecker, N., Saad, M.M., Schmitz, R.A., Schulten, H., Voolstra, C.R., Weiland-Bräuer, N., Ziegler, M., Bosch, T.C.G., 2018. Metaorganisms in extreme environments: do microbes play a role in organismal adaptation? *Zoology* 127, 1–19.
- Baumann, P., Moran, N.A., Baumann, L., 1997. The evolution and genetics of aphid endosymbionts. *BioScience* 47, 12–20.
- Berendsen, R.L., Pieterse, C.M.J., Bakker, P.A.H.M., 2012. The rhizosphere microbiome and plant health. *Trends Plant Sci.* 17, 478–486.
- Bertrand, D., Shaw, J., Narayan, M., Ng, H.Q.A., Kumar, S., Li, C., Dvornicic, M., Soldo, J.P., Kho, J.Y., Ng, O.T., Barkham, T., Young, B., Marimuthu, K., Chng, K.R., Sikic, M., Nagarajan, N., 2018. Nanopore sequencing enables high-resolution analysis of resistance determinants and mobile elements in the human gut microbiome. *bioRxiv*. <https://doi.org/10.1101/456905>.
- Bishara, A., Moss, E.L., Kolmogorov, M., Parada, A.E., Weng, Z., Sidow, A., Dekas, A.E., Batzoglou, S., Bhatt, A.S., 2018. High-quality genome sequences of uncultured microbes by assembly of read clouds. *Nat. Biotechnol.* <https://doi.org/10.1038/nbt.4266>.

- Boehm, A.-M., Khalturin, K., Anton-Erxleben, F., Hemmrich, G., Klostermeier, U.C., Lopez-Quintero, J.A., Oberg, H.-H., Puchert, M., Rosenstiel, P., Wittlieb, J., Bosch, T.C.G., 2012. FoxO is a critical regulator of stem cell maintenance in immortal *Hydra*. *Proc. Natl. Acad. Sci. U. S. A.* 109, 19697–19702.
- Boscaro, V., Kolisko, M., Felletti, M., Vannini, C., Lynn, D.H., Keeling, P.J., 2017. Parallel genome reduction in symbionts descended from closely related free-living bacteria. *Nat. Ecol. Evol.* 1, 1160–1167.
- Bosch, T.C.G., McFall-Ngai, M.J., 2011. Metaorganisms as the new frontier. *Zoology* 114, 185–190.
- Bosch, T., Miller, D.J., 2016. *The Holobiont Imperative*. Springer, Vienna.
- Brooks, A.W., Kohl, K.D., Brucker, R.M., van Opstal, E.J., Bordenstein, S.R., 2016. Phyllosymbiosis: relationships and functional effects of microbial communities across host evolutionary history. *PLoS Biol.* 14, e2000225.
- Brooks, J.P., Edwards, D.J., Harwich Jr., M.D., Rivera, M.C., Fettweis, J.M., Serrano, M.G., Reis, R.A., Sheth, N.U., Huang, B., Girerd, P., Vaginal Microbiome Consortium, Strauss 3rd, J.F., Jefferson, K.K., Buck, G.A., 2015. The truth about metagenomics: quantifying and counteracting bias in 16S rRNA studies. *BMC Microbiol.* 15, 66.
- Brucker, R.M., Bordenstein, S.R., 2012. Speciation by symbiosis. *Trends Ecol. Evol.* 27, 443–451.
- Byrd, A.L., Belkaid, Y., Segre, J.A., 2018. The human skin microbiome. *Nat. Rev. Microbiol.* 16, 143–155.
- Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* 13, 581–583.
- Callahan, B.J., Wong, J., Heiner, C., Oh, S., Theriot, C.M., Gulati, A.S., McGill, S.K., Dougherty, M.K., 2018. High-throughput amplicon sequencing of the full-length 16S rRNA gene with single-nucleotide resolution. *bioRxiv*. <https://doi.org/10.1101/392332>.
- Chandler, J.A., Lang, J.M., Bhatnagar, S., Eisen, J.A., Kopp, A., 2011. Bacterial communities of diverse *Drosophila* species: ecological context of a host-microbe model system. *PLoS Genet.* 7, e1002272.
- Chin, C.-S., Peluso, P., Sedlazeck, F.J., Nattestad, M., Concepcion, G.T., Clum, A., Dunn, C., O'Malley, R., Figueroa-Balderas, R., Morales-Cruz, A., Cramer, G.R., Delledonne, M., Luo, C., Ecker, J.R., Cantu, D., Rank, D.R., Schatz, M.C., 2016. Phased diploid genome assembly with single-molecule real-time sequencing. *Nat. Methods* 13, 1050–1054.
- Cho, I., Blaser, M.J., 2012. The human microbiome: at the interface of health and disease. *Nat. Rev. Genet.* 13, 260–270.
- Collins, S.M., Surette, M., Bercik, P., 2012. The interplay between the intestinal microbiota and the brain. *Nat. Rev. Microbiol.* 10, 735–742.
- Daims, H., Ramsing, N.B., Schleifer, K.H., Wagner, M., 2001. Cultivation-independent, semiautomatic determination of absolute bacterial cell numbers in environmental samples by fluorescence in situ hybridization. *Appl. Environ. Microbiol.* 67, 5810–5818.
- Degnan, S.M., 2014. Think laterally: horizontal gene transfer from symbiotic microbes may extend the phenotype of marine sessile hosts. *Front. Microbiol.* 5, 638.
- DeLong, E.F., Wickham, G.S., Pace, N.R., 1989. Phylogenetic stains: ribosomal RNA-based probes for the identification of single cells. *Science* 243, 1360–1363.
- Denton, J.F., Lugo-Martinez, J., Tucker, A.E., Schrider, D.R., Warren, W.C., Hahn, M.W., 2014. Extensive error in the number of genes inferred from draft genome assemblies. *PLoS Comput. Biol.* 10, e1003998.
- Donaldson, G.P., Lee, S.M., Mazmanian, S.K., 2016. Gut biogeography of the bacterial microbiota. *Nat. Rev. Microbiol.* 14, 20–32.
- Dunn, S.R., Phillips, W.S., Green, D.R., Weis, V.M., 2007. Knockdown of actin and caspase gene expression by RNA interference in the symbiotic anemone *Aiptasia pallida*. *Biol. Bull.* 212, 250–258.
- Franzenburg, S., Walter, J., Künzel, S., Wang, J., Baines, J.F., Bosch, T.C.G., Fraune, S., 2013. Distinct antimicrobial peptide expression determines host species-specific bacterial associations. *Proc. Natl. Acad. Sci. U. S. A.* 110, E3730–E3738.
- Fraune, S., Anton-Erxleben, F., Augustin, R., Franzenburg, S., Knop, M., Schröder, K., Willoweit-Ohl, D., Bosch, T.C.G., 2015. Bacteria-bacteria interactions within the microbiota of the ancestral metazoan *Hydra* contribute to fungal resistance. *ISME J.* 9, 1543–1556.
- Fraune, S., Bosch, T.C.G., 2007. Long-term maintenance of species-specific bacterial microbiota in the basal metazoan *Hydra*. *Proc. Natl. Acad. Sci. U. S. A.* 104, 13146–13151.
- Friesen, T.L., Stukenbrock, E.H., Liu, Z., Meinhardt, S., Ling, H., Faris, J.D., Rasmussen, J.B., Solomon, P.S., McDonald, B.A., Oliver, R.P., 2006. Emergence of a new disease as a result of interspecific virulence gene transfer. *Nat. Genet.* 38, 953–956.
- Gamble, L.J., Anderton, C.R., 2016. Secondary Ion Mass Spectrometry Imaging of tissues, cells, and microbial systems. *Microsc. Today* 24, 24–31.
- Gloor, G.B., Macklaim, J.M., Pawlowsky-Glahn, V., Egozcue, J.J., 2017. Microbiome datasets are compositional: and this is not optional. *Front. Microbiol.* 8, 2224.
- Goltsman, E., Ho, I., Rokhsar, D., 2017. Meraculous-2D: Haplotype-sensitive assembly of highly heterozygous genomes. *arXiv [q-bio.GN]*.
- Henss, A., Rohnke, M., El Khassawna, T., Govindarajan, P., Schlewitz, G., Heiss, C., Janek, J., 2013. Applicability of ToF-SIMS for monitoring compositional changes in bone in a long-term animal model. *J. R. Soc. Interface* 10, 20130332.
- Hernandez-Agreda, A., Gates, R.D., Ainsworth, T.D., 2017. Defining the core microbiome in corals' microbial soup. *Trends Microbiol.*
- Hillyer, K.E., Dias, D.A., Lutz, A., Roessner, U., Davy, S.K., 2017. Mapping carbon fate during bleaching in a model cnidarian symbiosis: the application of 13 C metabolomics. *New Phytol.* 214, 1551–1562.
- Human Microbiome Project Consortium, 2012. Structure, function and diversity of the healthy human microbiome. *Nature* 486, 207–214.
- Husnik, F., Nikoh, N., Koga, R., Ross, L., Duncan, R.P., Fujie, M., Tanaka, M., Satoh, N., Bachtrog, D., Wilson, A.C.C., von Dohlen, C.D., Fukatsu, T., McCutcheon, J.P., 2013. Horizontal gene transfer from diverse bacteria to an insect genome enables a tripartite nested mealybug symbiosis. *Cell* 153, 1567–1578.
- Ikmi, A., McKinney, S.A., Delventhal, K.M., Gibson, M.C., 2014. TALEN and CRISPR/Cas9-mediated genome editing in the early-branching metazoan *Nematostella vectensis*. *Nat. Commun.* 5, 5486.
- Ivanov, I.I., Atarashi, K., Manel, N., Brodie, E.L., Shima, T., Karaoz, U., Wei, D., Goldfarb, K.C., Santee, C.A., Lynch, S.V., Tanoue, T., Imaoka, A., Itoh, K., Takeda, K., Umesaki, Y., Honda, K., Littman, D.R., 2009. Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell* 139, 485–498.
- Jahn, M.T., Markert, S.M., Ryu, T., Ravasi, T., Stigloher, C., Hentschel, U., Moitinho-Silva, L., 2016. Shedding light on cell compartmentation in the candidate phylum *Poribacteria* by high resolution visualisation and transcriptional profiling. *Sci. Rep.* 6, 35860.
- Jain, C., Rodriguez-R, L.M., Phillippy, A.M., Konstantinidis, K.T., Aluru, S., 2018. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat. Commun.* 9, 5114.
- Jaspers, C., Fraune, S., Consortium of Australian Academy of Science Boden Research Conference Participants, Arnold, A., Elizabeth, Miller, D.J., Bosch, T., Voolstra, C.R., 2019. Resolving structure and function of metaorganisms through a holistic framework combining reductionist and integrative approaches. *Zoology*. <https://doi.org/10.1016/j.zool.2019.02.007>.
- Jian, C., Luukkonen, P., Yki-Jarvinen, H., Salonen, A., Korpela, K., 2018. Quantitative PCR provides a simple and accessible method for quantitative microbiome profiling. *bioRxiv*. <https://doi.org/10.1101/478685>.
- Kajitani, R., Tshimoto, K., Noguchi, H., Toyoda, A., Ogura, Y., Okuno, M., Yabana, M., Harada, M., Nagayasu, E., Maruyama, H., Kohara, Y., Fujiyama, A., Hayashi, T., Itoh, T., 2014. Efficient de novo assembly of highly heterozygous genomes from whole-genome shotgun short reads. *Genome Res.* 24, 1384–1395.
- Keeling, P.J., Palmer, J.D., 2008. Horizontal gene transfer in eukaryotic evolution. *Nat. Rev. Genet.* 9, 605–618.
- Kelley, S.T., Dobler, S., 2011. Comparative analysis of microbial diversity in *Longitarsus* flea beetles (Coleoptera: Chrysomelidae). *Genetica* 139, 541–550.
- Kennedy, K., Hall, M.W., Lynch, M.D.J., Moreno-Hagelsieb, G., Neufeld, J.D., 2014. Evaluating bias of illumina-based bacterial 16S rRNA gene profiles. *Appl. Environ. Microbiol.* 80, 5717–5722.
- Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., Glöckner, F.O., 2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res.* 41, e1.
- Klitgaard, K., Mølbak, L., Jensen, T.K., Lindboe, C.F., Boye, M., 2005. Laser capture microdissection of bacterial cells targeted by fluorescence in situ hybridization. *Biotechniques* 39, 864–868.
- Konstantinidis Konstantinos, T., Alban, Ramette, Tiedje James, M., 2006. The bacterial species definition in the genomic era. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 361, 1929–1940.
- Kronenberg, Z.N., Hall, R.J., Hiendleder, S., Smith, T.P.L., Sullivan, S.T., Williams, J.L., Kingan, S.B., 2018. FALCON-Phase: Integrating PacBio and Hi-C data for phased diploid genomes. *bioRxiv*. <https://doi.org/10.1101/327064>.
- Laursen, M.F., Dalgaard, M.D., Bahl, M.I., 2017. Genomic GC-content affects the accuracy of 16S rRNA gene sequencing based microbial profiling due to PCR bias. *Front. Microbiol.* 8, 1934.
- Lindgreen, S., Adair, K.L., Gardner, P.P., 2016. An evaluation of the accuracy and speed of metagenome analysis tools. *Sci. Rep.* 6, 19233.
- Liu, M., Darling, A., 2015. Metagenomic chromosome conformation capture (3C): techniques, applications, and challenges. *F1000Res.* 4, 1377.
- Lovén, J., Orland, D.A., Sigova, A.A., Lin, C.Y., Rahl, P.B., Burge, C.B., Levens, D.L., Lee, T.I., Young, R.A., 2012. Revisiting global gene expression analysis. *Cell* 151, 476–482.
- Magrini, V., Gao, X., Rosa, B.A., McGrath, S., Zhang, X., Hallsworth-Pepin, K., Martin, J., Hawdon, J., Wilson, R.K., Mitreva, M., 2018. Improving eukaryotic genome annotation using single molecule mRNA sequencing. *BMC Genomics* 19, 172.
- Mark Welch, J.L., Rossetti, B.J., Rieken, C.W., Dewhurst, F.E., Borisy, G.G., 2016. Biogeography of a human oral microbiome at the micron scale. *Proc. Natl. Acad. Sci. U. S. A.* 113, E791–E800.
- Moeller, A.H., Caro-Quintero, A., Mjunga, D., Georgiev, A.V., Lonsdorf, E.V., Muller, M.N., Pusey, A.E., Peeters, M., Hahn, B.H., Ochman, H., 2016. Cospeciation of gut microbiota with hominids. *Science* 353, 380–382.
- Mohamed, A.R., Cumbo, V., Harii, S., Shinzato, C., Chan, C.X., Ragan, M.A., Bourne, D.G., Willis, B.L., Ball, E.E., Satoh, N., Miller, D.J., 2016. The transcriptomic response of the coral *Acropora digitifera* to a competent *Symbiodinium* strain: the symbiosome as an arrested early phagosome. *Mol. Ecol.* 25, 3127–3141.
- Moran, N.A., Yun, Y., 2015. Experimental replacement of an obligate insect symbiont. *Proc. Natl. Acad. Sci. U. S. A.* 112, 2093–2096.
- Mortzfeld, B.M., Taubenheim, J., Fraune, S., Klimovich, A.V., Bosch, T.C.G., 2018. Stem cell transcription factor FoxO controls microbiome resilience in *Hydra*. *Front. Microbiol.* 9, 629.
- Nakatsuji, T., Chiang, H.-I., Jiang, S.B., Nagarajan, H., Zengler, K., Gallo, R.L., 2013. The microbiome extends to subepidermal compartments of normal skin. *Nat. Commun.* 4, 1431.
- Neave, M.J., Apprill, A., Ferrier-Pagès, C., Voolstra, C.R., 2016. Diversity and function of prevalent symbiotic marine bacteria in the genus *Endozoicomonas*. *Appl. Microbiol. Biotechnol.* 100, 8315–8324.
- Neave, M.J., Michell, C.T., Apprill, A., Voolstra, C.R., 2017a. *Endozoicomonas* genomes reveal functional adaptation and plasticity in bacterial strains symbiotically associated with diverse marine hosts. *Sci. Rep.* 7, 40579.
- Neave, M.J., Rachmawati, R., Xun, L., Michell, C.T., Bourne, D.G., Apprill, A., Voolstra,

- C.R., 2017b. Differential specificity between closely related corals and abundant *Endozoicomonas* endosymbionts across global scales. *ISME J.* 11, 186–200.
- Nicholls, S.M., Quick, J.C., Tang, S., Loman, N.J., 2018. Ultra-deep, long-read nanopore sequencing of mock microbial community standards. *bioRxiv*. <https://doi.org/10.1101/487033>.
- Oakley, C.A., Ameismeier, M.F., Peng, L., Weis, V.M., Grossman, A.R., Davy, S.K., 2016. Symbiosis induces widespread changes in the proteome of the model cnidarian *Aiptasia*. *Cell. Microbiol.* 18, 1009–1023.
- O'Brien, P.A., Webster, N.S., Miller, D.J., Bourne, D.G., 2019. Host-microbe coevolution: applying evidence from model systems to complex marine invertebrate holobionts. *mBio*. <https://doi.org/10.1128/mBio.02241-18>.
- Ott, A., Schnable, J.C., Yeh, C.-T., Wu, L., Liu, C., Hu, H.-C., Dalgard, C.L., Sarkar, S., Schnable, P.S., 2018. Linked read technology for assembling large complex and polyploid genomes. *BMC Genomics* 19, 651.
- Pogoreutz, C., Rådecker, N., Cárdenas, A., Gärdes, A., Wild, C., Voolstra, C.R., 2018. Dominance of *Endozoicomonas* bacteria throughout coral bleaching and mortality suggests structural inflexibility of the *Pocillopora verrucosa* microbiome. *Ecol. Evol.* 8, 2240–2252.
- Pollock, F.J., McMinds, R., Smith, S., Bourne, D.G., Willis, B.L., Medina, M., Thurber, R.V., Zaneveld, J.R., 2018. Coral-associated bacteria demonstrate phyllosymbiosis and cophylogeny. *Nat. Commun.* 9, 4921.
- Poulsen, M., Hu, H., Li, C., Chen, Z., Xu, L., Otani, S., Nygaard, S., Nobre, T., Klaubauf, S., Schindler, P.M., Hauser, F., Pan, H., Yang, Z., Sonnenberg, A.S.M., de Beer, Z.W., Zhang, Y., Wingfield, M.J., Grimmelikhuijzen, C.J.P., de Vries, R.P., Korb, J., Aanen, D.K., Wang, J., Boomsma, J.J., Zhang, G., 2014. Complementary symbiont contributions to plant decomposition in a fungus-farming termite. *Proc. Natl. Acad. Sci. U. S. A.* 111, 14500–14505.
- Props, R., Kerckhof, F.-M., Rubbens, P., De Vrieze, J., Hernandez Sanabria, E., Waegeman, W., Monsieus, P., Hammes, F., Boon, N., 2017. Absolute quantification of microbial taxon abundances. *ISME J.* 11, 584–587.
- Qin, J., Li, R., Raes, J., Arumugam, M., Burgdorf, K.S., Manichanh, C., Nielsen, T., Pons, N., Levenez, F., Yamada, T., Mende, D.R., Li, J., Xu, J., Li, S., Li, D., Cao, J., Wang, B., Liang, H., Zheng, H., Xie, Y., Tap, J., Lepage, P., Bertalan, M., Batto, J.-M., Hansen, T., Le Paslier, D., Linneberg, A., Nielsen, H.B., Pelletier, E., Renault, P., Sicheritz-Ponten, T., Turner, K., Zhu, H., Yu, C., Li, S., Jian, M., Zhou, Y., Li, Y., Zhang, X., Li, S., Qin, N., Yang, H., Wang, J., Brunak, S., Doré, J., Guarner, F., Kristiansen, K., Pedersen, O., Parkhill, J., Weissenbach, J., MetaHIT Consortium, Bork, P., Ehrlich, S.D., Wang, J., 2010. A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* 464, 59–65.
- Rådecker, N., Raina, J.-B., Pernice, M., Perna, G., Guagliardo, P., Kilburn, M.R., Aranda, M., Voolstra, C.R., 2018. Using *Aiptasia* as a model to study metabolic interactions in cnidarian-symbiodinium symbioses. *Front. Physiol.* 9, 214.
- Raina, J.-B., Clode, P.L., Cheong, S., Bougoure, J., Kilburn, M.R., Reeder, A., Forêt, S., Stat, M., Beltran, V., Thomas-Hall, P., Tapiolas, D., Motti, C.M., Gong, B., Pernice, M., Marjo, C.E., Seymour, J.R., Willis, B.L., Bourne, D.G., 2017. Subcellular tracking reveals the location of dimethylsulfoniopropionate in microalgae and visualises its uptake by marine bacteria. *Elife* 6. <https://doi.org/10.7554/eLife.23008>.
- Rawls, J.F., Mahowald, M.A., Ley, R.E., Gordon, J.L., 2006. Reciprocal gut microbiota transplants from zebrafish and mice to germ-free recipients reveal host habitat selection. *Cell* 127, 423–433.
- Richards, S., 2018. Full disclosure: Genome assembly is still hard. *PLoS Biol.* 16, e2005894.
- Richier, S., Sabourault, C., Courtiade, J., Zucchini, N., Allemand, D., Furla, P., 2006. Oxidative stress and apoptotic events during thermal stress in the symbiotic sea anemone, *Anemonia viridis*. *FEBS J.* 273, 4186–4198.
- Rintala, A., Pietilä, S., Munukka, E., Eerola, E., Pursiheimo, J.-P., Laiho, A., Pekkala, S., Huovinen, P., 2017. Gut microbiota analysis results are highly dependent on the 16S rRNA gene target region, whereas the impact of DNA extraction is minor. *J. Biomol. Tech.* 28, 19–30.
- Russell, C.W., Bouvaine, S., Newell, P.D., Douglas, A.E., 2013. Shared metabolic pathways in a coevolved insect-bacterial symbiosis. *Appl. Environ. Microbiol.* 79, 6117–6123.
- Salzberg, S.L., Phillippy, A.M., Zimin, A., Puiu, D., Magoc, T., Koren, S., Treangen, T.J., Schatz, M.C., Delcher, A.L., Roberts, M., Marçais, G., Pop, M., Yorke, J.A., 2012. GAGE: A critical evaluation of genome assemblies and assembly algorithms. *Genome Res.* 22, 557–567.
- Sanders, J.G., Powell, S., Kronauer, D.J.C., Vasconcelos, H.L., Frederickson, M.E., Pierce, N.E., 2014. Stability and phylogenetic correlation in gut microbiota: lessons from ants and apes. *Mol. Ecol.* 23, 1268–1283.
- Schmitt, S., Angermeier, H., Schiller, R., Lindquist, N., Hentschel, U., 2008. Molecular microbial diversity survey of sponge reproductive stages and mechanistic insights into vertical transmission of microbial symbionts. *Appl. Environ. Microbiol.* 74, 7694–7708.
- Shakya, M., Quince, C., Campbell, J.H., Yang, Z.K., Schadt, C.W., Podar, M., 2013. Comparative metagenomic and rRNA microbial diversity characterization using archaeal and bacterial synthetic communities. *Environ. Microbiol.* 15, 1882–1899.
- Sieber, M., Pita, L., Weiland-Bräuer, N., Dirksen, P., Wang, J., Mortzfeld, B., Franzenburg, S., Schmitz, R.A., Baines, J.F., Fraune, S., Hentschel, U., Schulenburg, H., Bosch, T.C.G., Traulsen, A., 2018. The neutral metaorganism. *bioRxiv*. <https://doi.org/10.1101/367243>.
- Spraker, J.E., Sanchez, L.M., Lowe, T.M., Dorrestein, P.C., Keller, N.P., 2016. *Ralstonia solanacearum* lipopeptide induces chlamyospore development in fungi and facilitates bacterial entry into fungal tissues. *ISME J.* 10, 2317–2330.
- Starr, A.E., Deeke, S.A., Li, L., Zhang, X., Daoud, R., Ryan, J., Ning, Z., Cheng, K., Nguyen, L.V.H., Abou-Samra, E., Lavallée-Adam, M., Figeys, D., 2018. Proteomic and Metaproteomic approaches to understand host-microbe interactions. *Anal. Chem.* 90, 86–109.
- Tropini, C., Earle, K.A., Huang, K.C., Sonnenburg, J.L., 2017. The gut microbiome: connecting spatial organization to function. *Cell Host Microbe* 21, 433–442.
- Valm, A.M., Mark Welch, J.L., Rieken, C.W., Hasegawa, Y., Sogin, M.L., Oldenbourg, R., Dewhirst, F.E., Borisy, G.G., 2011. Systems-level analysis of microbial community organization through combinatorial labeling and spectral imaging. *Proc. Natl. Acad. Sci. U. S. A.* 108, 4152–4157.
- van Ham, R.C.H.J., Kamerbeek, J., Palacios, C., Rausell, C., Abascal, F., Bastolla, U., Fernández, J.M., Jiménez, L., Postigo, M., Silva, F.J., Tamames, J., Viguera, E., Latorre, A., Valencia, A., Morán, F., Moya, A., 2003. Reductive genome evolution in *Buchnera aphidicola*. *Proc. Natl. Acad. Sci. U. S. A.* 100, 581–586.
- Voolstra, C.R., 2013. A journey into the wild of the cnidarian model system *Aiptasia* and its symbionts. *Mol. Ecol.* 22, 4366–4368.
- Wada, N., Pollock, F.J., Willis, B.L., Ainsworth, T., Mano, N., Bourne, D.G., 2016. In situ visualization of bacterial populations in coral tissues: pitfalls and solutions. *PeerJ* 4, e2424.
- Wagner, M., Horn, M., Daims, H., 2003. Fluorescence in situ hybridisation for the identification and characterisation of prokaryotes. *Curr. Opin. Microbiol.* 6, 302–309.
- Wallberg, A., Bunikis, I., Pettersson, O.V., Mosbech, M.-B., Childers, A.K., Evans, J.D., Mikheyev, A.S., Robertson, H.M., Robinson, G.E., Webster, M.T., 2018. A hybrid de novo genome assembly of the honeybee, *Apis mellifera*, with chromosome-length scaffolds. *bioRxiv*. <https://doi.org/10.1101/361469>.
- Webster, N.S., Taylor, M.W., Behnam, F., Lückner, S., Rattei, T., Whalan, S., Horn, M., Wagner, M., 2010. Deep sequencing reveals exceptional diversity and modes of transmission for bacterial sponge symbionts. *Environ. Microbiol.* 12, 2070–2082.
- Westermann, A.J., Barquist, L., Vogel, J., 2017. Resolving host-pathogen interactions by dual RNA-seq. *PLoS Pathog.* 13, e1006033.
- Wiles, T.J., Jemielita, M., Baker, R.P., Schlomann, B.H., Logan, S.L., Ganz, J., Melancon, E., Eisen, J.S., Guillemin, K., Parthasarathy, R., 2016. Host gut motility promotes competitive exclusion within a model intestinal microbiota. *PLoS Biol.* 14, e1002517.
- Woese, C.R., Fox, G.E., 1977. Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proc. Natl. Acad. Sci. U. S. A.* 74, 5088–5090.
- Xu, Y., Zhao, F., 2018. Single-cell metagenomics: challenges and applications. *Protein Cell* 9, 501–510.
- Xue, M., Wu, L., He, Y., Liang, H., Wen, C., 2018. Biases during DNA extraction affect characterization of the microbiota associated with larvae of the Pacific white shrimp, *Litopenaeus vannamei*. *PeerJ* 6, e5257.
- Yarza, P., Yilmaz, P., Pruesse, E., Glöckner, F.O., Ludwig, W., Schleifer, K.-H., Whitman, W.B., Euzéby, J., Amann, R., Rosselló-Móra, R., 2014. Uniting the classification of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nat. Rev. Microbiol.* 12, 635.
- Yilmaz, P., Parfrey, L.W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J., Ludwig, W., Glöckner, F.O., 2014. The SILVA and “All-species Living Tree Project (LTP)” taxonomic frameworks. *Nucleic Acids Res.* 42, D643–D648.
- Young, J.Z., 1938. The functioning of the giant nerve fibres of the squid. *J. Exp. Biol.* 15, 170–185.
- Zhang, X., Goodsell, J., Norgren Jr., R.B., 2012. Limitations of the rhesus macaque draft genome assembly and annotation. *BMC Genomics* 13, 206.
- Ziegler, M., Grupstra, C.G.B., Barreto, M.M., Eaton, M., BaOmar, J., Zubier, K., Al-Sofyani, A., Turki, A.J., Ormond, R., Voolstra, C.R., 2019. Coral bacterial community structure responds to environmental change in a host-specific manner. *Nat. Commun.* 10, 3092.