


Universität Konstanz



New Convexity Conditions in the Calculus of Variations and Compensated Compactness Theory

Krzysztof Chelminski
Agnieszka Kalamajska

Konstanzer Schriften in Mathematik und Informatik

Nr. 192, Juni 2003

ISSN 1430–3558

New Convexity Conditions in the Calculus of Variations and Compensated Compactness Theory

Krzysztof Chelmiński and Agnieszka Kałamajska *

University of Constance, Universitätsstr. 10, 78464 Konstanz, Germany

permanent address: Cardinal Stefan Wyszyński University, ul. Dewajtis 5,
01-815 Warszawa, Poland

email: chelminski@uksw.edu.pl

and

Institute of Mathematics of the Polish Academy of Sciences, ul. Śniadeckich 8,
00-950 Warszawa, Poland

permanent address: Institute of Mathematics, Warsaw University, ul. Banacha 2,
02-097 Warszawa, Poland

email: kalamajs@mimuw.edu.pl

Abstract

We consider the lower semicontinuous functional of the form $I_f(u) = \int_{\Omega} f(u) dx$ where u satisfies a given conservation law defined by differential operator of degree one with constant coefficients. We show that under certain constraints the well known Murat and Tartar's Λ -convexity condition for the integrand f extends to the new geometric conditions satisfied on four dimensional simplexes. Similar conditions on three dimensional simplexes were recently obtained by the second author. New conditions apply to quasiconvex functions.

AMS classification 2000: 49J10, 49J45

Key words: quasiconvexity, rank-one convexity, semicontinuity.

1 Introduction

Let $\Omega \subseteq \mathbf{R}^n$ be a bounded domain, $f : \mathbf{R}^{n \times m} \rightarrow \mathbf{R}$ be a continuous function and consider the variational functional

$$I_f(u) = \int_{\Omega} f(\nabla u) dx, \quad (1.1)$$

*The work of the second author is supported by a KBN grant no. 2-PO3A-028-22.

where u belongs to the Sobolev space $W^{1,\infty}(\Omega, \mathbf{R}^m)$. It was proved by Morrey in 1952, [51] that the functional $I_f(u)$ is sequentially lower semicontinuous with respect to weak-* convergence in $W^{1,\infty}$ (i.e. $\liminf_{\nu \rightarrow \infty} I_f(u^\nu) \geq I_f(u)$ provided that $u^\nu \rightarrow u$ in $L^1(\Omega, \mathbf{R}^m)$ and $\nabla u^\nu \rightharpoonup \nabla u$ weakly-* in $L^\infty(\Omega, \mathbf{R}^{n \times m})$) if and only if f is quasiconvex, which is by definition

$$\int_Q f(A + \nabla v) dx \geq f(A) \quad (1.2)$$

for every cube $Q \subseteq \mathbf{R}^n$, every matrix $A \in \mathbf{R}^{n \times m}$ and every function $v \in C_0^\infty(Q, \mathbf{R}^m)$. The quasiconvexity condition is very hard to be verified in practice. It is easier to verify the so-called rank-one convexity condition which is a consequence of quasiconvexity. This condition means that every mapping of the form $\mathbf{R} \ni t \mapsto f(A + ta \otimes b)$ is convex for arbitrary $A \in \mathbf{R}^{n \times m}$ and arbitrary rank one matrix $a \otimes b = (a_i b_j)_{i=1, \dots, n; j=1, \dots, m} \in \mathbf{R}^{n \times m}$. (see e.g. [17],[51],[52],[55],[64]). Those two notions agree if $\min\{n, m\} = 1$, in such cases every quasiconvex function is even convex, or when f is a quadratic form (see e.g. [17],[51], [53, Section 3], [78, Theorem 11]).

It has been conjectured by Morrey in 1952 ([51]) that rank-one convexity does not imply quasiconvexity. This conjecture has been confirmed by Šverák 40 years later in [74] in dimensions $n \geq 2, m \geq 3$ with the example of a polynomial of degree four which is rank-one convex but not quasiconvex. The conjecture is still open in the remaining dimensions $n \geq 2, m = 2$.

However, an alternative to (1.2) algebraic description of quasiconvex functions is known (see [14]), and some numerical approaches to face Morrey's rank-one conjecture are known (see e.g. [18], [19]), but it is still not possible in general to verify it in practice. There are so far few ways to investigate the quasiconvexity condition directly. It is known that in dimensions $n \geq 2, m \geq 3$ this condition is nonlocal (see [40]). In the same dimensions it is also not invariant with respect to the transposition ([41],[57]). For some other related approaches we refer e.g to [1],[5],[10],[25],[33],[34],[49],[53],[63],[67],[68],[71],[72],[73],[75],[81],[82], and references therein. None of the above mentioned properties except the rank-one condition can be described in geometric way.

Recently, the second author has found necessary geometric conditions for quasiconvexity satisfied on certain three dimensional simplexes in $\mathbf{R}^{n \times m}$, [34]. Roughly speaking these conditions, defined as tetrahedral convexity conditions, express the following property: if f agrees with a certain polynomial A on the tetrahedron D from certain class of tetrahedrons in $\mathbf{R}^{n \times m}$ on its vertices and three more other points (where the polynomial and those points are determined by D) then $f \leq A$ inside D . Then the natural question is whether we can expect similar geometric conditions holding on four dimensional simplexes in $\mathbf{R}^{n \times m}$. In particular in the case $n = m = 2$ the dimension of the simplex is the same as the dimension of the domain of f . In this paper we find such conditions. Both mentioned geometric conditions (three and four dimensional) are familiar to the usual convexity conditions, as every convex function which agrees with an affine function in the endpoints of the interval is not bigger than this affine function in its interior. Obviously an interval is a one-dimensional simplex. Note also that our conditions are convenient for a numerical treatment ([35]), and they are between quasiconvexity and rank-one convexity.

Let us mention that our geometric conditions generalize the so-called Λ -convexity conditions due to Murat and Tartar appearing in the more general problem. Let $P =$

$(P_1, \dots, P_N) : C^\infty(\Omega, \mathbf{R}^m) \rightarrow C^\infty(\Omega, \mathbf{R}^N)$ be a differential operator with constant coefficients, given by

$$P_k u = \sum_{i=1, \dots, n, j=1, \dots, m} a_{i,j}^k \frac{\partial u_j}{\partial x_i}, \quad k = 1, \dots, N, \quad (1.3)$$

and let $f : \mathbf{R}^m \rightarrow \mathbf{R}$ be a continuous function. Consider instead of (1.1) the functional

$$I_f(u) = \int_{\Omega} f(u(x)) dx, \quad u \in \text{Ker} P \cap L^\infty(\Omega, \mathbf{R}^m), \quad (1.4)$$

where $\text{Ker} P$ is the distributional kernel of the operator P .

In particular, when $P = \text{curl}$ is applied to each column of u (and $u \in \mathbf{R}_n^k$, $k \cdot n = m$) in a simply connected domain, we recover the classical functional of the calculus of variations.

In general the necessary and sufficient conditions for a function f to define a lower semicontinuous functional with respect to sequential weak- $*$ -convergence in $L^\infty(\Omega, \mathbf{R}^m) \cap \text{Ker} P$ are not known (we refer e.g. to [53, Section 3], [78, Theorem 11], [16, page 26], [78], [53], [65], [33], [56], [78], [80], [25] for special cases). The only known general condition is so-called Λ -convexity necessary condition due to Murat and Tartar (see e.g. [53, Theorem 2.1], [16, Theorem 3.1], [64, Theorem 10.1], [79, Corollary 9]). It reads as follows.

Theorem 1.1 *Define*

$$V = \{(\xi, \lambda) : \xi \in \mathbf{R}^n, \xi \neq 0, \lambda \in \mathbf{R}^m, \sum_{i,j} a_{i,j}^k \xi_i \lambda_j = 0, \text{ for } k = 0, \dots, N\},$$

$$\Lambda = \{\lambda \in \mathbf{R}^m : \text{there exists } \xi \in \mathbf{R}^n, \xi \neq 0, \text{ such that } (\xi, \lambda) \in V\}.$$

If I_f given by (1.4) is lower semicontinuous (continuous) with respect to L^∞ -weak $$ -convergence in $L^\infty(\Omega, \mathbf{R}^m) \cap \text{Ker} P$, then f is Λ -convex (Λ -affine), which means that for every $A \in \mathbf{R}^m$ and every $\lambda \in \Lambda$ the function $\mathbf{R} \ni t \mapsto f(A + t\lambda)$ is convex (affine).*

We try to contribute to both approaches: the general one and the special variational one. As a result we obtain a general condition stated in Theorem 4.2 and its special case related to the variational approach (Theorem 4.3).

Let us mention that the rank-one problem is strongly related to the important and long standing problems in the theory of quasiconformal mappings, as has been recently pointed out by Iwaniec and Astala ([29],[4]).

The paper is organized as follows. At first we study functionals of the form: $I_f(u) = \int_{[0,1]^2} f(u_1(\tau_1), u_2(\tau_2), u_3(\tau_1 + \tau_2), u_4(\tau_1 - \tau_2)) d\tau_1 d\tau_2$ and after some preparations made in Section 2 and 3 we see that if I_f is lower semicontinuous then f satisfies certain conditions on four dimensional simplexes in \mathbf{R}^4 . Then the general functional given by (1.4) is reduced to that special one after restricting the general functional to subspaces in the kernel of the operator P of the form $\{A + \sum_{i=1}^4 u_i(\langle x, \xi \rangle) \lambda_i\}$, where $(\xi_i, \lambda_i) \in V$ and V is defined in Theorem 1.1. The main result and the discussion are presented in Sections 4 and 5 respectively.

The results of this paper and that of [34] are the continuation of the program started by the second author mainly in [33] where she studied functionals of the general form (1.4) for systems like (1.3) where the kernel of P is the solution of the system of equations like

$$\partial_{V_j} u_j = 0 \text{ for } j = 1, \dots, m, \quad (1.5)$$

where V_1, \dots, V_m are linear subspaces of \mathbf{R}^n and $\partial_V u = 0$ means that for every $v \in V$ we have $\partial_v u = 0$. It has been pointed out in [33] that the knowledge about lower semicontinuous functionals on solutions of (1.5) should yield new conditions in the Compensated Compactness Theory and Calculus of Variations. This is because subspaces like (1.5) can be found in kernels of generally defined operators like (1.3), so we can restrict our general functional to such subspaces. The main result of [34] is based on investigation of functional like $I_f(u) = \int_{[0,1]^2} f(u_1(\tau_1), u_2(\tau_2), u_3(\tau_1 + \tau_2)) d\tau_1 d\tau_2$, while here the special role is played by the functional $I_f(u) = \int_{[0,1]^2} f(u_1(\tau_1), u_2(\tau_2), u_3(\tau_1 + \tau_2), u_4(\tau_1 - \tau_2)) d\tau_1 d\tau_2$. Both studied models bring new geometric conditions for lower semicontinuous functionals in the general model (1.4).

We believe that the number of various versions of geometric convexity-like conditions for quasiconvex functions like ours will decrease with the time. They require systematic investigation. Perhaps one of them will lead to the confirmation of Morrey's conjecture in some cases, or perhaps impossibility to find an example of a rank-one convex function which does not satisfy a geometric condition in the remaining cases of the rank-one conjecture of Morray or will encourage someone to find the proof that quasiconvexity is the same as rank-one convexity.

Acknowledgements Part of this research was done during the second author's visit at Konstanz University in December 2002. This author would like to warmly thank Konstanz University for hospitality. We also would like to thank Pietro Celada for the argument presented in Remark 4.5 and to Bernd Kirchheim and Martin Kružík for helpful conversations.

2 Notation and preliminaries

2.1 The basic notation

For a measurable function $u : [0, 1) \rightarrow \mathbf{R}$ we denote by \hat{u} its periodic extension outside $[0, 1)$. If $A \subseteq [0, 1)$ is any subset, we write $A^1 = A$ and $A^0 = [0, 1) \setminus A$. Let $[s]_1$ stand for an integer part of $s \in \mathbf{R}$. In the sequel we will assume that $\Omega \subseteq \mathbf{R}^n$ is an open bounded domain. If A is a measurable subset of \mathbf{R}^n , by $|A|$ we denote its Lebesgue's measure. By n -dimensional cube in \mathbf{R}^n we mean an arbitrary set of the form $I_1 \times \dots \times I_n$ where the I_k 's are closed intervals. If X is a topological space, by $C(X)$ we denote the space of continuous functions on X . By e_1, \dots, e_n we denote the standard basis in \mathbf{R}^n , while $\langle \cdot, \cdot \rangle$ stands for the inner product. If $\xi \in \mathbf{R}^n$ and $a \in \mathbf{R}^m$ by $\xi \otimes a$ we denote the rank one matrix $(\xi_i a_j)_{i=1, \dots, n, j=1, \dots, m}$. By χ_B we denote the characteristic function of the set B . Let $j = (j_1, \dots, j_k) \in \{0, 1\}^k$. We describe its length by $|j| = \sum_{i=1}^k j_i$. If H is a given group of transformations of \mathbf{R}^n and $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is a mapping, we denote $f_h(x) := f(hx)$. By $\mathbf{R}^{n \times m}$ we denote the space of $n \times m$ matrices.

2.2 Combinatorial objects

In this subsection we introduce the space \mathcal{S}^k , and some operations on this space. This space will play an essential role subsequently. Although our point of interest is the case

$k \in \{3, 4\}$ only, the calculations for this special case are not essentially shorter.

The \mathcal{S}^k spaces.

By \mathcal{S}^k we define the space of sequences with indices in $\{0, 1\}^k$. As $\{0, 1\}^k$ consists of 2^k elements, it follows that the space \mathcal{S}^k is isomorphic to \mathbf{R}^N with $N = 2^k$. Let $i \in \{1, \dots, k-1\}$, $\epsilon \in \{0, 1\}$, and let us define the transformation of indices $s_i^\epsilon : \{0, 1\}^{k-1} \rightarrow \{0, 1\}^k$ by putting ϵ on the i -th place:

$$(s_i^\epsilon(j))_s = \begin{cases} j_k & \text{for } s < i \\ \epsilon & \text{for } s = i \\ j_{k-1} & \text{for } s \geq i+1. \end{cases} \quad (2.6)$$

For example if $j = (1, 0, 1)$, $i = 2$ and $\epsilon = 0$, we have $s_2^0(j) = (1, 0, 0, 1)$.

Then we define the two related operators $\Pi_i^\epsilon, \Pi_i : \mathcal{S}^k \rightarrow \mathcal{S}^{k-1}$ by

$$(\Pi_i^\epsilon\{h\})_j = h_{s_i^\epsilon(j)}, \quad \Pi_i\{h\} = \Pi_i^0 + \Pi_i^1.$$

For example when $i = k$ we have $(\Pi_k\{h\})_l = h_{(l,0)} + h_{(l,1)}$, for every $l \in \{0, 1\}^{k-1}$. Note that the operator Π_i can be regarded as discrete directional integration of $\{h\}$ in the given i -th direction.

We denote

$$\begin{aligned} \bigcirc_{i \in \{r_1, \dots, r_t\}} \Pi_i &= \sum_{j: j_{r_1} = a_1, \dots, j_{r_t} = a_t, a_1, \dots, a_t \in \{0, 1\}} h_j \in \mathcal{S}^{k-t} \\ \bigcirc_{i \in \{r_1, \dots, r_t\}} \Pi_i^\epsilon &= \sum_{j: j_{r_1} = \dots = j_{r_t} = \epsilon} h_j \in \mathcal{S}^{k-t}. \end{aligned}$$

More generally, if $A = \{r_1, \dots, r_t\}$, $B = \{l_1, \dots, l_s\}$ and $C = \{t_1, \dots, t_m\}$ are disjoint subsets of $\{1, \dots, k\}$, we set

$$\bigcirc_{i \in A} \Pi_i \bigcirc_{i \in B} \Pi_i^1 \bigcirc_{i \in C} \Pi_i^0 = \sum_{j \in D} h_j \in \mathcal{S}^{k-(r+s+l)}, \quad (2.7)$$

where

$$D = \{j \in \{0, 1\}^k : j_{t_1} = 0, \dots, j_{t_m} = 0, j_{l_1} = 1, \dots, j_{l_s} = 1, j_{r_1}, \dots, j_{r_t} \in \{0, 1\}\}.$$

For example for $k = 3$ we have $\bigcirc_{i \notin \{1, 2\}} \Pi_i \bigcirc_{i \in \{1, 2\}} \Pi_i^1\{h\} = h_{(1,1,0)} + h_{(1,1,1)}$. Note that this notation agrees with composition of operators of the type Π_i and Π_i^ϵ if at first we write numbers $r_1, \dots, r_t, l_1, \dots, l_s, t_1, \dots, t_m$ in increasing order: $a_1 < a_2 < \dots < a_{t+s+m}$ and then consider the composition $\Pi_{a_1}^{n_1} \circ \Pi_{a_2}^{n_2} \circ \dots \circ \Pi_{a_N}^{n_N}$ where $N = t + s + m$ and $\Pi_a^n = \Pi_a$ if $a \in A$, $\Pi_a^n = \Pi_a^1$ if $a \in B$ and $\Pi_a^n = \Pi_a^0$ if $a \in C$. The other compositions may not have sense.

The following lemma summarizes obvious, but useful properties of the operators Π_i and Π_i^ϵ . Its proof is left to the reader as a simple exercise.

Lemma 2.1 1) For every $\{h\} \in \mathcal{S}^k$ we have

$$\bigcirc_{i \in \{1, \dots, k\}} \Pi_i\{h\} = \sum_{j \in \{0, 1\}^k} h_j.$$

2) The operator Π_i preserves the sum: if $\{h\} \in \mathcal{S}^k$ sums up to A , then also $\{\Pi_i\{h\}\}$ sums up to A , which is expressed by

$$\sum_{j \in \{0,1\}^{k-1}} (\Pi_i\{h\})_j = \sum_{j \in \{0,1\}^k} h_j.$$

Remark 2.1 One can easily check that usually there is no equality $\Pi_i \circ \Pi_j = \Pi_j \circ \Pi_i$ for $i \neq j$.

By $\mathcal{S}^k(\Omega)$ we will denote the space of all measurable functions $\{h\} : \Omega \rightarrow \mathcal{S}^k$. If $\{h\} \in \mathcal{S}^k(\Omega)$ is the given function, we use the same expressions: $\Pi_i^\epsilon\{h\}$ and $\Pi^i\{h\}$ to denote functions induced by Π_i^ϵ and Π_i , namely $(\Pi_i^\epsilon\{h\})(x) = \Pi_i^\epsilon\{h(x)\}$ and $(\Pi_i\{h\})(x) = \Pi_i\{h(x)\}$.

2.3 Algebraic and geometric objects

Polynomials and projections.

By \mathcal{A} we denote the 11 dimensional subspace of polynomials in \mathbf{R}^4 spanned by $\{1, x_i, x_i x_j : i, j \in \{1, 2, 3, 4\}, i \neq j\}$. Let us describe the following operator $\mathcal{P} : C(\mathbf{R}^4) \rightarrow \mathcal{A}$ expressed in terms of differences of f : $\Delta_i f(x) = f(x + e_i) - f(x)$:

$$\begin{aligned} Pf(x) &= f(0, 0, 0, 0) + \sum_{i=1}^4 \Delta_i f(0, 0, 0, 0) x_i \\ &+ \sum_{(i,j) \in \{(1,3), (1,4), (2,3)\}} \Delta_i \Delta_j f(0, 0, 0, 0) x_i x_j + \Delta_1 \Delta_2 f(0, 0, 1, 0) x_1 x_2 \\ &+ \Delta_2 \Delta_4 f(1, 0, 1, 0) x_2 x_4 + \left(\frac{1}{2} \Delta_3 \Delta_4 f(0, 0, 0, 0) + \frac{1}{2} \Delta_3 \Delta_4 f(1, 0, 0, 0) \right) x_3 x_4 \\ &:= Qf(x) + \frac{1}{2} (R_1 f(x) + R_2 f(x)), \end{aligned} \tag{2.8}$$

where $x = (x_1, x_2, x_3, x_4)$, $R_1 f(x) = \Delta_3 \Delta_4 f(0, 0, 0, 0) x_3 x_4$, the projectors R_1, R_2 are defined by $R_2 f(x) = \Delta_3 \Delta_4 f(1, 0, 0, 0) x_3 x_4$, and $Qf(x)$ is the remaining term in the expression above.

We have the following lemma. Its simple proof is left to the reader.

Lemma 2.2 1) The operator Pf coincides with f in the following 9 vertices of the cube $Q = [0, 1]^4$: $(0, 0, 0, 0)$, $(1, 0, 0, 0)$, $(0, 1, 0, 0)$, $(0, 0, 1, 0)$, $(0, 0, 0, 1)$, $(1, 0, 1, 0)$, $(1, 0, 0, 1)$, $(0, 1, 1, 0)$, $(1, 1, 1, 0)$.

2) Pf depends only on values of f in 12 vertices of the cube Q : nine vertices described above and: $(1, 0, 1, 1)$, $(1, 1, 1, 1)$, $(0, 0, 1, 1)$.

3) If $f \in \mathcal{A}$ we have $Pf = f$, in particular Pf is the projection operator onto the space \mathcal{A} .

Remark 2.2 Note that the class of projection operators from $C(\mathbf{R}^4)$ to \mathcal{A} is large. For example we can define it by taking an arbitrary $f \in C(\mathbf{R}^4)$ and prescribing to it the uniquely defined polynomial which agrees with f in the following 11 vertices of the cube Q : the first one taken arbitrary and the remaining ones linked with the first one by at most two edges of the cube. As convex combination of arbitrary two projection operators is again a projection operator, it follows that the set of projection operators from $C(\mathbf{R}^4)$ to \mathcal{A} is convex.

Remark 2.3 Let us look at the projection operator Pf from Lemma 2.2 more closely. Note that Pf is the convex combination of two projection operators: $P_1f = Qf + R_1f$, and $P_2f = Qf + R_2f$. The operator P_2 agrees with f in the given 11 vertices of the cube Q : 9 vertices described in part 1) of Lemma 2.2 and two more: $(1, 0, 1, 1)$ and $(1, 1, 1, 1)$. The first operator agrees with f in 10 vertices of the cube Q only: the 9 described in part 1) of Lemma 2.2 and in $(0, 0, 1, 1)$.

Special group of invariances.

We will consider the group G of linear transformations of \mathbf{R}^4 generated by the following ones:

- translations: $y \mapsto b + y$ where $b, y \in \mathbf{R}^4$,
- dilations: $y = (y_1, y_2, y_3, y_4) \mapsto (t_1y_1, t_2y_2, t_3y_3, t_4y_4)$ where $t_1, \dots, t_4 \in \mathbf{R}$
- permutations $\pi_{1,2}$ and $\pi_{3,4}$ defined by $\pi_{1,2}(x_1, x_2, x_3, x_4) = (x_2, x_1, x_3, x_4)$, and $\pi_{3,4}(x_1, x_2, x_3, x_4) = (x_1, x_2, x_4, x_3)$.

The group G described above will be called the *special group of invariances*.

We will consider also its subgroup \tilde{G} consisting of isometries of $[0, 1]^4$. It is generated by the following 6 transformations: 4 symmetries $s_i: x_i \mapsto 1 - x_i$ where $i \in \{1, \dots, 4\}$ and two permutations: π_{12} and π_{34} (note that every its element is of order 2). It is easy to see that the subgroup H_1 of \tilde{G} generated by two symmetries s_1, s_2 , and permutation π_{12} is the normal subgroup of \tilde{G} . Moreover, the quotient group \tilde{G}/H_1 is isomorphic to the subgroup H_2 of \tilde{G} generated by symmetries s_3, s_4 and the permutation π_{34} , and both subgroups H_1 and H_2 consist of 8 elements. Then \tilde{G} is isomorphic to $\tilde{G}/H_1 \otimes H_1$, hence it is isomorphic to $H_2 \otimes H_1$ (see e.g. [43, Proposition 1 on page 12]). This implies that the group \tilde{G} consists of 64 elements.

The principal simplex and its G -similar cousins.

Special role in our development will play the following simplex in \mathbf{R}^4 :

$$R_1 = \{x \in Q : x_3 > x_1 + x_2, x_1 \leq 1/2, 2x_1 > x_3 + x_4, x_2 \geq 0, x_4 \geq 0\}. \quad (2.9)$$

The simplex R_1 will be called the *principal simplex*.

Let H be the given group of transformations of \mathbf{R}^4 and $D \subseteq \mathbf{R}^4$ be the subset. We will say that the subset $\tilde{D} \subseteq \mathbf{R}^4$ is H -similar to D if there is such $h \in H$ that $\tilde{D} = h(D)$. Let us denote by $Caus(D, H)$ the set of all H -similar to D subsets of \mathbf{R}^4 . This set will be called the set of H -cousins of D . Let us consider the set $Caus(R_1, G)$ where R_1 is the principal simplex and G is the special group of invariances. One may ask what kind of

symplexes will be found there. Obviously, every such simplex is of the form $D \circ Tr(R)$, where R is \tilde{G} -cousin of R_1 (note that the set $Caus(R_1, \tilde{G})$ consists of 64 symplexes), Tr is some translation and D is some dilation in \mathbf{R}^4 . Let us denote vertices of R_1 by: $W_1 = (0, 0, 0, 0)$, $W_2 = (1/2, 0, 1, 0)$, $W_3 = (1/2, 0, 1/2, 0)$, $W_4 = (1/2, 0, 1/2, 1/2)$, and $W_5 = (1/2, 1/2, 1, 0)$ (note that only W_1 is the vertex of the cube $Q = [0, 1]^4$). Then R_1 is the convex Minkowski's combination of two sets: the tetrahedron $T \subseteq \mathbf{R}^4$ spanned by W_2, W_3, W_4, W_5 , and W_1 , which means that $R_1 = \{tx + (1 - t)W_1 : t \in [0, 1], x \in T\}$.




Figure 1 : $T = R_1 \cap \{x_1 = \frac{1}{2}\} \subset T_3$

The tetrahedron T lives on the hyperplane $x_1 = 1/2$ (so $x_1 = const$) and has three axes perpendicular to each other: W_2W_5 is parallel to e_2 , W_2W_3 is parallel to e_3 , W_3W_4 is parallel to e_4 . Those axes form the polyline in \mathbf{R}^4 : $W_5W_2W_3W_4$ which uniquely defines the simplex T . Now if we apply the dilation on \mathbf{R}^4 we see that the new polyline obtained by this dilation again has three axes parallel to e_2, e_3, e_4 respectively, moreover, the new tetrahedron obtained from T by dilation also lives on the hyperplane $x_1 = const$. Obviously, those mentioned properties: $R_1 = conv\{W, T\}$, where a) W is one of the vertices of Q and b) T lives on the hyperplane $x_i = const$ for some i and has three axes perpendicular to each other, remain unchanged under the action of G (where instead of Q we consider its G -cousin).

2.4 The functional setting

The special functional

Let $Q = [0, 1]^2$ be the unit cube, and $f : \mathbf{R}^4 \rightarrow \mathbf{R}$ be continuous. The functional

$$I_f(u) = \int_Q f(u_1(x_1), u_2(x_2), u_3(x_1 + x_2), u_4(x_1 - x_2)) dx_1 dx_2, \quad (2.10)$$

where $u_i \in L^\infty(\mathbf{R})$ for $i = 1, 2, 3, 4$ will play an essential role in our investigations. For this reason such functionals will be called *special*.

We have the following lemma.

Lemma 2.3 Let $f \in C(\mathbf{R}^4)$, $\xi_1, \xi_2 \in \mathbf{R}^n$ be the given two independent vectors, $\xi = (\xi_1, \xi_2)$, and consider the functional

$$J_{f,\Omega,\xi}(u) = \int_{\Omega} f(u_1(\langle x, \xi_1 \rangle), u_2(\langle x, \xi_2 \rangle), u_3(\langle x, \xi_1 + \xi_2 \rangle), u_4(\langle x, \xi_1 - \xi_2 \rangle)) dx, \quad (2.11)$$

where $u_i \in L^\infty(\mathbf{R})$ for $i = 1, \dots, 4$. The following statements are equivalent:

- 1) For an arbitrary bounded domain $\Omega \subseteq \mathbf{R}^n$ with $n \geq 2$ the functional $J_{f,\Omega,\xi}(\cdot)$ given by (2.11) is lower semicontinuous with respect to the weak * convergence of the u_i 's in $L^\infty(\mathbf{R})$.
- 2) The special functional $I_f(\cdot)$ given by (2.10) is lower semicontinuous with respect to the weak * convergence of the u_i 's in $L^\infty(\mathbf{R})$.

Proof. We use almost the same arguments as that in [34], but for the reader's convenience we include the proof of the lemma. The implication **1**) \Rightarrow **2**) is obvious. Hence we prove the implication **2**) \Rightarrow **1**) only. By an easy approximation argument it suffices to prove **1**) for an arbitrary domain Ω with the smooth boundary. Note that if **2**) holds then Q can be replaced by any cube with edges parallel do the axes. Let us set $y_1 = \langle x, \xi_1 \rangle$ and $y_2 = \langle x, \xi_2 \rangle$. The functional $J_{f,\Omega,\xi}$ is lower semicontinuous if and only if

$$J_f(u) = \int_{\tilde{\Omega}} |\Omega_{(y_1, y_2)}| f(u_1(y_1), u_2(y_2), u_3(y_1 + y_2), u_4(y_1 - y_2)) dy_1 dy_2$$

is lower semicontinuous, where $\Omega_{(y_1, y_2)} = \{x \in \Omega : y_1 = \langle x, \xi_1 \rangle \text{ and } y_2 = \langle x, \xi_2 \rangle\}$, $|\Omega_{(y_1, y_2)}|$ is its $n - 2$ dimensional Hausdorff measure and $\tilde{\Omega} = \{(y_1, y_2) \in \mathbf{R}^2 : \Omega_{(y_1, y_2)} \neq \emptyset\}$. Let us cover the set $\tilde{\Omega}$ by countable family of disjoint cubes $Q_i = Q_i(y^i)$ where $y^i = (y_1^i, y_2^i)$ whose edges are parallel to the axes and consider the functional

$$J_f^N(u) = \sum_{i=1}^N |\Omega_{(y^i)}| \int_{Q_i} f(u_1(y_1), u_2(y_2), u_3(y_1 + y_2), u_4(y_1 - y_2)) dy_1 dy_2.$$

If the diameter of each Q_i is small enough, then the continuity of f and continuity of the mapping $y \mapsto |\Omega_{(y)}|$ yields: for all $\varepsilon > 0$ and N sufficiently large

$$|J_f^N(u) - J_f(u)| < \varepsilon$$

for all functions $u \in L^\infty$ with $\|u\|_\infty \leq R$. This and the lower semicontinuity of J_f^N complete the proof. \square

Now we are going to concentrate on obtaining some basic properties of special functionals. Let \mathcal{M} be the set of all continuous functions on \mathbf{R}^4 which define lower semicontinuous special functional, and \mathcal{C} be the space of those integrands which define weakly continuous special functionals (with respect to the sequential weak-* convergence of the u_i 's in L^∞). One can easily check that the following property holds.

Lemma 2.4 The set \mathcal{M} is invariant with respect to the action of the special group of invariances G (see Section 2.3). This means that if $g \in G$ is taken arbitrary and $f \in \mathcal{M}$ then the mapping $(f_g(y) = f(gy))$ also belongs to \mathcal{M} .

Remark 2.4 We will see later that $\mathcal{C} = \mathcal{A}$.

The following lemma describes the restriction of special functional to the set of periodic extensions of characteristic functions of cubes in \mathbf{R}^4 .

Lemma 2.5 *Let $x = (x_1, x_2, x_3, x_4) \in Q = [0, 1]^4$, $u_i(\tau, x) = \hat{\chi}_{[0, x_i]}(\tau)$ where $\tau \in \mathbf{R}^2$, $i \in \{1, 2, 3, 4\}$ and $u : \mathbf{R}^2 \times Q \rightarrow \mathbf{R}^4$ be defined by $u(\tau, x) = (u_1(\tau_1), u_2(\tau_2), u_3(\tau_1 + \tau_2), u_4(\tau_1 - \tau_2))$. Set for $j \in \{0, 1\}^4$ $h_j(x) =$*

$$|\{(\tau_1, \tau_2) \in [0, 1]^2 : \tau_1 \in [0, x_1]^{j_1}, \tau_2 \in [0, x_2]^{j_2}, [\tau_1 + \tau_2]_1 \in [0, x_3]^{j_3}, [\tau_1 - \tau_2]_1 \in [0, x_4]^{j_4}\}|.$$

Then the following properties hold.

i) *For arbitrary continuous function $f : \mathbf{R}^4 \times \mathbf{R}$ we have*

$$\int_{[0, 1]^2} f(u(\tau, x)) d\tau = \sum_{j \in \{0, 1\}^4} h_j(x) f(j).$$

ii) *Functions $\{h_j(x)\}$ define the distribution of the probability measure concentrated in 16 points $j \in \{0, 1\}^4$, in particular all the h_j 's are nonnegative.*

iii) *For every $j \in \{0, 1\}^4$ the mapping $x \mapsto h_j(x)$ is continuous.*

Some further properties of functions $\{h_j(x)\}$ and the computation of their values in selected subregions of the cube Q will be presented in Section 3. From now functions $\{h_j(x)\}$ and their shiftings $\{h_j^i(x)\} = \{\Pi_i h_j(x)\}$ will be called *special distributions of measures* (note that the shiftings also define distributions of probability measures).

The key point in our argumentation will be the following lemma.

Lemma 2.6 *Assume that $f \in C(\mathbf{R}^4)$ and f defines the special functional, which is lower semicontinuous with respect to the weak-* convergence of the u_i 's in $L^\infty(\Omega)$. Then if u_1, u_2, u_3, u_4 are bounded and periodic of period 1, we have*

$$\int_{[0, 1]^2} f(u_1(\tau_1), u_2(\tau_2), u_3(\tau_1 + \tau_2), u_4(\tau_1 - \tau_2)) d\tau_1 d\tau_2 \geq f\left(\int_0^1 u_1(\tau) d\tau, \dots, \int_0^1 u_4(\tau) d\tau\right).$$

Proof. This follows from the Riemann–Lebesgue's Theorem applied to functions $u^\nu(x) = u(\nu x)$, where $u = (u_1, u_2, u_3, u_4)$ and $f^\nu(x) = f(u^\nu(x))$ (see e.g. Lemma 1.2 on page 8 in [16]). It only remains to check that $\int_{[0, 1]^2} u_3(\tau_1 + \tau_2) d\tau_1 d\tau_2 = \int_0^1 u_3(\tau) d\tau$ and $\int_{[0, 1]^2} u_4(\tau_1 - \tau_2) d\tau_1 d\tau_2 = \int_0^1 u_4(\tau) d\tau$. \square

3 Special distributions of measures

3.1 Invariances

Through this section we assume that $Q = [0, 1]^4$ is the unit cube and h_j 's are the same as in Lemma 2.5. Let $\{h_j^i(x)\}_{j \in \{0,1\}^3} = \{\Pi_i h_j(x)\}_{j \in \{0,1\}^3}$. Note that $h_j^i(x)$ depends on three coordinates of x only, so we will treat it as the function defined on $[0, 1]^3$. On the other hand in some places we will denote coordinates of $x \in [0, 1]^3$ in $h_j^i(x)$ in the following way: in $h_j^1(x)$ by (x_2, x_3, x_4) , in $h_j^2(x)$ by (x_1, x_3, x_4) , in $h_j^3(x)$ by (x_1, x_2, x_4) , in $h_j^4(x)$ by (x_1, x_2, x_3) to indicate that in fact h_j^i 's are functions of four variables, but we forget about the given one. The choice of the notation will be obvious from the context.

Let $k \in \mathbf{N}$ and $A : \mathbf{R}^k \rightarrow \mathbf{R}^k$ be an arbitrary affine transformation such that A restricted to $\{0, 1\}^k$ is the bijection. Then A preserves the whole cube $Q = [0, 1]^k$, and A defines the mapping $\mathcal{S}^k(Q) \rightarrow \mathcal{S}^k(Q)$ by expression

$$A\{h_j(x)\}_{j \in \{0,1\}^k} = \{h_{Aj}(Ax)\}_{j \in \{0,1\}^k}. \quad (3.12)$$

Our goal is to look for those affine transformations of \mathbf{R}^3 and \mathbf{R}^4 which leave the distributions $\{h_j^i(x)\}_{j \in \{0,1\}^3}$ and $\{h_j(x)\}_{j \in \{0,1\}^4}$ invariant. We start with the following lemma.

Lemma 3.1 (Lemma about invariances) *The distributions*

$\{h_j^i(x)\}_{j \in \{0,1\}^3}$'s and $\{h_j(x)\}_{j \in \{0,1\}^4}$ are invariant with respect to the following isometries of \mathbf{R}^3 and \mathbf{R}^4 respectively.

- 1) $\{h_j^4(x)\}_{j \in \{0,1\}^3}$ is invariant under the permutation of axes $\pi_{12}(x_1, x_2, x_3) = (x_2, x_1, x_3)$, and symmetry with respect to the point $(1/2, 1/2, 1/2)$, $S(x_1, x_2, x_3) = (1 - x_1, 1 - x_2, 1 - x_3)$.
- 2) $\{h_j^1(x)\}_{j \in \{0,1\}^3}$ is invariant with respect to isometries of \mathbf{R}^3 , $B_1, B_2 : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ given by $B_1(x_1, x_2, x_3) = (x_1, 1 - x_3, 1 - x_2)$, $B_2(x_1, x_2, x_3) = (1 - x_1, x_3, x_2)$.
- 3) Let $C_3 : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ be an affine isometry given by $C_3(x_1, x_2, x_3) = (x_1, 1 - x_2, x_3)$. Then for every $x \in [0, 1]^3$ and every $j \in \{0, 1\}^3$ we have

$$h_j^3(x) = h_{C_3j}^4(C_3x).$$

In particular $\{h_j^3(x)\}_{j \in \{0,1\}^3}$ is invariant with respect to mappings

$$D_1(x_1, x_2, x_3) = C_3\pi_{12}C_3(x_1, x_2, x_3) = (1 - x_2, 1 - x_1, x_3) \text{ and } S.$$

- 4) Let $C_1 : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ be an affine isometry given by $C_1(x_1, x_2, x_3) = (x_1, x_2, 1 - x_3)$. Then for every $x \in [0, 1]^3$ and every $j \in \{0, 1\}^3$ we have

$$h_j^2(x) = h_{C_1j}^1(C_1x).$$

In particular $\{h_j^2(x)\}_{j \in \{0,1\}^3}$ is invariant with respect to the following permutation of axes $\pi_{23}(x_1, x_2, x_3) = (x_1, x_3, x_2)$ and the mapping $S \circ \pi_{23}$.

5) $\{h_j(x)\}_{j \in \{0,1\}^4}$ is invariant with respect to orientation preserving isometries of \mathbf{R}^4 given by

$$A_1(x) = (1 - x_1, x_2, 1 - x_4, 1 - x_3), \quad A_2(x) = (x_2, x_1, x_3, 1 - x_4),$$

where $x = (x_1, x_2, x_3, x_4)$.

Proof. This follows from the following changes of variables in the calculation of $\{h_j\}$'s: 1) $\bar{\tau}_1 = \tau_2, \bar{\tau}_2 = \tau_1$, and $\bar{\tau}_1 = 1 - \tau_1, \bar{\tau}_2 = 1 - \tau_2$; 2) $\bar{\tau}_1 = 1 - \tau_1, \bar{\tau}_2 = \tau_2$, and $\bar{\tau}_1 = \tau_1, \bar{\tau}_2 = 1 - \tau_2$; 3) $\bar{\tau}_1 = \tau_1, \bar{\tau}_2 = 1 - \tau_2$; 4) $\bar{\tau}_1 = \tau_2, \bar{\tau}_2 = \tau_1$; 5) $\bar{\tau}_1 = 1 - \tau_1, \bar{\tau}_2 = \tau_2$ and $\bar{\tau}_1 = \tau_2, \bar{\tau}_2 = \tau_1$. \square

3.2 Special isomorphism

We are now showing a possibly shorter way to calculate special distributions of measures. Our techniques will be based on the lemma about isomorphism presented below. Before we state it, let us introduce some useful objects.

The \mathcal{T}^k spaces.

Let $r, k \in \mathbf{N}$ and $r \leq k$. By $s(r, k)$ we denote the set of all subsets of $\{1, \dots, k\}$ consisting of r elements; we will identify this subsets with ordered sequences: $l_1 \dots l_r$ where $1 \leq l_1 < \dots < l_r \leq k$. This set consists of $\binom{k}{r}$ elements.

By $\mathcal{T}^{r,k}$ we denote all functions from $s(r, k)$ to \mathbf{R} , that is all finite sequences $\{t_s\}_{s \in s(r,k)}$. Note that the space $\mathcal{T}^{r,k}$ is isomorphic to \mathbf{R}^N with $N = \binom{k}{r}$, in particular $\mathcal{T}^{k,k}$ can be identified with \mathbf{R} . For our convenience we also denote $\mathcal{T}^{0,k} = \mathbf{R}$.

Let $j = (j_1, \dots, j_k) \in \{0, 1\}^k$. If $|j| = r$ and j has 1 on place t_1, \dots, t_r where $t_1 < t_2 < \dots < t_r$, we denote $d(j) = \{t_1, \dots, t_r\} \in s(r, k)$; in particular every $j \in \{0, 1\}^k$ determines uniquely an element of $s(|j|, k)$.

Let us introduce the special subsets of $s(r, k)$:

$$s(r, k, j) = \{s \in s(r, k) : d(j) \subset s\}.$$

(note that $s(r, k, j) = \emptyset$ if $|j| > r$), and define the following operations on the space $\mathcal{T}^{r,k}$: if $A = \{A_s\}_{s \in s(r,k)} \in \mathcal{T}^{r,k}$, we set

$$\sum A = \sum_{s \in s(r,k)} A_s, \quad \sum^{+j} A = \sum_{s \in s(r,k,j)} A_s \quad (3.13)$$

(if $|j| > r$ the second sum is zero).

We set $\mathcal{T}^k = \mathcal{T}^{0,k} \times \mathcal{T}^{1,k} \times \dots \times \mathcal{T}^{k,k}$. As this space is isomorphic to \mathbf{R}^N with $N = 2^k$, it follows that it is also isomorphic to \mathcal{S}^k . Elements $\{A\} \in \mathcal{T}^k$ will be identified with long vectors $\{A\} = \{A^r\}_{r=0}^k$ where $A^r = \{A_s^r\}_{s \in s(r,k)} \in \mathcal{T}^{r,k}$.

Now we are in the position to state the following lemma.

Lemma 3.2 (Lemma about isomorphism) Let $k \in \mathbb{N}$, $\{A\} = \{A^r\}_{r=0,\dots,k} \in \mathcal{T}^k$, \bigcirc and \sum^{+j} be as in Section 2.2 and (3.13), and consider the system of 2^k linear equations with unknown $\{h\} \in \mathcal{S}^k$:

$$\begin{cases} \bigcirc_{i \in \{1,\dots,k\}} \Pi_i \{h\} = A^0, \\ \bigcirc_{i \notin s} \Pi_i \bigcirc_{i \in s} \Pi_i^1 \{h\} = A_s^r, \quad r \in \{1, \dots, k\}, \quad s \in s(r, k). \end{cases} \quad (3.14)$$

The solutions to this system are uniquely determined by the formulae

$$h_j = (-1)^{|j|} \sum_{r=|j|}^k (-1)^r \sum^{+j} A^r. \quad (3.15)$$

Proof. Let $B : \mathcal{S}^k \rightarrow \mathcal{T}^k$ be the linear mapping associated to the system (3.14), so that (3.14) reads as $B\{h\} = \{A\}$. Let $j_0 \in \{0, 1\}^k$, and $\{\delta_{j_0}\} \in \mathcal{S}^k$ be such that $(\delta_{j_0})_j = 0$ if $j \neq j_0$ and $(\delta_{j_0})_j = 1$ if $j = j_0$. Obviously, vectors $\{\delta_j\}_{j \in \{0,1\}^k}$ form the basis in \mathcal{S}^k . Moreover, an easy calculation shows that $B\{\delta_{j_0}\} = \{A(j_0)\}$ where

$$(A(j_0))_s^r = \begin{cases} 1 & \text{if } r = 0 \\ 1 & \text{if } r \geq 1 \text{ and } s \subseteq d(j_0) \\ 0 & \text{if } r \geq 1 \text{ and } s \not\subseteq d(j_0), \end{cases}$$

for $r \in \{0, \dots, k\}$ and $s \in s(r, k)$, in particular vectors $\{A(j_0)\}_{j_0 \in \{0,1\}^k}$ are linearly independent. This implies that B is the isomorphism. To justify that the formulae (3.15) holds true at first we write it in the form: $C\{A\} = \{h\}$ where $\{A\} \in \mathcal{T}^k$ and $\{h\} \in \mathcal{S}^k$; then we show that $C = B^{-1}$. This will be done if we prove that

$$C\{A(j_0)\} = \{\delta_{j_0}\} \text{ for every } j_0 \in \{0, 1\}^k, \quad (3.16)$$

as this implies that C agrees with B^{-1} on the basis $\{A(j_0)\}_{j_0 \in \{0,1\}^k}$ of \mathcal{T}^k . The justification of (3.16) follows by easy calculations which are left to the reader. \square

Remark 3.1 For the reader's convenience we list the solutions to the system (3.14) for $k = 3$ and $k = 4$ in tables below, putting for simplicity $A_{123}^3 = A^3$ and $A_{1234}^4 = A^4$.

j	h_j
$(0, 0, 0)$	$A^0 - \sum A^1 + \sum A^2 - A^3$
$d(j) = \{s\}$	$A_s^1 - \sum_{\{j_1, j_2: s \in \{j_1, j_2\}\}} A_{j_1 j_2}^2 + A^3$
$d(j) = \{s_1, s_2\}$	$A_{s_1 s_2}^2 - A^3$
$j = (1, 1, 1)$	A^3

Solutions for $k = 3$.

j	h_j
$(0, 0, 0, 0)$	$A^0 - \sum A^1 + \sum A^2 - \sum A^3 + A^4$
$d(j) = \{s\}$	$A_s^1 - \sum_{\{j_1, j_2: s \in \{j_1, j_2\}\}} A_{j_1 j_2}^2 + \sum_{\{j_1, j_2, j_3: s \in \{j_1, j_2, j_3\}\}} A_{j_1 j_2 j_3}^3 - A^4$
$d(j) = \{s_1, s_2\}$	$A_{s_1 s_2}^2 - \sum_{\{j_1, j_2, j_3: s_1, s_2 \in \{j_1, j_2, j_3\}\}} A_{j_1 j_2 j_3}^3 + A^4$
$d(j) = \{s_1, s_2, s_3\}$	$A_{s_1 s_2 s_3}^3 - A^4$
$j = (1, 1, 1, 1)$	A^4

Solutions for $k = 4$.

Remark 3.2 However for every $j \in \{0, 1\}^4$ the function $h_j(x)$ is a piecewise polynomial function of degree not bigger than two, in general it does not belong to the space \mathcal{A} in the region where it behaves polynomially.

Now we will compute the expressions $h_{111}^3(x)$ and $h_{111}^4(x)$, leaving the more complicated calculations for $h_{111}^1(x)$, $h_{111}^2(x)$ and that of $h_{1111}(x)$ to remaining subsections.

Lemma 3.4 *The mapping $x = (x_1, x_2, x_3) \mapsto h_j^4(x)$ belongs to \mathcal{A} for every $j \in \{0, 1\}^3$ if and only if $x \in D_4 = \{(x_1, x_2, x_3) \in [0, 1]^3 : x_1 + x_2 < 1, x_3 > x_1 + x_2\}$ or $x \in S(D_4)$. The respective values of $\{h_j^4(x)\}$ in the sets D_4 and $S(D_4)$ are given in the two tables below.*

j	$(0, 0, 0)$	$(0, 0, 1)$	$(1, 0, 0)$	$(1, 0, 1)$
$h_j^4(x)$	$(1 - x_3)(1 - (x_1 + x_2))$	$x_3(1 - x_1) + x_2(x_1 - x_3)$	$(1 - x_3)x_1$	$(x_3 - x_2)x_1$
j	$(0, 1, 0)$	$(0, 1, 1)$	$(1, 1, 0)$	$(1, 1, 1)$
$h_j^4(x)$	$(1 - x_3)x_2$	$x_2(x_3 - x_1)$	0	x_1x_2

h_j^4 in the set D_4

j	$(0, 0, 0)$	$(0, 0, 1)$	$(1, 0, 0)$	$(1, 0, 1)$
$h_j^4(x)$	$(1 - x_1)(1 - x_2)$	0	$(1 - x_2)(x_1 - x_3)$	$x_3(1 - x_2)$
j	$(0, 1, 0)$	$(0, 1, 1)$	$(1, 1, 0)$	$(1, 1, 1)$
$h_j^4(x)$	$(1 - x_1)(x_2 - x_3)$	$x_3(1 - x_1)$	$(1 - x_2)(x_3 - x_1) + (1 - x_3)x_1$	$x_3(x_1 + x_2 - 1)$

h_j^4 in the set $S(D_4)$

The proof of this fact is given in [34], but for the reader's convenience we include the sketch.

Proof. According to Lemmas 3.2 and 3.3 it suffices to calculate $h_{111}^4(x)$ and use table given in Remark 3.1 for $k = 3$. Then we use the result about invariances (Lemma 3.1), which shows that calculations can be reduced to the region $\{x_1 + x_2 < 1, x_1 > x_2\}$. \square

Our next lemma immediately follows from Lemma 3.1 and Lemma 3.4.

Lemma 3.5 *We have $x = (x_1, x_2, x_4) \mapsto h_j^3(x) \in \mathcal{A}$ for every $j \in \{0, 1\}^3$ if and only if $x \in D_3$ or $x \in S(D_3)$, where $D_3 = \{(x_1, x_2, x_4) \in [0, 1]^3 : x_1 - x_2 < 0, x_4 > x_1 + 1 - x_2\}$ and there we have $h_{111}^3(x) = h_{101}^4(x_1, 1 - x_2, x_4) = (x_4 + x_2 - 1)x_1$, while for $x \in S(D_3)$ the value of $h_{111}^3(x)$ equals x_2x_4 .*

3.4 The computation of h_{111}^1 and h_{111}^2

We are now computing the functions h_{111}^1 and h_{111}^2 in the selected regions of $[0, 1]^3$. The values of h_{111}^1 are obtained with the help of the following lemma.

Lemma 3.6 *The mapping $x = (x_2, x_3, x_4) \mapsto h_{111}^1(x)$ belongs to the space \mathcal{A} if and only if $x \in \bigcup_{k=1}^4 T_k$, where T_k for $k = 1, 2, 3, 4$ are disjoint tetrahedrons*

$$T_1 = \{(x_2, x_3, x_4) \in [0, 1]^3 : x_2 > 1 + \frac{x_3 - x_4}{2}\},$$

$$T_2 = T_1 - (\frac{1}{2}, 0, 0) = \{\frac{1}{2} > x_2 > \frac{1+x_3-x_4}{2}\}, T_3 = S(T_1), T_4 = S(T_2).$$

Under the above notation we have

$$h_{111}^1(x_2, x_3, x_4) = \begin{cases} x_3(x_4 + x_2 - 1) & \text{if } (x_2, x_3, x_4) \in T_1 \\ x_3(x_2 - \frac{1-x_4}{2}) & \text{if } (x_2, x_3, x_4) \in T_2 \\ x_2x_4 & \text{if } (x_2, x_3, x_4) \in T_3 \\ x_4(x_2 - \frac{1-x_3}{2}) & \text{if } (x_2, x_3, x_4) \in T_4. \end{cases}$$

Remark 3.3 The figure presented below shows the position of sets T_i in the cube $[0, 1]^3$.




Figure 3: The tetrahedrons T_1, T_2, T_3, T_4

Proof. The proof follows by steps: 1) We assume that $x_3 \leq x_4, x_3 + x_4 < 1$. 2) We complete the proof of the lemma.

Proof of Step 1: Let us denote for simplicity $I = h_{111}^1(x_2, x_3, x_4)$, and consider the following five sets

$$\begin{aligned} A_0 &= \{(\tau_1, \tau_2) : \tau_1 + \tau_2 < x_3\} \\ A_1 &= \{(\tau_1, \tau_2) : 0 < \tau_1 + \tau_2 - 1 < x_3\} \\ B_0 &= \{(\tau_1, \tau_2) : 0 < \tau_1 - \tau_2 < x_4\} \\ B_1 &= \{(\tau_1, \tau_2) : 0 < 1 + \tau_1 - \tau_2 < x_4\} \\ H_2 &= \{(\tau_1, \tau_2) : \tau_2 \in [0, x_2]\} \end{aligned}$$

Figure 4 represents geometrically the position of the sets A_0, A_1, B_0 and B_1 in the cube $[0, 1]^2$.




Figure 4: The sets A_0, A_1, B_0, B_1

Then $I = I_1 + I_2 + I_3 + I_4$ where $I_1 = |A_0 \cap B_0 \cap H_2|$, $I_2 = |A_0 \cap B_1 \cap H_2|$, $I_3 = |A_1 \cap B_0 \cap H_2|$, $I_4 = |A_1 \cap B_1 \cap H_2|$. As for every $a, b, c, d, \alpha, \beta \in \mathbf{R}$, we have

$$\begin{aligned} \mathcal{L}(a, b, c, d, \alpha, \beta) &= \int_a^b \left(\int_{c+\alpha\tau_2}^{d+\beta\tau_2} d\tau_1 \right) d\tau_2 = \\ &= (b-a) \left[(d-c) + \frac{1}{2}(\beta-\alpha)(b+a) \right]. \end{aligned} \quad (3.17)$$

After easy computation we find that

$$I_1 = \int_0^{\min\{x_2, \frac{x_3}{2}\}} \left(\int_{\tau_2}^{x_3-\tau_2} d\tau_1 \right) d\tau_2 = \begin{cases} x_2 x_3 - x_2^2 & \text{if } x_2 \leq \frac{x_3}{2} \\ \frac{x_3^2}{4} & \text{if } x_2 > \frac{x_3}{2} \end{cases}.$$

To calculate I_2 we observe that $A_0 \cap B_1 = \emptyset$, so immediately $I_2 = 0$.

Let us calculate I_3 . Subtracting the inequalities defining sets A_1 and B_0 we get $1 - x_4 < 2\tau_2$, so for $2x_2 < 1 - x_4$ we obtain $I_3 = 0$. Now, for $x_2 > (1 - x_4)/2$ we conclude that I_3 has the following form

$$I_3 = \int_{\frac{1-x_4}{2}}^{\min\{x_2, \frac{1+x_3}{2}\}} \left(\int_{\max\{\tau_2, 1-\tau_2\}}^{\min\{x_3+1-\tau_2, x_4+\tau_2\}} d\tau_1 \right) d\tau_2.$$

Let us decompose $[0, 1]$ into 5 subintervals $C_i = [a_{i-1}, a_i]$ where $i \in \{1, \dots, 5\}$, with the help of 6 ordered numbers: $a_0 = 0 < a_1 = (1 - x_4)/2 < a_2 = (1 + x_3 - x_4)/2 < a_3 = 1/2 < a_4 = (1 + x_3)/2 < a_5 = 1$. Then

$$\max\{\tau_2, 1 - \tau_2\} = \begin{cases} 1 - \tau_2 & \text{on } C_1 \cup C_2 \cup C_3 \\ \tau_2 & \text{on } C_4 \cup C_5 \end{cases}, \text{ while}$$

$$\min\{x_3 + 1 - \tau_2, x_4 + \tau_2\} = \begin{cases} x_4 + \tau_2 & \text{on } C_1 \cup C_2 \\ x_3 + 1 - \tau_2 & \text{on } C_3 \cup C_4 \cup C_5 \end{cases}.$$

Now using (3.17) we easily compute the values of I_3 on every set C_i and present them in table below.

C_j	$[0, \frac{1-x_4}{2}]$	$[\frac{1-x_4}{2}, \frac{1+x_3-x_4}{2}]$	$[\frac{1+x_3-x_4}{2}, \frac{1}{2}]$	$[\frac{1}{2}, \frac{1+x_3}{2}]$	$[\frac{1+x_3}{2}, 1]$
I_3	0	$(x_2 + \frac{x_4-1}{2})^2$	$-\frac{x_3^2}{4} + x_3(x_2 - \frac{1-x_4}{2})$	$\frac{x_3 x_4}{2} - (\frac{1+x_3}{2} - x_2)^2$	$\frac{x_3 x_4}{2}$

In a similar manner we calculate I_4 . First we observe that for all $(\tau_1, \tau_2) \in A_1 \cap B_1$ we have $2 - x_4 < 2\tau_2$, so for $2x_2 < 2 - x_4$ we have $I_4 = 0$, while for $x_2 > (2 - x_4)/2$

$$I_4 = \int_{1-\frac{x_4}{2}}^{x_2} \left(\int_{1-\tau_2}^{\min\{x_4-1+\tau_2, 1+x_3-\tau_2\}} d\tau_1 \right) d\tau_2.$$

After the simple calculation based on (3.17) we find the values of I_4 on subintervals D_i of $[0, 1]$, and present them in table below

D_j	$[0, 1 - \frac{x_4}{2}]$	$[1 - \frac{x_4}{2}, 1 - \frac{x_4}{2} + \frac{x_3}{2}]$	$[1 - \frac{x_4}{2} + \frac{x_3}{2}, 1]$
I_4	0	$(x_2 + \frac{x_4}{2} - 1)^2$	$-\frac{x_3^2}{4} + x_3(x_2 - 1 + \frac{x_4}{2})$

Adding all pieces together we obtain the following result

$$I = I_1 + I_2 + I_3 + I_4 \in \mathcal{A} \Leftrightarrow \begin{aligned} &x_2 > 1 + \frac{x_3 - x_4}{2} \quad \text{and} \quad I = x_3(x_4 + x_2 - 1) \\ &\text{or} \quad \frac{1 + x_3 - x_4}{2} < x_2 \leq \frac{1}{2} \quad \text{and} \quad I = x_3(\frac{x_4}{2} + x_2 - \frac{1}{2}). \end{aligned}$$

Proof of Step 2: Using Lemma 3.3 and Lemma 3.2 for $k = 3$ we calculate all the h_j^1 's in the regions $T_1 \cap \{x_3 \leq x_4\} \cap \{x_3 + x_4 \leq 1\}$ and $T_2 \cap \{x_3 \leq x_4\} \cap \{x_3 + x_4 \leq 1\}$. The obtained values are presented in the next two tables below.

j	(0, 0, 0)	(0, 0, 1)
$h_j^1(x)$	$(1 - x_2)(1 - x_4)$	$(1 - x_2)(x_4 - x_3)$
j	(1, 0, 0)	(1, 0, 1)
$h_j^1(x)$	$(1 - x_4)(x_2 - x_3)$	$x_2(x_4 - x_3) + x_3(1 - x_4)$
j	(0, 1, 0)	(0, 1, 1)
$h_j^1(x)$	0	$x_3(1 - x_2)$
j	(1, 1, 0)	(1, 1, 1)
$h_j^1(x)$	$x_3(1 - x_4)$	$x_3(x_4 + x_2 - 1)$

$h_j^1(x_2, x_3, x_4)$ in the set T_1

j	(0, 0, 0)	(0, 0, 1)
$h_j^1(x)$	$(1 - x_4)(1 - x_2 - \frac{x_3}{2})$	$x_4(1 - \frac{x_3}{2} - x_2) + x_3(x_2 - \frac{1}{2})$
j	(1, 0, 0)	(1, 0, 1)
$h_j^1(x)$	$(1 - x_4)(x_2 - \frac{x_3}{2})$	$x_2(x_4 - x_3) + \frac{x_3}{2}(1 - x_4)$
j	(0, 1, 0)	(0, 1, 1)
$h_j^1(x)$	$\frac{x_3}{2}(1 - x_4)$	$\frac{1}{2}x_3x_4 + x_3(\frac{1}{2} - x_2)$
j	(1, 1, 0)	(1, 1, 1)
$h_j^1(x)$	$\frac{x_3}{2}(1 - x_4)$	$\frac{1}{2}x_3x_4 - x_3(\frac{1}{2} - x_2)$

$h_j^1(x_2, x_3, x_4)$ in the set T_2

Now we use Lemma 3.1. Since the isometry $B_1(x_2, x_3, x_4) = (x_2, 1 - x_4, 1 - x_3)$ transforms the region $\{x_3 \leq x_4\}$ into itself and the region $\{x_3 + x_4 < 1\}$ into $\{x_3 + x_4 > 1\}$, we find the whole region where $h_{111}^1 \in \mathcal{A}$ in the set $\{x_3 \leq x_4\}$. It is easy to see (Remark 3.3) that this is the set $T_1 \cup T_2$. Moreover, using the identity $h_j(x) = h_{B_1j}(B_1x)$ we obtain the

formulas for all h_j^1 in $T_1 \cup T_2$. Those formulas are the same as presented in the two tables above. Now having described the whole region where $h_{111}^1 \in \mathcal{A}$ in the set $\{x_3 \leq x_4\}$ we use the symmetry with respect to the point $(1/2, 1/2, 1/2)$: $S = B_1 \circ B_2$, and according to the formulae $h_j^1(x) = h_{Sj}^1(Sx)$ we find all functions h_j^1 in sets T_3 and T_4 . Obviously, T_1, T_2, T_3 and T_4 are the only regions where we have $h_j \in \mathcal{A}$ for all j 's.

j	$(0, 0, 0)$	$(0, 0, 1)$
$h_j^1(x)$	$(1 - x_3)(1 - x_2 - x_4)$	$(1 - x_3)x_4$
j	$(1, 0, 0)$	$(1, 0, 1)$
$h_j^1(x)$	$(1 - x_3)x_2$	0
j	$(0, 1, 0)$	$(0, 1, 1)$
$h_j^1(x)$	$(1 - x_2)(x_3 - x_4) + x_4(1 - x_3)$	$x_4(x_3 - x_2)$
j	$(1, 1, 0)$	$(1, 1, 1)$
$h_j^1(x)$	$x_2(x_3 - x_4)$	x_2x_4

$h_j^1(x_2, x_3, x_4)$ in the set T_3

j	$(0, 0, 0)$	$(0, 0, 1)$
$h_j^1(x)$	$(1 - x_3)(1 - x_2 - \frac{x_4}{2})$	$\frac{1}{2}x_4(1 - x_3)$
j	$(1, 0, 0)$	$(1, 0, 1)$
$h_j^1(x)$	$(1 - x_3)(x_2 - \frac{x_4}{2})$	$\frac{1}{2}x_4(1 - x_3)$
j	$(0, 1, 0)$	$(0, 1, 1)$
$h_j^1(x)$	$(1 - x_2)(x_3 - x_4) + \frac{1}{2}x_4(1 - x_3)$	$x_4(\frac{1}{2} - x_2 + \frac{x_3}{2})$
j	$(1, 1, 0)$	$(1, 1, 1)$
$h_j^1(x)$	$(x_2 - \frac{1}{2})(x_3 - x_4) + \frac{1}{2}x_3(1 - x_4)$	$x_4(x_2 - \frac{1}{2} + \frac{x_3}{2})$

$h_j^1(x_2, x_3, x_4)$ in the set T_4

□

Now we will calculate $h_j^2(x_1, x_3, x_4)$. We have

Lemma 3.7 *The mapping $x \mapsto h_j^2(x)$ belongs to the space \mathcal{A} for every j if and only if $x \in \cup_{i=1}^4 \hat{S}(T_i)$ where T_i are the tetrahedrons from Lemma 3.6 and \hat{S} denotes the symmetry with respect to the plane $\{x_4 = \frac{1}{2}\}$. Moreover, we have*

$$h_{111}^2(x) = \begin{cases} x_3x_4 & \text{if } x \in \hat{S}(T_1) = \{x_1 > \frac{x_3+x_4+1}{2}\} \\ \frac{1}{2}x_3x_4 & \text{if } x \in \hat{S}(T_2) = \{\frac{1}{2} > x_1 > \frac{x_3+x_4}{2}\} \\ x_1(x_3 + x_4 - 1) & \text{if } x \in \hat{S}(T_3) = \{x_1 < \frac{x_3+x_4-1}{2}\} \\ (x_1 - \frac{1}{2})(x_3 + x_4 - 1) + \frac{1}{2}x_3x_4 & \text{if } x \in \hat{S}(T_4) = \{\frac{1}{2} < x_1 < \frac{x_3+x_4}{2}\} \end{cases}$$

where $x = (x_1, x_3, x_4)$.

Proof. This follows immediately from Lemmas 3.1 and 3.6. □

3.5 The computation of h_{1111}

For simplicity let us introduce the following definition.

Definition 3.1 We will say that the set D is \mathcal{A} -regular if there is decomposition $D = \cup_{i=1}^r D_i$ for some $r \in \mathbb{N}$, where for every $i \in \{1, \dots, r\}$ the set D_i is connected and the mapping $D_i \ni x \mapsto h_j(x)$ is represented by an element of \mathcal{A} for every $j \in \{0, 1\}^4$. Sets D_i will be called regular components of D .

We will say that D is maximal \mathcal{A} -regular subset of Ω if $D \subset \Omega$, D is \mathcal{A} -regular, and an arbitrary \mathcal{A} -regular subset of Ω is contained in D .

Now we are in the position to formulate the following result.

Lemma 3.8 *The maximal \mathcal{A} -regular subset of $Q = [0, 1]^4$ equals $R = \cup_{i=1}^8 R_i$, where simplexes R_i are regular components of R , R_1 is the principal simplex given by (2.9) R_2, \dots, R_8 are obtained by the condition $R_j = E_j(R_1)$ where E_j 's are isometries given by $E_1 = id$, $E_2 = A_1$, $E_3 = A_2$, $E_4 = A_2 \circ A_1$, $E_5 = (A_1 \circ A_2)^2$, $E_6 = (A_1 \circ A_2)^2 \circ A_1$, $E_7 = A_1 \circ A_2 \circ A_1$, $E_8 = A_1 \circ A_2$, and A_1 and A_2 are the same as in Lemma 3.1. The functions $h_j(x)$ on the set R_1 is described in the table below.*

$(0, 0, 0, 0)$	$(1, 0, 0, 0)$	$(0, 1, 0, 0)$	$(0, 0, 1, 0)$
$1 - (x_1 + x_2 + x_3 + x_4) + x_1x_3 + x_1x_4 + x_2x_3 + 1/2x_3x_4$	$x_1 - x_1x_3 - x_1x_4 + \frac{1}{2}x_3x_4$	$x_2(1 - x_3)$	$x_3 - x_1x_3 - x_2x_3 - \frac{1}{2}x_3x_4 + x_1x_2$
$(0, 0, 0, 1)$	$(1, 1, 0, 0)$	$(1, 0, 1, 0)$	$(1, 0, 0, 1)$
$x_4 - x_1x_4 - \frac{1}{2}x_3x_4$	0	$x_1x_3 - x_1x_2 - \frac{1}{2}x_3x_4 + x_2x_4$	$x_1x_4 - \frac{1}{2}x_3x_4$
$(0, 1, 1, 0)$	$(0, 1, 0, 1)$	$(0, 0, 1, 1)$	$(1, 1, 1, 0)$
$x_2x_3 - x_1x_2$	0	$\frac{1}{2}x_3x_4$	$x_1x_2 - x_2x_4$
$(0, 1, 1, 1)$	$(1, 0, 1, 1)$	$(1, 1, 0, 1)$	$(1, 1, 1, 1)$
0	$\frac{1}{2}x_3x_4 - x_2x_4$	0	x_2x_4

Functions h_j defined on R_1

The functions $h_j(x)$ for $x \in R_i$ and $i \in \{2, \dots, 8\}$ are obtained from those on R_1 according to the role given in Lemma 3.1: $h_j(x) = h_{E_i j}(E_i x)$.

Proof. The proof follows by steps: 1) We show that the maximal \mathcal{A} -regular subset in

$$\Omega_1 = \{(x_1, x_2, x_3, x_4) : x_1 + x_2 < 1, x_2 \leq x_1, x_3 + x_4 < 1\}$$

is contained in

$$R_1 = \{x \in Q : x_1 + x_2 < 1, x_3 + x_4 < 1, x_3 > x_1 + x_2, x_1 \leq 1/2, 2x_1 > x_3 + x_4\}.$$

2) R_1 is the maximal \mathcal{A} -regular subset of Ω_1 , and for $x \in R_1$ we have

$$h_{1111}(x) = x_2x_4.$$

- 3) We compute h_j 's for $x \in R_1$.
4) We complete the proof of the lemma.

Proof of Step 1. Let us denote the maximal \mathcal{A} -regular set under consideration by P_1 . According to Lemma 3.4 we have

$$P_1 \subseteq \Omega_2 = \Omega_1 \cap \{x \in \Omega : x_3 > x_1 + x_2\}.$$

Then by Lemma 3.5

$$P_1 \subseteq \Omega_3 = \Omega_2 \cap \{x \in Q : x_1 > x_2 + x_4\}.$$

An easy computation shows that if $x \in P_1$ we have $1 > x_3 > x_1 + x_2 > 2x_2 + x_4$, which implies $x_2 \leq 1/2$, and $x_4 \leq x_3$. Now we apply Lemma 3.6. As $x_2 \leq 1/2$, we have $P_1 \subseteq \Omega_3 \cap (T_2 \cup T_3)$ (we identify T_i 's with its embeddings in $Q \subseteq \mathbf{R}^4$). But $\Omega_3 \cap T_2 = \emptyset$, because $T_2 \subseteq \{x_4 \geq x_3\}$ (see Figure 3). Hence

$$P_1 \subseteq \Omega_4 = \Omega_3 \cap \{x \in Q : 2x_2 < x_3 - x_4\}.$$

Then we use Lemma 3.7, which implies that

$$P_1 \subseteq \Omega_4 \cap (\hat{S}(T_1) \cup \hat{S}(T_2) \cup \hat{S}(T_3) \cup \hat{S}(T_4)).$$

Now it suffices to verify that

$$\Omega_4 \cap \hat{S}(T_i) = \emptyset, \text{ for } i = 1, 3, 4. \quad (3.18)$$

The property (3.18) for $i = 1$ follows from the sequence of inequalities: $2x_1 > x_3 + x_4 + 1 > x_1 + x_2 + x_4 + 1$, which implies the impossible one: $x_1 > x_2 + x_4 + 1 > 1$; (3.18) for $i = 3$ and $i = 4$ contradicts the fact that $x_3 + x_4 < 1$ on P_1 . Summing up all conditions describing P_1 we get

$$\begin{aligned} P_1 &\subseteq R_1 = \{x \in Q : x_1 + x_2 < 1, x_2 \leq x_1, x_3 + x_4 < 1, \\ &x_4 \leq x_3, x_3 > x_1 + x_2, x_1 > x_2 + x_4, x_4 \leq 1/2, \\ &x_2 \leq 1/2, 2x_2 < x_3 - x_4, x_1 \leq 1/2, 2x_1 > x_3 + x_4\}. \end{aligned}$$

Now it suffices to eliminate those conditions in the description of P_1 , which are implied by the other's. To do this let us denote inequalities defying P_1 by 1)–11) respectively. Then it suffices to note that 6) implies 2), 5) and 6) imply 4) and 9), 6) and 10) imply 7) and 8), 5) and 11) imply 6). Since R_1 is the nonempty intersection of 5 halfspaces, it is the simplex in \mathbf{R}^4 .

Proof of Step 2. According to Lemma 3.2, Corollary 3.1 and Lemma 3.3 we have $D \ni x \mapsto h_j(x) \in \mathcal{A}$ for every j if and only if $D \ni x \mapsto h_{111}^i(x) \in \mathcal{A}$ for $i = \{1, \dots, 4\}$ and $D \ni x \mapsto h_{1111}(x) \in \mathcal{A}$. We have shown in the proof of Step 1 that the mapping $R_1 \ni x \mapsto h_{111}^i(x)$ is represented by elements of \mathcal{A} . Hence it suffices to show that $h_{1111}(x) \in \mathcal{A}$ for $x \in R_1$. We will show it by direct computation of $h_{1111}(x)$. For simplicity we put $I = h_{(1,1,1,1)}(x)$. Then an easy computation shows that for $x \in R_1$ we have $I = \int_0^1 \int_0^1 g(\tau_1, \tau_2) d\tau_1 d\tau_2$, where $g(\tau) = f_0(\tau)f_1(\tau)f_2(\tau)f_3(\tau)$, and for $\tau = (\tau_1, \tau_2)$

$$\begin{aligned} f_0(\tau) &= \chi_{\tau_1 \in [0, x_1], \tau_2 \in [0, x_2]}, \\ f_1(\tau) &= \chi_{(\tau_1 + \tau_2) < x_3}, \\ f_2(\tau) &= \chi_{[\tau_1 - \tau_2]_1 \in [0, x_4]}, \\ f_3(\tau) &= \chi_{\tau_1 > \tau_2}. \end{aligned}$$

(as for $\tau_1 \leq \tau_2$ and $x_4 \leq 1/2$ the function f_2 vanishes). Note that $f_0 f_3 \neq 0$ only if $\tau_2 < x_1$, while $f_1 \neq 0$ if $\tau_2 < x_3$ and $\tau_1 < x_3 - \tau_2$, so $f_1 f_3 \neq 0$ only if $\tau_2 < x_3 - \tau_2$, which is equivalent to: $\tau_2 < x_3/2$. Hence

$$I = \int_0^{\min(x_1, x_2, x_3/2)} \int_{\tau_2}^{h(\tau_2)} d\tau_1 d\tau_2, \quad \text{where } h(\tau) = \min(x_1, x_3 - \tau, x_4 + \tau).$$

But for $x \in R_1$ we have $\min\{x_1, x_2, x_3/2\} = x_2$, and $\min\{x_1, x_3 - \tau, x_4 + \tau\} = x_4 + \tau$ if $\tau < x_2$. Thus $I = x_2 x_4$.

Proof of Step 3. Lemmas: 3.4, 3.5, 3.6, 3.7 give for $x \in R_1$: $h_{111}^1(x) = x_2 x_4$, $h_{111}^2(x) = 1/2 x_3 x_4$, $h_{111}^3(x) = x_2 x_4$, $h_{111}^4(x) = x_1 x_2$, $h_{1111}(x) = x_2 x_4$. Thus we can use Remark 3.1 and calculate all the remaining coefficients h_j on R_1 . Note that according to our notation we have $A^0 = 1$, $A_j^1 = x_j$, $A_{j_1 j_2}^2 = x_{j_1} x_{j_2}$, $A_{123}^3 = h_{111}^4$, $A_{124}^3 = h_{111}^3$, $A_{134}^3 = h_{111}^2$, $A_{234}^3 = h_{111}^1$, $A_{1234}^4 = h_{1111}$.

Proof of Step 4. Consider the following sets

$$\begin{aligned} C_1 &= \{x_1 + x_2 < 1, x_2 \leq x_1, x_3 + x_4 < 1\} \\ C_2 &= \{x_1 + x_2 < 1, x_2 \leq x_1, x_3 + x_4 > 1\}. \end{aligned}$$

Since the isometry A_1 in Lemma 3.1 transforms C_1 onto C_2 , it follows from Lemma 3.1 and previous steps that $R_1 \cup (R_2 = A_1(R_1))$ is the maximal \mathcal{A} -regular subset of

$$\{x_1 + x_2 < 1, x_2 \leq x_1\}.$$

Now, if D is the maximal regular subset of $\{x_1 + x_2 < 1, x_2 \leq x_1\}$, then $A_2(D)$ is the maximal \mathcal{A} -regular subset of $\{x_1 + x_2 < 1, x_2 \geq x_1\}$. In particular $R_1 \cup R_2 \cup (R_3 = A_2(R_1)) \cup (R_4 = A_2(R_2))$ is the maximal \mathcal{A} -regular subset of $\{x_1 + x_2 < 1\}$. Finally, it suffices to use the isometry $A = A_1 \circ A_2 \circ A_1 \circ A_2$, which transforms the region $\{x_1 + x_2 < 1\}$ onto $\{x_1 + x_2 > 1\}$. Then we obtain the maximal \mathcal{A} -regular subset of $Q: \cup_{i=1}^8 R_i$ where $R_{4+i} = A(R_i)$, where $i = 1, \dots, 4$. The coefficients on every simplex R_i for $i = 2, \dots, 8$ are calculated according to the role $h_j(x) = h_{A_j}(Ax)$ in Lemma 3.1 from those given on R_1 , where we take $A = E_2, E_3, \dots, E_8$ for calculations on R_2, R_3, \dots, R_7 and R_8 respectively. \square

Remark 3.4 It is easy to check that the set of coefficients $\{h_j(x)\}_j$ for all simplex R_i where $i = 1, \dots, 8$ is different for different i .

4 The main result

We start with the following lemma.

Lemma 4.1 *Let us define the following operator*

$$C(\mathbf{R}^4) \ni f \mapsto \sum_{j \in \{0,1\}^4} \bar{h}_j(x) f(j) \in C(\mathbf{R}^4), \quad (4.19)$$

where $\bar{h}_j = h_j|_{R_1}$, \bar{h}_j 's are described by Lemma 3.8, and R_1 is the principal simplex given by (2.9). Then the operator described above is the projection operator onto the space \mathcal{A} , and agrees with the operator Pf described by (2.8).

Proof. The fact that the range of the operator given by (4.19) is contained in the space \mathcal{A} follows directly from Lemma 3.8. Now it suffices to describe the operator given by (4.19) in the basis of space \mathcal{A} . \square

Remark 4.1 Note that for $j = (1, 1, 1, 1)$ we have $\bar{h}_{1111}(1, 1, 1, 1) = 1$ but not all other coefficients $\bar{h}_j(1, 1, 1, 1)$ are equal to zero. In particular $Pf(1, 1, 1, 1) \neq f(1, 1, 1, 1)$.

The following lemma will be crucial to obtain our main results.

Lemma 4.2 *Let G be the special group of invariances, R_1 be the principal simplex (see Section 2.3), and assume that the simplex $R \subseteq \mathbf{R}^4$ is G -similar to R_1 , that is $R = g(R_1)$ for some $g \in G$. Suppose that f defines a special functional which is lower semicontinuous with respect to the weak- $*$ convergence of the u_i 's in L^∞ . Then for every $x \in R$ we have*

$$f(x) \leq (g^*Pf)(x),$$

where $g^*Pf(x) = \sum_{j \in \{0,1\}^4} \bar{h}_j(g^{-1}x)f(gj)$, P is the same as in (2.8).

Proof. The proof follows by steps: 1) we obtain the result for $R = R_1$ and 2) we complete the proof of the theorem.

Proof of Step 1): Let $x \in R_1$ and $u(\tau, x)$ be the same as in Lemma 2.5. According to Lemma 2.6 we have

$$f(x) \leq \int_{[0,1]^2} f(u(\tau, x))d\tau.$$

Using Lemmas 2.5 and 3.8 we see that the right hand side equals $\sum_{j \in \{0,1\}^4} \bar{h}_j(x)f(j)$.

Proof of Step 2): Let $f_g(y) = f(gy)$. By Lemma 2.4 and by Step 1 applied to f_g we get $f_g(y) \leq Pf_g(y)$ for every $y \in R_1$. Now it suffices to substitute $x = gy \in R$. \square

Remark 4.2 Let $\tilde{G} \subseteq G$ be the subgroup of G of isometries of $[0, 1]^4$ introduced in Section 2.3. It consists of 64 elements, let us denote them by g_1, g_2, \dots, g_{64} , where $g_k = E_k$, for $k = 1, \dots, 8$, E_k 's are the same as in Lemma 3.8, and g_9, \dots, g_{64} are the remaining elements of \tilde{G} written in an arbitrary order. Define $R_i = g_i(R_1)$ for $i = 1, \dots, 64$. According to Lemma 4.2 the inequality $f(x) \leq (g_i^*Pf)(x)$ holds for every $x \in R_i$ and for every $i \in \{1, \dots, 64\}$. Note that Lemma 3.8 guarantees such inequality only for $i = 1, \dots, 8$, as coefficients $h_j(x)$ belong to the space \mathcal{A} only for x being in simplex R_1, \dots, R_8 . In particular the application of Lemma 2.4 gives stronger result than that which follows directly from Lemmas 3.8 and Lemma 2.5.

As a corollary we obtain the following result.

Corollary 4.1 *If we assume additionally in Lemma 4.2 that $f \in \mathcal{C}$ then we have $f(x) = (Pf)(x)$ for every $x \in \mathbf{R}^4$, and consequently the space \mathcal{C} agrees with \mathcal{A} .*

First proof of Corollary 4.1. At first we note that for arbitrary open and bounded set $U \subseteq \mathbf{R}^4$ we will find $g \in G$ such that $U \subset g(R_1)$. Thus according to Lemma 4.2 applied to f and $-f$ we see that f agrees in U with a certain polynomial belonging to the space \mathcal{A} . As two polynomials which agree on an open subset of \mathbf{R}^n must be the same, we see that $f \in \mathcal{A}$. This implies $\mathcal{C} \subseteq \mathcal{A}$. The reverse inclusion is obtained by direct computation. \square

The above result could also be obtained as the consequence of the following known result due to Murat and Tartar (see e.g. [16, Theorem 3.3 on page 27], [53, Theorem 5.1], [78, Theorem18]).

Theorem 4.1 Assume that $f : \mathbf{R}^m \rightarrow \mathbf{R}$ defines a weakly continuous functional I_f given by (1.4). Then f is a polynomial of degree $\min\{n, m\}$, moreover, the following property is satisfied:

$$\left\{ \begin{array}{l} \text{Given arbitrary } r \in \mathbf{N} \text{ and } (\xi_1, \lambda_1), \dots, (\xi_r, \lambda_r) \in V \text{ such that} \\ \text{rank}\{\xi_1, \dots, \xi_r\} \leq r - 1, \text{ we have for arbitrary } s \in \mathbf{R}^m \\ f^{(r)}(s)(\lambda_1, \dots, \lambda_r) = 0. \end{array} \right. \quad (4.20)$$

Second proof of Corollary 4.1. It reminds to note that in our case we have $\Lambda = \cup_{i=1}^4 \text{span}\{e_i\} \subseteq \mathbf{R}^4$, and $V \subseteq \mathbf{R}^2 \times \mathbf{R}^4$. Hence $\mathcal{C} \subseteq \mathcal{A}$, while the reverse inclusion is obtained by direct computation. \square

Let us introduce the following definition.

Definition 4.1 Let $P : C(\mathbf{R}^4) \rightarrow \mathcal{A}$ be the projection operator defined by (2.8), R_1 be the principal simplex given by (2.9), and G be the special group of invariances. We will say that $f \in C(\mathbf{R}^4)$ is sub \mathcal{A} -function with respect to the projection operator P on G -similar sets to R_1 (in short $\text{sub}(\mathcal{A}, P, R_1, G)$) if the inequality

$$f(x) \leq (g^* P f)(x)$$

holds for every $g \in G$ and every $x \in g(R_1)$ where $(g^* P f)(x) = \sum_{j \in \{0,1\}^4} h_j(g^{-1}x) f(gj)$.

Remark 4.3 Note that the above definition is related to the classical definition of convexity. Let us take instead of \mathcal{A} the space of affine functions on \mathbf{R} and denote it by A . Take $R = [0, 1]$, and let P be the projection operator from $C(\mathbf{R})$ to A , given by

$$P f(x) = (1 - x)f(0) + x f(1) := h_1(x)f(0) + h_2(x)f(1),$$

and consider the group G_1 of transformations of \mathbf{R} generated by translations $x \mapsto a + x$ and dilations $x \mapsto bx$. Then one could define in analogous way the class of sub A -functions with respect to the projection operator P on G_1 -similar sets to R by saying that $f \in \text{sub}(A, P, R, G_1)$ if the inequality

$$f(x) \leq (h_1(g^{-1}(x))f(g(0)) + h_2(g^{-1}(x))f(g(1)))$$

holds for every $g \in G_1$ and $x \in g(R)$ where $g \in G_1$. As an arbitrary $g \in G_1$ is of the form: $g(t) = a + bt$, the above inequality is equivalent to the inequality:

$$f((1 - t)a + t(a + b)) \leq (1 - t)f(a) + t f(a + b),$$

which holds for arbitrary $a, b \in \mathbf{R}$, and $t \in [0, 1]$. But this is nothing else than convexity.

Now we are in the position to state our main result.

Theorem 4.2 *Assume that $f \in C(\mathbf{R}^m)$ defines a weakly lower semicontinuous functional I_f given by (1.4), V and Λ are the same as in Theorem 1.1. Then f satisfies the following condition:*

Let $(\xi_1, \lambda_1), (\xi_2, \lambda_2), (\xi_3, \lambda_3), (\xi_4, \lambda_4) \in V$ are such that λ_i 's are linearly independent, ξ_1 and ξ_2 are linearly independent, $\xi_3 = \xi_1 + \xi_2$ and $\xi_4 = \xi_1 - \xi_2$. Then for arbitrary $A \in \mathbf{R}^m$ the mapping

$$\mathbf{R}^4 \ni x = (x_1, x_2, x_3, x_4) \mapsto \tilde{f}(x) = f\left(A + \sum_{i=1}^4 x_i \lambda_i\right)$$

is the sub(\mathcal{A}, P, R_1, G) function.

In the case when f defines a weakly continuous functional I_f the described mapping above belongs to the space \mathcal{A} .

Proof. It suffices to note that for arbitrary $u_i \in L^\infty(\mathbf{R})$ ($i = 1, 2, 3, 4$) the set $\{A + \sum_{i=1}^4 u_i \langle x, \xi_i \rangle \lambda_i\}$ is the subset of $\text{Ker} P$. In particular \tilde{f} defines special functional which is lower semicontinuous with respect to the weak-* convergence of the u_i 's in L^∞ . \square

Remark 4.4 Obviously, the second statement can be deduced directly from Theorem 4.1. A similar result for \tilde{f} depending on three variables only is given in Theorem 3.2 in [34].

Our next results apply directly to the variational case. As the consequence of Theorem 4.2 we obtain the following theorem.

Theorem 4.3 *Assume that $\Omega \subseteq \mathbf{R}^n$ is a bounded domain, and $f \in C(\mathbf{R}^{n \times m})$ defines a sequentially weakly-* lower semicontinuous functional on the Sobolev space $W^{1,\infty}(\Omega, \mathbf{R}^m)$: $\tilde{I}_f(w) = \int_\Omega f(\nabla w(x)) dx$, where $w : \Omega \rightarrow \mathbf{R}^m$. Let us take $A \in \mathbf{R}^{n \times m}$, two independent vectors: $\xi_1, \xi_2 \in \mathbf{R}^n$, and $a_1, a_2, a_3, a_4 \in \mathbf{R}^m$ and define $\lambda_i = \xi_i \otimes a_i$ for $i \in \{1, 2\}$ and $\lambda_3 = (\xi_1 + \xi_2) \otimes a_3$, $\lambda_4 = (\xi_1 - \xi_2) \otimes a_4$, where $\xi \otimes a = (\xi_i a_j)_{i=1,\dots,n, j=1,\dots,m} \in \mathbf{R}^{n \times m}$. Assume that all λ_i 's are linearly independent. Then the mapping*

$$\mathbf{R}^4 \ni x = (x_1, \dots, x_4) \mapsto \tilde{f}(x) = f\left(A + \sum_{i=1}^4 x_i \lambda_i\right)$$

is the sub(\mathcal{A}, P, R_1, G) function.

Proof. In this case the manifold V consists of pairs $(\xi, \xi \otimes a)$ where $\xi \in \mathbf{R}^n \setminus \{0\}$ and $a \in \mathbf{R}^m$. Hence the result follows directly from Theorem 4.2. \square

Remark 4.5 Continuity assumptions can be violated in all our statements. It suffices to assume that f is Borel measurable and bounded. The argument is the following. Assume at first that $f : \mathbf{R} \rightarrow \mathbf{R}$ is Borel measurable, locally bounded and defines the lower semicontinuous functional $I_f(u) = \int_{[0,1]} f(u(x)) dx$ defined on the class of measurable and

bounded functions $u : \mathbf{R} \rightarrow \mathbf{R}$. Let $u(x)$ be the periodic extension of $\chi_{[0,t]}A + \chi_{[t,1]}B$ where $t \in (0, 1)$ and $A, B \in \mathbf{R}$ are taken arbitrary, and let $u^\nu(x) = u(\nu x)$ where $\nu \in \mathbf{N}$. If we apply lower semicontinuity assumption to this sequence we see (the argument follows directly from Riemann–Lebesgue’s Theorem, see e.g. Lemma 1.2 in [16]) that f must be convex, so also continuous. This argument can be used in the proof of Theorem 1.1, where one restricts the general functional I_f to the subspace in $\text{Ker}P$ given by $\{A + v(\langle x, \xi \rangle)\lambda, v \in L^\infty\}$ where $(\xi, \lambda) \in V$. In particular, every Borel measurable and bounded integrand which defines a lower semicontinuous functional given by (1.4) must be Λ -convex, so also continuous in all Λ -directions. In particular if Λ spans all of \mathbf{R}^m then f must be continuous. In our arguments we study the behaviour of f along subspaces spanned by Λ only (starting from an arbitrary point in \mathbf{R}^m), so continuity assumption is satisfied there. This remark was presented to us by Pietro Celada.

5 Final conclusions and remarks

There are a series of questions and remarks naturally arising after following these results. We state them below with the hope that the proposed research program will contribute to further development of the Calculus of Variations and Compensation Compactness Theory.

Three dimensional conditions.

Remark 5.1 Let D be an arbitrary tetrahedron in \mathbf{R}^3 with three edges parallel to the axes. It is obtained by translation and dilation I of the cutted corner of the standard cube $Q = [0, 1]^3$ (see picture below).




Figure 5: $D_{(0,0,1)}^+$ and $D_{(0,0,1)}^-$

Let us denote $D = I(D_{(0,0,1)}^+)$, where $D_{(0,0,1)}^+ = \text{conv}\{(0, 0, 1), (0, 1, 1), (1, 0, 1), (0, 0, 0)\}$. Let $P_D^+ f$ be the only polynomial from 7 dimensional space of polynomials $\text{span}\{1, x_i, x_i x_j : i, j \in \{1, 2, 3\}, i \neq j\}$, which equals with f in 7 corners of the cube $\tilde{Q} = I(Q)$, all except $I(1, 1, 0)$.

Let $f \in C(\mathbf{R}^3)$ and consider the following functional

$$I_f(u) = \int_{[0,1]^2} f(u_1(x_1), u_2(x_2), u_3(x_1 + x_2)) dx_1 dx_2, \quad (5.21)$$

where $u_i \in L^\infty(\mathbf{R})$.

It was proved in [34] that if the functional I_f is lower semicontinuous then the following inequality

$$f(x) \leq P_D^+ f(x) \quad (5.22)$$

holds for every $x \in D$.

Obviously, such an inequality implies analogous inequalities for integrands defining lower semicontinuous functionals having the general form (1.4), satisfied on three dimensional simplexes, see Theorem 3.2 in [34]. We do not know if it is possible to obtain three dimensional conditions analogous to (5.22) as the direct consequence of our new four-dimensional conditions obtained in Lemma 4.2 and Theorem 4.2.

Remark 5.2 Three dimensional conditions (5.22) obtained in Theorem 3.1 in [34] were satisfied on all cutted corners of the standard cube $[0, 1]^3$ (as for example $D_{(0,0,1)}^+$ under notation of Remark 5.1). Our new four dimensional conditions among this which use corners of the standard cube $[0, 1]^4$ obtained in Lemma 4.2 are satisfied on R_1 and his \tilde{G} similar cousins. None of this simplexes is the cutted corner of the standard cube in \mathbf{R}^4 .

Remark 5.3 Assume that $f \in C(\mathbf{R}^3)$ defines a lower semicontinuous functional of the general form (1.3). The calculations in Section 3.4 suggest that there is a whole family of new conditions of the form $f(x) \leq Pf(x)$, where P is a certain projection operator onto the space of sufficiently good polynomials, which are satisfied on three dimensional simplexes. We can easily deduce such one's with similar but different statements to (5.22) on simplexes obtained by transformations of T_2 in Lemma 3.7. This shows that even three dimensional conditions still require further investigations.

Perspectives for further geometric conditions

Remark 5.4 We think that if one considers functionals of the form

$$I_f(u) = \int_{\Omega} f(u_1(A_1x), \dots, u_m(A_mx)) dx \quad (5.23)$$

where $u_i : \mathbf{R} \rightarrow \mathbf{R}$ and $A_i : \mathbf{R}^n \rightarrow \mathbf{R}$ are some linear operators then additional geometric conditions will appear. These conditions will be satisfied on simplexes with the specially prescribed geometry and probably will have a form of inequality between f on the simplex and some projection operator onto the space of weakly continuous functionals given by (5.23) like in Lemma 4.2. The relation between the structure of the A_i 's, the geometry of these simplexes and the projection operators in this relation requires systematic study. Such issue should lead to the generalization of Theorems 4.2, 4.3 and Theorem 3.2 in [34].

New conditions and CW-structures.

Remark 5.5 Note that the new geometric conditions have CW–structural nature: they are given inside simplexes, and on their lower order skeleton’s. For example Λ –convexity condition is the condition given on the subset of the 1–skeleton of the simplex.

Some other estimates.

Remark 5.6 In [34] also other conditions were obtained. It was shown in Theorem 3.2 there that under notations of Remark 5.1 we have for every $x \in D^- = I(D_{(0,0,1)}^-)$

$$f(x) \leq (P_D^- f)(x), \quad (5.24)$$

where $D_{(0,0,1)}^- = \text{conv}\{(1/2, 1/2, 1/2), (1/2, 0, 1/2), (0, 1/2, 1/2), (1/2, 1/2, 1)\}$, and $P_D^- f(x)$ is the disturbance of $P_D^+ f(x)$: $P_D^- f(x) = P_D^+ f(x) + Rf(x)$, where $Rf(x) = \text{dist}(x, D)^2 \cdot \Delta^3 f$, $\text{dist}(x, D)$ is the distance function, and $\Delta^3 f$ is certain combination of values of f in corners of \tilde{Q} (the exact definition is given in [34]). In particular $P_D^- f(x)$ does not define weakly continuous special functional of the form (5.21), as it was the case of $P_D^+ f(x)$. It would be interesting to know if one may obtain (5.24) as the consequence of inequalities like (5.22).

One can also obtain variants of (5.24) on four dimensional simplexes (they cannot be relatives to the nobel R_1), with regular term disturbed by square roots of distances from walls of those simplexes. We have seen this in our calculations when writing this paper, but because of technical difficulties we did not take care about their exact form.

Geometric conditions and elliptic systems.

Remark 5.7 Let us look at the inequality in Definition 4.1: $f(x) \leq (g^* P f)(x)$ more closely. Its right hand side, $G = g^* P f$, is the solution to the elliptic system: $\frac{\partial^2}{\partial x_i^2} G = 0$ for $i \in \{1, 2, 3, 4\}$, $\frac{\partial}{\partial x_i} \frac{\partial}{\partial x_j} \frac{\partial}{\partial x_k} G = 0$ for different i, j, k , and G agrees with f on certain given set (related to $g(R_1)$, it consists of corners of $g(Q)$ where Q is the standard cube). Inequality holds in the certain set of dimension 4. We think we have to do with a kind of subsolutions to elliptic systems. Recall that if one deals with a single elliptic equation then the subsolution has the property that if it coincides with the solution to this equation on the boundary of the sufficiently regular set Ω then inside Ω it is less or equal than this solution (see e.g. [26], [28]). Here we deal with similar property. We refer e.g. to [2], [3], [12], [13], [27], [45], [50], and their references for various maximum principles for solutions of linear and nonlinear elliptic systems.

Numerical treatment and Morrey’s conjecture.

Remark 5.8 In the literature there are several of examples of functions which are known to be rank–one convex but it is not confirmed whether they are quasiconvex (see e.g. [1, 10, 18, 19, 29, 72]). Now one could check at least numerically if those functions satisfy the new three and four dimensional geometric conditions obtained in [34] and in Theorem 4.2. The numerical verification is in progress now (see [35]).

Šverák’s example and locality.

Remark 5.9 It was shown in [34] that the rank–one convex function which is not quasi-convex constructed by Šverák in his famous paper [74] does not satisfy the new geometric conditions on three dimensional simplex (see Remark 4.1 in [34]). It is possible to verify that also four dimensional conditions of our Theorem 4.3 cannot be satisfied by this function either. The sketch of the proof is the following. If we relax the dependence of f on x_4 in Lemma 4.2 we arrive at the same inequality as (5.22) but on a subsimplex of D . Now we can use the same arguments as in Remark 4.1 in [34] and show that Šverák’s function (after the slight modification to make it strongly rank–one convex) does not satisfy this three dimensional condition. Thus it cannot satisfy the four dimensional conditions either. This shows also that our four dimensional conditions cannot be local: there exists strongly rank–one convex function which is not quasiconvex and does not satisfy this condition. But as was shown by Kristensen (see [40] for details) every strongly rank–one convex function agrees with quasiconvex functions on balls covering $\mathbf{R}^{n \times m}$. As each of them satisfies four dimensional conditions, it follows that our conditions restricted to functions defined on $n \times m$ matrices where $n \geq 2$ and $m \geq 3$ cannot be local (see also Remark 4.5 in [34]).

New conditions and null–lagrangians.

Remark 5.10 It is known that if $f : \mathbf{R}^{n \times m} \rightarrow \mathbf{R}$ defines weakly continuous variational functional given by (1.1) then f is a null–lagrangian, that is f belongs to the linear space spanned by $\{1, \lambda_{ij}, m_{I,J}\}$ where $m_{I,J}$ are minors of matrices in $\mathbf{R}^{n \times m}$ (see e.g. [5], [6], [11], [16], [17], [24], [78], [79]). In particular if $n = m = 2$ the space of null–lagrangians is 6 dimensional. As the determinant function is affine along every rank–one direction and is the polynomial of degree 2, we see that under the notation of Theorem 4.3 the function \tilde{f} belongs to \mathcal{A} for arbitrary four rank–one matrices $\lambda_1, \dots, \lambda_4$ like in Theorem 4.3. Since \mathcal{A} is 11 dimensional space, we do not think we can expect that for an arbitrary quasiconvex function \tilde{f} the mapping $P\tilde{f}$ will be a null–lagrangian. It would be interesting to understand the relations between projections $P\tilde{f}$ of quasiconvex integrands and null–lagrangians. We refer e.g. to [30], [31], [32], [46], [54], [77] and their references for selected results on null–lagrangians.

Are our conditions sufficient for lower semicontinuity?

Remark 5.11 Let us consider the variational case. If we deal with gradients of scalar functions of two variables, that is $m = 1$ under the notation of (1.1) then quasiconvexity condition is equivalent to the usual convexity. It would be interesting to know what is the answer on two related questions:

1. Is it true that if we deal with second gradients of scalar functions of two variables, that is $\nabla u \in \mathbf{R}^{2 \times 2}$ and ∇u is symmetric, then rank–one tetrahedral convexity condition (the condition on three dimensional simplexes) of Theorem 4.1 in [34] is equivalent to quasiconvexity?
 2. Is it true that in the case $n = m = 2$ (gradients are 2×2 matrices) our new geometric necessary conditions on four dimensional simplexes are also sufficient for quasiconvexity?
- For answers on both questions it would be helpful to know if our new convexity conditions on three and for dimensional simplexes from Theorem 3.1 in [34] and Lemma 4.2 are also sufficient for lower semicontinuity of functionals $I_f(u) = \int_{[0,1]^2} f(u_1(\tau_1), u_2(\tau_2), u_3(\tau_1 + \tau_2)) d\tau$ and $I_f(u) = \int_{[0,1]^2} f(u_1(\tau_1), u_2(\tau_2), u_3(\tau_1 + \tau_2), u_4(\tau_1 - \tau_2)) d\tau$ respectively.

Hulls, envelops and other remarks.

Remark 5.12 Let $f \geq 0$ and f^{qc} and f^{rc} be the quasiconvex and rank–one convex envelop of f , that is the largest quasiconvex and rank–one convex minorant of f respectively. By the Fundamental Relaxation Theorem we know that the minimum value of I_f given by (4.3) is the same as the minimum of $I_{f^{qc}}$ in the class $\{u \in W^{1,\infty}(\Omega, \mathbf{R}^m) : u \equiv Fx \text{ on } \partial\Omega\}$ (see e.g. [16], [70]). It is not known how to compute in general the quasiconvex envelop of f (see e.g. [37], [38], [69], [44] for some of the very few results), but as quasiconvexity implies rank–one convexity we have $f^{qc} \leq f^{rc} \leq f$. Hence the minimum of I_f is the same as the minimum of $I_{f^{rc}}$ and it becomes important to be able to compute the rank–one convex envelop of f when one looks for minimum of I_f . Now we can introduce envelopes of f which will be related to the new geometric conditions. Let us denote them by f^x . As these conditions imply rank–one convexity condition we will have $f^{qc} \leq f^x \leq f^{rc} \leq f$, so f^x will be closer to the quasiconvex envelope of f than f^{rc} . Perhaps the new envelopes will be more helpful for numerical computations, for example it will be easier to find the minimizing sequence for I_{f^x} than for $I_{f^{rc}}$. We refer e. g. to [9], [20], [21], [37], [38], [47],[66] and their references for the approach related to rank–one convex and quasiconvex envelopes and their computation.

Remark 5.13 Let $K \subseteq \mathbf{R}^{n \times m}$ be the closed subset. In variational problems of martensitic phase transitions and material microstructures, in the general theory of Partial Differential Inclusions solved by the method of Convex Integration due to Gromow and its applications to construct wild solutions of nonlinear elliptic and parabolic systems (see e.g. [7], [8], [15], [76], [58], [59], [60], [61], [62], [36]) one introduces various semiconvex hulls of sets. Those hulls are defined as quasiconvex, rank–one convex and lamination convex hulls respectively. The first two hulls are defined as cosets of quasiconvex and rank–one convex functions respectively (see e.g. [84], [76] for details). The lamination convex hull is defined as the smallest set \hat{K} with the following property: $K \subseteq \hat{K}$ and for all $A, B \in \hat{K}$ which satisfy $\text{rank}(A - B) = 1$ where the interval $[A, B]$ is also contained in \hat{K} . Now one can define hulls described with the help of new geometric conditions. For example instead of adding the rank–one intervals in the construction of rank–one convex hulls one can add simplexes instead and obtain richer sets. This may be more efficient in particular in the construction of solutions of Partial Differential Inclusions by methods of Convex Integration, where one solves at first the inclusion: $Du \in \hat{K}$ on the “convexified” set (using the nonstandard convexification) and uses the in–approximation method to reduce the set \hat{K} to the original one. In particular if the “convexified” set \hat{K} is bigger, it is easier to solve Partial Differential Inclusion. We also refer e. g. to [39], [42], [84], [23], [22] and their references for the related works.

Remark 5.14 Some other questions related to three dimensional conditions for integrands defying lower semicontinuous functionals were stated in [34]. One can forward them and use four dimensional conditions instead.

References

- [1] J.J. ALIBERT, B. DACOROGNA, An example of a quasiconvex function that is not polyconvex in two dimensions two. *Arch. Ration. Mech. Anal.* **117** (1992), 155–166.

- [2] S. AGMON, Maximum theorems for solutions of higher order elliptic equations. *Bull. Am. Math. Soc.* **66** (1960), 77–80.
- [3] S. AGMON, L. NIRENBERG, M. H. PROTTER, A maximum principle for a class of hyperbolic equations and applications to equations of mixed elliptic–hyperbolic type. *Commun. Pure Appl. Math.* **6** (1953), 455–470.
- [4] K. ASTALA, Analytic aspects of quasiconformality. *Doc. Math. J. DMV, Extra Volume ICM 1998 Vol. II*, 617–626.
- [5] J.M. BALL, Convexity conditions and existence theorems in nonlinear elasticity. *Arch. Ration. Mech. Anal.* **63** (1977), 337–403.
- [6] J.M. BALL, Constitutive inequalities and existence theorems in nonlinear elastostatics. *Nonlin. Anal. Mech., Heriot–Watt Symp. Vol. I (R. Knops, Ed.)*, 1977, Pitman, London, 187–241.
- [7] J. M. BALL, R. D. JAMES, Fine phase mixtures as minimizers of energy. *Arch. Ration. Mech. Anal.* **100** (1987), 13–52.
- [8] J. M. BALL, R. D. JAMES, Proposed experimental tests of a theory of fine microstructure and the two–well problem. *Philos. Trans. R. Soc. Lond.*, **338**(A) (1992), 389–450.
- [9] J. M. BALL, B. KIRCHHEIM, J. KRISTENSEN, Regularity of quasiconvex envelopes. *Calc. Var. Partial Differ. Equ.* **11** (2000), 333–359.
- [10] J.M. BALL, F. MURAT, Remarks on rank-one convexity and quasiconvexity. *Ordinary and Partial Differential Equations, eds. B.D. Sleeman and R.J. Jarvis, Vol. III, Pitman Research Notes in Mathematics Series 254*, 1991, Longman, New York, 25–37.
- [11] J.M. BALL, J. C. CURRIE, P. J. OLVER, Null Lagrangians, weak continuity, and variational problems of arbitrary order. *J. Funct. Anal.* **41** (1981), 135–174.
- [12] A. V. BITSADZE, A system of nonlinear partial differential equations. *Differ. Uravn.* **15** (1979), no. 7, 1267–1270 (in Russian).
- [13] A. CANFORA, Teorema del massimo modulo e teorema di esistenza per il problema di Dirichlet relativo ai sistemi fortemente ellittici. *Ric. Mat.* **15** (1966), 249–294.
- [14] E. CASADIO–TARABUSI, An algebraic characterization of quasiconvex functions. *Ric. Mat.* **42** (1993), 11–24.
- [15] M. CHIPOT, D. KINDERLEHRER, Equilibrium configurations of crystals. *Arch. Ration. Mech. Anal.* **103** (1988), 237–277.
- [16] B. DACOROGNA, *Weak continuity and weak lower semicontinuity for nonlinear functionals*. Lecture Notes in Mathematics 922, Berlin-Heidelberg-New York, Springer, 1982.
- [17] B. DACOROGNA, *Direct methods in the calculus of variations*. Springer, Berlin 1989.
- [18] B. DACOROGNA, J. DOUCHET, W. GANGBO, J.RAPPAZ, Some examples of rank–one convex functions in dimension two. *Proc. R. Soc. Edinb.* **114** (1990), 135–150.

- [19] B. DACOROGNA, J.-P. HAEBERLY, Some numerical methods for the study of the convexity notions arising in the calculus of variations. *M²AN, Mathematical Modelling and Numerical Analysis* **32** (1998), 153–175.
- [20] G. DOLZMANN, Numerical computation of rank–one convex envelopes. *SIAM J. Numer. Anal.* **36**(5), (1999) 1621–1635.
- [21] G. DOLZMANN, *Variational methods for crystalline microstructure–analysis and computation*. Lecture Notes in Mathematics, 1803, Springer–Verlag, Berlin, 2003.
- [22] G. DOLZMANN, B. KIRCHHEIM, S. MÜLLER, V. ŠVERÁK, The two–well problem in three dimensions. *Calc. Var. Partial Differ. Equ.* **10** (2000), 21–40.
- [23] G. DOLZMANN, B. KIRCHHEIM, J. KRISTENSEN, Conditions for equality of hulls in the calculus of variations. *Arch. Ration. Mech. Anal.* **154** (2000), 93–100.
- [24] D. G. B. EDELEN, The null set of the Euler–Lagrange operator. *Arch. Ration. Mech. Anal.* **11** (1962), 117–121.
- [25] I. FONSECA, S. MÜLLER, A –quasiconvexity, lower semicontinuity and Young measures. *SIAM J. Math. Anal.* **30** (1999), 1355–1390.
- [26] L. E. FRAENKEL, *An introduction to maximum principles and symmetry in elliptic problems*. Cambridge University Press, Cambridge, 2000.
- [27] M. GIAQUINTA, E. GIUSTI, Quasi–minima, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **1**(2) (1984), 79–107.
- [28] D. GILBARG, N.S. TRUDINGER, *Elliptic partial differential equations of second order*. Springer–Verlag, Berlin–Heidelberg–New York 1977.
- [29] T. IWANIEC, Nonlinear Cauchy–Riemann operators in \mathbf{R}^n , *Trans. Am. Math. Soc.* **354** (2002), 1961–1995.
- [30] T. IWANIEC, Integrability theory of the Jacobians. *Lipshitz Lectures, preprint Univ. Bonn Sonderforschungsbereich* **256** (1995).
- [31] T. IWANIEC, Nonlinear differential forms. *Series in Lectures at the International School in Jyväskylä*, published by *Math. Inst. Univ. Jyväskylä* (1998), 1–207.
- [32] T. IWANIEC, A. LUTOBORSKI, Integral estimates for null–lagrangians. *Arch. Ration. Mech. Anal.* **125** (1993), 25–79.
- [33] A. KALAMAJSKA, On Λ –convexity conditions in the theory of lower semicontinuous functionals, to appear in *J. Convex Anal.*
- [34] A. KALAMAJSKA, On new geometric conditions for some weakly lower semicontinuous functionals with applications to the rank–one conjecture of Morrey, preprint 636 of the Institute of Mathematics of the Polish Academy of Sciences, to appear in *Proc. R. Soc. Edinb.*
- [35] A. KALAMAJSKA, K. MOSZYŃSKI, R. STEFANIUK, From rank–one convexity to tetrahedral convexity. Theoretical and numerical approach. In preparation.

- [36] B. KIRCHHEIM, S. MÜLLER, V. ŠVERÁK, Studing nonlinear pde by geometry in matrix space. in: *Geometric Analysis and Nonlinear Differential Equations* (H. Karcher, S. Hildebrandt eds.), Springer 2003, 347–395.
- [37] V. KOHN, G. STRANG, Optimal design and relaxation of variational problems I. *Commun. Pure Appl. Math.* **39** (1986), 113–137.
- [38] V. KOHN, G. STRANG, Optimal design and relaxation of variational problems II. *Commun. Pure Appl. Math.* **39** (1986), 139–182.
- [39] J. KOLÁŘ, Non-compact lamination convex hulls. *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **20**(3) (2003), 391–403.
- [40] J. KRISTENSEN, On the non-locality of quasiconvexity. *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **16** (1999), 1–13.
- [41] M. KRUŽÍK, On the composition of quasiconvex functions and the transposition. *J. Convex Anal.* **6** (1999), 207–213.
- [42] M. KRUŽÍK, Bauer’s maximum principle and hulls of sets. *Calc. Var. Partial Differ. Equ.* **11** (2000), 321–332.
- [43] S. LANG, *Algebra*. Addison–Wesley Publishing Company, New York 1965.
- [44] H. LE DRET, A. RAOULT, The quasiconvex envelope of the Saint–Venant–Kirchhoff stored energy function. *Proc. R. Soc. Edinb.* **125** (1995), 1179–1192.
- [45] F. LEONETTI, Maximum principle for vector-valued minimizers of some integral functionals. *Boll. Unione Mat. Ital.* **7**(5–A) (1991), 51–56.
- [46] P. L. LIONS, Jacobians and Hardy spaces. *Ric. Mat. Suppl.* **40** (1991), 255–260.
- [47] M. LUSKIN, On the computation of crystalline microstructure. *Acta Numerica* **5** (1996), 191–257.
- [48] J. J. MANFREDI, Weakly monotone functions. *J. Geom. Anal.* **4**(2) (1994), 393–402.
- [49] P. MARCELLINI, Quasiconvex quadratic forms in two dimensions. *Appl. Math. Optimization* **11** (1984), 183–189.
- [50] M. MIRANDA, Maximum principles and minimal surfaces. *Ann. Sc. Norm. Super. Pisa, Cl. Sci.* (4) Vol. XXV (1997), 667–681.
- [51] C. B. MORREY, Quasi-convexity and the lower semicontinuity of multiple integrals. *Pac. J. Math.* **2** (1952), 25–53.
- [52] C. B. MORREY, *Multiple integrals in the calculus of variations*. Springer–Verlag, Berlin–Heidelberg–New York, 1966.
- [53] F. MURAT, A survey on compensated compactness. *Contributions to modern calculus of variations* (L. Cesari ed.), *Pitman Research Notes in Mathematics Series* **148**, Longman, Harlow, 1987, 145–183.

- [54] S. MÜLLER, A surprising higher integrability property of mappings with positive determinant. *Bull. Am. Math. Soc.* **21** (1989), 245–248.
- [55] S. MÜLLER, *Variational models for microstructure and phase transitions, Collection: Calculus of variations and geometric evolution problems (Cetraro 1996), 85–210, Lecture Notes in Math., 1713, Springer, Berlin 1999.*
- [56] S. MÜLLER, Rank–one convexity implies quasiconvexity on diagonal matrices. *Int. Math. Res. Not.*, **20**, (1999), 1087–1095.
- [57] S. MÜLLER, Quasiconvexity is not invariant under transposition. *Proc. R. Soc. Edinb.*, **130**, (2000), 389–395.
- [58] S. MÜLLER, V. ŠVERÁK, Attainment results for the two–well problem by convex integration. *Geom. Anal. and the Calc. Variations, (J. Jost, ed.), International Press, 1996, 239–251.*
- [59] S. MÜLLER, V. ŠVERÁK, Unexpected solutions of first and second order partial differential equations. *Doc. Math. J. DMV, Special volume Proc. ICM 1998, Vol. II, 691–702.*
- [60] S. MÜLLER, V. ŠVERÁK, Convex integration for Lipschitz mappings and counterexamples to regularity, preprint MPI–MIS 26/1999.
- [61] S. MÜLLER, V. ŠVERÁK, Convex integration with constraints and applications to phase transitions and partial differential equations. *J. Eur. Math. Soc. (JEMS)* **1/4** (1999), 393–422.
- [62] S. MÜLLER, M. O. RIEGER, V. ŠVERÁK, Parabolic systems with nowhere smooth solutions, preprint, <http://www.math.cmu.edu/nwOz/publications/02-CNA-014/014abs/>
- [63] G. P. PARRY, On the planar rank–one convexity condition. *Proc. R. Soc. Edinb.* **125A** (1995), 247–264.
- [64] P. PEDREGAL, *Parametrized measures and variational principles. Birkhäuser, 1997.*
- [65] P. PEDREGAL, Weak continuity and weak lower semicontinuity for some compensation operators. *Proc. R. Soc. Edinb.* **113(A)** (1989), 267–279.
- [66] P. PEDREGAL, Laminates and microstructure. *Eur. J. Appl. Math.* **4** (1993), 121–149.
- [67] P. PEDREGAL, Some remarks on quasiconvexity and rank–one convexity. *Proc. R. Soc. Edinb.* **126(A)** (1996), 1055–1065.
- [68] P. PEDREGAL, V. ŠVERÁK, A note on quasiconvexity and rank–one convexity for 2×2 Matrices. *J. Convex Anal.* **5** (1998), 107–117.
- [69] A. C. PIPKIN, Elastic materials with two preferred states. *Q. J. Mech. Appl. Math.* **44** (1991), 1–15.
- [70] T. ROUBÍČEK, *Relaxation in optimization theory and variational calculus. Berlin, W. de Gruyter, 1997.*

- [71] J. SIVALOGANATHAN, Implications of rank one convexity. *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **5,2** (1988), 99–118.
- [72] V. ŠVERÁK, Examples of rank–one convex functions. *Proc. R. Soc. Edinb.* **114A** (1990), 237–242.
- [73] V. ŠVERÁK, Quasiconvex functions with subquadratic growth. *Proc. R. Soc. Lond.* **433A**, (1991), 723–725.
- [74] V. ŠVERÁK, Rank–one convexity does not imply quasiconvexity. *Proc. R. Soc. Edinb.* **120** (1992) 185–189.
- [75] V. ŠVERÁK, Lower semicontinuity of variational integrals and compensated compactness. *Proceedings of the Internaional Congress of Mathematicians, Zürich, Switzerland 1994, Birkhäuser Verlag, Basel, Switzerland 1995*, 1153–1158.
- [76] V. ŠVERÁK, On the problem of two wells. In: *Microstructures and phase transitions, IMA Vol. Appl. Math.* **54**, (D. Kinderlehrer, R. D. James, M. Luskin and J. Ericksen, eds., Springer, 1993, 183–189.
- [77] J. ROBBIN, R. C. ROGERS, B. TEMPLE, On weak continuity and Hodge decomposition. *Trans. Am. Math. Soc.* **303** (1987), 609–618.
- [78] L. TARTAR, Compensated compactness and applications to partial differential equations. *Nonlinear Analysis and Mechanics: Heriot–Watt Symp., vol. IV (ed. R. Knops), Pitman Res. Notes Math.* **39**, 1979, 136–212.
- [79] L. TARTAR, The compensated compactness method applied to systems of conservation laws. *Systems of Nonlinear Partial Differential Eq. (ed. J.M. Ball) , Reidel, (1983), 263–285.*
- [80] L. TARTAR, Some remarks on separately convex functions, Microstructure and Phase Transitions. *IMA Vol. Math. Appl.* **54** (eds. D. Kinderlehrer, R.D. James, M. Luskin and J.L. Ericksen), Springer 1993, 191–204.
- [81] B. YAN, On rank–one convex and polyconvex conformal energy functions with slow growth. *Proc. R. Soc. Edinb.* **127** (1997) 651–663.
- [82] K. W. ZHANG, A construction of quasiconvex functions with linear growth at infinity. *Ann. Sc. Norm. Super. Pisa, Cl. Sci. Ser. IV* **XIX** (1992), 313–326.
- [83] K. W. ZHANG, Biting theorems for Jacobians and their applications. *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **7** (1990), 345–365 .
- [84] K. W. ZHANG, On various semiconvex hulls in the calculus of variations. *Calc. Var. Partial Differ. Equ.* **6** (1998), 143–160.