

EXISTENTIAL \emptyset -DEFINABILITY OF HENSELIAN VALUATION RINGS

ARNO FEHM

Abstract. In [1], Anscombe and Koenigsmann give an existential \emptyset -definition of the ring of formal power series $F[[t]]$ in its quotient field in the case where F is finite. We extend their method in several directions to give general definability results for henselian valued fields with finite or pseudo-algebraically closed residue fields.

§1. Introduction. The question of first order definability of valuation rings in their quotient fields has a long history. Given a valued field K , one is interested in whether there exists a first order formula φ in the language $\mathcal{L} = \{+, -, \cdot, 0, 1\}$ of rings such that the set $\varphi(K)$ defined by φ in K is precisely the valuation ring, and what complexity such formula must have.

Many results of this kind are known for *henselian* valued fields, like fields of formal power series $K = F((t))$ over a field F , and their valuation ring $F[[t]]$. In this setting, a definition going back to Julia Robinson gives an existential definition of the valuation ring using the parameter t . Later, Ax [2] gave a definition of the valuation ring, which uses no parameters, but is not existential.

Recently, Anscombe and Koenigsmann [1] succeeded to give an existential and parameter-free definition of $F[[t]]$ in $F((t))$ in the special case where $F = \mathbb{F}_q$ is a finite field. Their proof uses the fact that \mathbb{F}_q can be defined in $\mathbb{F}_q((t))$ by the quantifier-free formula $x^q - x = 0$. In particular, their result does not apply to any infinite field F , and their formula depends heavily on q .

In this note we simplify and extend their method. As a first application we get the following general definability result for henselian valued fields with finite or pseudo-algebraically closed residue fields (Theorem 2.6 and Theorem 3.5), which generalizes [1, Theorem 1.1] on $\mathbb{F}_q((t))$ and [5, Theorem 6] on finite extensions of \mathbb{Q}_p :

THEOREM 1.1. *Let K be a henselian valued field with valuation ring \mathcal{O} and residue field F . If F is finite or pseudo-algebraically closed and the algebraic part of F is not algebraically closed, then there exists an \exists - \emptyset -definition of \mathcal{O} in K .*

As a further application, in Section 4, we find definitions of the valuation ring which are uniform for large (infinite) families of finite residue fields, like the following one for finite prime fields (Theorem 4.3):

Received August 22, 2013.

Key words and phrases. Henselian valuation, existential definability, finite fields, pseudo-algebraically closed fields.

© 2015, Association for Symbolic Logic
0022-4812/15/8001-0015
DOI:10.1017/jsl.2014.13

THEOREM 1.2. *For every $\varepsilon > 0$ there exists an \exists - \emptyset -formula φ and a set P of prime numbers of Dirichlet density at least $1 - \varepsilon$ such that for any henselian valued field K with valuation ring \mathcal{O} and residue field F with $|F| \in P$, the formula φ defines \mathcal{O} in K .*

In particular, this applies to power series fields $\mathbb{F}_p((t))$ and p -adic fields \mathbb{Q}_p . Theorem 1.2 is in a sense optimal, see the discussion at the end of this note.

§2. Defining subsets of the valuation ring. Let K be a henselian valued field with valuation ring $\mathcal{O} \subseteq K$, maximal ideal $\mathfrak{m} \subseteq \mathcal{O}$ and residue field $F = \mathcal{O}/\mathfrak{m}$. For $a \in \mathcal{O}$ we let $\bar{a} = a + \mathfrak{m} \in F$ be its residue class and write $\bar{f} \in F[X]$ for the reduction of a polynomial $f \in \mathcal{O}[X]$.

We start by simplifying the key lemma of [1], thereby generalizing it to arbitrary henselian valuations. This proof follows Helbig [10]. Here, and in what follows, by $f(K)^{-1}$ we mean the set $\{f(x)^{-1} : x \in K\}$ and implicitly claim that $f(x) \neq 0$ for all $x \in K$.

LEMMA 2.1. *Let $f \in \mathcal{O}[X]$ be a monic polynomial such that \bar{f} has no zero in F , and let $a \in K$. Let $U_{f,a} := f(K)^{-1} - f(a)^{-1}$. Then the following holds:*

- a) $f(K)^{-1} \subseteq \mathcal{O}$,
- b) $U_{f,a} \subseteq \mathcal{O}$,
- c) *If in addition $a \in \mathcal{O}$ and $f'(a) \notin \mathfrak{m}$, then $\mathfrak{m} \subseteq U_{f,a}$.*

PROOF. a) We have that $f(K) \cap \mathfrak{m} = \emptyset$: If $x \in K$ with $f(x) \in \mathfrak{m}$, then $x \in \mathcal{O}$ since \mathcal{O} is integrally closed and f is monic, and hence $\bar{f}(\bar{x}) = 0$, contradicting the assumption that \bar{f} has no zero in F . Therefore, $f(K)^{-1} \subseteq (K \setminus \mathfrak{m})^{-1} = \mathcal{O}$.

b) From a) we get that $f(K)^{-1} \subseteq \mathcal{O}$, and in particular $f(a)^{-1} \in \mathcal{O}$. Thus, $U_{f,a} \subseteq \mathcal{O}$.

c) Now assume that $a \in \mathcal{O}$ and $f'(a) \notin \mathfrak{m}$. Let $x \in \mathfrak{m}$. Since $a \in \mathcal{O}$ we have $f(a) \in \mathcal{O}$, hence $f(a) \in \mathcal{O}^\times$. Define $g(X) = f(X) - (f(a) + x) \in \mathcal{O}[X]$. Then $g(a) = -x \in \mathfrak{m}$ and $g'(a) = f'(a) \notin \mathfrak{m}$, so by the assumption that \mathcal{O} is henselian there exists $b \in \mathcal{O}$ with $g(b) = 0$, i.e. $f(a) + x = f(b)$. Hence, $f(a) + \mathfrak{m} \subseteq f(K)$. Since $f(a) \in \mathcal{O}^\times$ we get that $f(a)^{-1} + \mathfrak{m} = (f(a) + \mathfrak{m})^{-1} \subseteq f(K)^{-1}$, and therefore $\mathfrak{m} \subseteq U_{f,a}$. ⊖

We observe that one can get rid of the element a even if it is not in the (model theoretic) algebraic closure of the prime field:

LEMMA 2.2. *Let $f \in \mathcal{O}[X]$ be a monic polynomial such that \bar{f} has no zero in F , and $a \in \mathcal{O}$ such that $f'(a) \notin \mathfrak{m}$. Then $U := f(K)^{-1} - f(K)^{-1}$ satisfies $\mathfrak{m} \subseteq U \subseteq \mathcal{O}$.*

PROOF. By Lemma 2.1a, $f(K)^{-1} \subseteq \mathcal{O}$, hence $U \subseteq \mathcal{O}$. Since $a \in \mathcal{O}$ and $f'(a) \notin \mathfrak{m}$, Lemma 2.1c implies that $\mathfrak{m} \subseteq U_{f,a} \subseteq U$. ⊖

Clearly, U can be defined in K by the \exists -formula

$$\varphi_f(x) \equiv (\exists y, z, y_1, z_1)(x = y_1 - z_1 \wedge y_1 f(y) = 1 \wedge z_1 f(z) = 1).$$

Note that if $f \in \mathbb{Z}[X]$, then φ_f is an \exists - \emptyset -formula.

LEMMA 2.3. *If $U, T \subseteq \mathcal{O}$ are such that $\mathfrak{m} \subseteq U$ and T meets all residue classes (i.e. $\bar{T} = F$), then $\mathcal{O} = U + T$.*

PROOF. If for $x \in \mathcal{O}$ we let $t \in T$ with $\bar{t} = \bar{x}$, then $x = u + t$ with $u := x - t \in \mathfrak{m} \subseteq U$. ⊖

Thus, if φ defines U and ψ defines T , then

$$\eta(x) \equiv (\exists u, t)(x = u + t \wedge \varphi(u) \wedge \psi(t))$$

defines \mathcal{O} . Note that if φ and ψ are \exists - \emptyset -formulas, then so is η .

We now give a first generalization of [1, Theorem 1.1]. We denote by F_0 the prime field of F and by F_{alg} the algebraic closure of F_0 in F . By abuse of notation we will consider polynomials $f \in \mathbb{Z}[X]$ as elements of $\mathcal{O}[X]$ via the canonical homomorphism $\mathbb{Z} \rightarrow \mathcal{O}$.

LEMMA 2.4. *For every prime p and positive integer m , there exists $f \in \mathbb{F}_p[X]$ monic, separable, and irreducible of degree m with $f'(0) \neq 0$.*

PROOF. Let $q = p^m$. Since $\mathbb{F}_q/\mathbb{F}_p$ is Galois it has a normal basis, i.e. there exists $\alpha \in \mathbb{F}_q$ such that the conjugates of α form an \mathbb{F}_p -basis of \mathbb{F}_q . In particular, α has degree m and nonzero trace over \mathbb{F}_p . Let $f \in \mathbb{F}_p[X]$ be the minimal polynomial of α^{-1} . Then f is irreducible of degree m and $f'(0) = \pm \text{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(\alpha) / \text{N}_{\mathbb{F}_q/\mathbb{F}_p}(\alpha) \neq 0$. \dashv

LEMMA 2.5. *If F is finite, then there exist $f \in F_0[X]$ monic, separable, and irreducible which has no zero in F , and $a \in F$ with $f'(a) \neq 0$.*

PROOF. Identify $F_0 = \mathbb{F}_p$, let m be any positive integer that does not divide $[F : F_0]$, choose f of degree m as in Lemma 2.4, and let $a = 0$. \dashv

THEOREM 2.6. *Let K be a henselian valued field with valuation ring \mathcal{O} and residue field F . If F is finite, then there exists an \exists - \emptyset -definition of \mathcal{O} in K .*

PROOF. If $F = \mathbb{F}_q$, let $g = X^q - X \in \mathbb{Z}[X]$ and $\psi(x) \equiv (g(x) = 0)$. Since $\bar{g}' = -1$, the assumption that \mathcal{O} is henselian gives that $T := \psi(K) \subseteq \mathcal{O}$ is a set of representatives of F . In particular, it meets all residue classes. Choose $f \in F_0[X]$ as in Lemma 2.5 and let $\tilde{f} \in \mathbb{Z}[X]$ be a monic lift of f . Since there exists $a \in F$ with $f'(a) \neq 0$, a lift $\tilde{a} \in \mathcal{O}$ of a satisfies $\tilde{f}'(\tilde{a}) \notin \mathfrak{m}$. Let $\varphi \equiv \varphi_{\tilde{f}}$. By Lemma 2.2, $U := \varphi(K)$ satisfies $\mathfrak{m} \subseteq U \subseteq \mathcal{O}$. Therefore, Lemma 2.3 shows that $\eta(K) = \mathcal{O}$. \dashv

The special case of Theorem 2.6 where F is a finite field and $K = F((t))$ was proven by Anscombe and Koenigsmann in [1, Theorem 1.1]. The special case where K is a finite extension of \mathbb{Q}_p was proven by Cluckers, Derakhshan, Leenknegt, and Macintyre in [5, Theorem 6].

§3. Pseudo-algebraically closed residue fields. We now consider assumptions on the residue field F under which we can define a set T as in Lemma 2.3. For basics on pseudo-algebraically closed (PAC) fields we refer to [8, Chapter 11]. For $d \in \mathbb{N}$ we fix the constant $c(d) = (2d - 1)^4$.

LEMMA 3.1. *Let $f \in F[X]$ be nonconstant and square-free (over the algebraic closure). Then $F = f(F)f(F) \cup \{0\}$ if F is PAC or F is finite with $|F| > c(\deg(f))$.*

PROOF. Let $0 \neq c \in F$. One checks that the polynomial $f(X)f(Y) - c \in F[X, Y]$ is absolutely irreducible, cf. [9, Proposition 1.1]. Thus, if F is PAC we can conclude that there exist $x, y \in F$ with $f(x)f(y) - c = 0$, i.e. $c \in f(F)f(F)$. If F is finite with $|F| > c(\deg(f))$ we come to the same conclusion by applying the Hasse-Weil bound, cf. [8, Corollary 5.4.2]. \dashv

LEMMA 3.2. *Let $f \in \mathcal{O}[X]$ be monic such that \bar{f} is square-free and has no zero in F . Then $T := f(K)^{-1}f(K)^{-1} \cup \{0\} \subseteq \mathcal{O}$. If in addition F is PAC or finite with $|F| > c(\deg(f))$, then T meets all residue classes.*

PROOF. By Lemma 2.1a, $f(K)^{-1} \subseteq \mathcal{O}$, hence $T \subseteq \mathcal{O}$. If F is PAC or finite with $|F| > c(\deg(f))$, then, since $F^\times \subseteq \bar{f}(F)\bar{f}(F)$ by Lemma 3.1, also

$$F^\times \subseteq (\bar{f}(F)\bar{f}(F))^{-1} \subseteq (\overline{f(\mathcal{O})} \cdot \overline{f(\mathcal{O})})^{-1} \subseteq \overline{f(K)^{-1}f(K)^{-1}},$$

hence T satisfies $F = \bar{T}$. ⊣

Clearly, the set T can be defined in K by the \exists -formula

$$\psi_f(x) \equiv (\exists y, z, y_1, z_1)(x = 0 \vee (x = y_1z_1 \wedge y_1f(y) = 1 \wedge z_1f(z) = 1)).$$

Let

$$\eta_f(x) \equiv (\exists u, t)(x = u + t \wedge \varphi_f(u) \wedge \psi_f(t)).$$

PROPOSITION 3.3. *Let $f \in \mathcal{O}[X]$ be monic such that \bar{f} is square-free and has no zero in F . Then $\eta_f(K) \subseteq \mathcal{O}$. If in addition there exists $a \in \mathcal{O}$ such that $f'(a) \notin \mathfrak{m}$ and F is PAC or finite with $|F| > c(\deg(f))$, then $\eta_f(K) = \mathcal{O}$.*

PROOF. Let $U = \varphi_f(K)$, so $U \subseteq \mathcal{O}$ by Lemma 2.1. By Lemma 3.2, $T := \psi_f(K) \subseteq \mathcal{O}$, so $\eta_f(K) = U + T \subseteq \mathcal{O}$. If in addition there exists $a \in \mathcal{O}$ such that $f'(a) \notin \mathfrak{m}$ and F is PAC or finite with $|F| > c(\deg(f))$, then Lemma 2.2 gives that $\mathfrak{m} \subseteq U$, and Lemma 3.2 gives that T meets all residue classes, hence $\eta_f(K) = \mathcal{O}$ by Lemma 2.3. ⊣

LEMMA 3.4. *If F is infinite and F_{alg} is not algebraically closed, then there exist $f \in F_0[X]$ monic, separable, and irreducible which has no zero in F , and $a \in F$ with $f'(a) \neq 0$.*

PROOF. Since F_{alg} is not algebraically closed, there exists a monic irreducible $f \in F_0[X]$ which has no zero in F_{alg} , hence in F . Since F_0 is perfect, f is separable, hence $f' \neq 0$. Therefore, since F is infinite, there exists $a \in F$ with $f'(a) \neq 0$. ⊣

THEOREM 3.5. *Let K be a henselian valued field with valuation ring \mathcal{O} and residue field F . If F is pseudo-algebraically closed and F_{alg} is not algebraically closed, then there exists an \exists - \emptyset -definition of \mathcal{O} in K .*

PROOF. Choose $f \in F_0[X]$ as in Lemma 3.4 and let $\tilde{f} \in \mathbb{Z}[X]$ be a monic lift of f . Since there exists $a \in F$ with $f'(a) \neq 0$, a lift $\tilde{a} \in \mathcal{O}$ of a satisfies $\tilde{f}'(\tilde{a}) \notin \mathfrak{m}$. By Proposition 3.3, $\eta_{\tilde{f}}(K) = \mathcal{O}$. ⊣

COROLLARY 3.6. *Let K be a henselian valued field with valuation ring \mathcal{O} and residue field F . If F is pseudo-real closed and F_{alg} is neither real closed nor algebraically closed, then there exists an \exists - \emptyset -definition of \mathcal{O} in K .*

PROOF. Let $K' = K(\sqrt{-1})$. Then the residue field $F' = F(\sqrt{-1})$ of K' is PAC by [11], and $F'_{\text{alg}} = F_{\text{alg}}(\sqrt{-1})$ is not algebraically closed by the Artin-Schreier theorem. By Theorem 3.5 there exists an \exists - \emptyset -definition of the unique prolongation \mathcal{O}' of \mathcal{O} in K' . By interpreting K' in K we get an \exists - \emptyset -definition of $\mathcal{O} = \mathcal{O}' \cap K$ in K . ⊣

REMARK 3.7. Note that as soon as F is infinite we cannot hope to have an \exists - \emptyset -definition of a set of representatives $T \subseteq \mathcal{O}$ of F : For example, if $K = F((t))$, then F is never \exists - \emptyset -definable in K unless it is finite, cf. [7, Corollary 9]. This explains why we rather define a set $T \subseteq \mathcal{O}$ that meets all residue classes.

REMARK 3.8. We point out that the assumption that F_{alg} is not algebraically closed in Theorem 3.5 is indeed necessary. For example, let K be the field of generalized

power series $F((\mathbb{Q}))$ over a field F . If F_{alg} is algebraically closed, then so is $K' := F_{\text{alg}}((\mathbb{Q}))$, cf. [6, 18.4.3]. Therefore, K' is existentially closed in K . So, if φ is an \exists - \emptyset -definition of the valuation ring in K , then $\varphi(K') = \varphi(K) \cap K'$ is a nontrivial valuation ring, contradicting the fact that definable subsets of an algebraically closed field are finite or cofinite.

§4. Uniform definitions. We now deal with definitions which are uniform over certain families of finite residue fields. We start with an example in fixed residue characteristic p :

THEOREM 4.1. *Given a prime number p and a positive integer m there exists an \exists - \emptyset -formula φ such that $\varphi(K) = \mathcal{O}$ for all henselian valued fields K with valuation ring \mathcal{O} and residue field $F = \mathbb{F}_{p^n}$ with $m \nmid n$.*

PROOF. Assume that $F = \mathbb{F}_{p^n}$ with $m \nmid n$. Choose $f \in \mathbb{F}_p[X]$ irreducible of degree m as in Lemma 2.4. Then f has no zero in F and there exists $a \in F$ with $f'(a) \neq 0$. Let $\tilde{f} \in \mathbb{Z}[X]$ be a monic lift of f . By Proposition 3.3, $\eta_{\tilde{f}}(K) \subseteq \mathcal{O}$, and $\eta_{\tilde{f}}(K) = \mathcal{O}$ for $p^n > c(m)$. For $k \in \mathbb{N}$ with $m \nmid k$ let $\psi_k(x) \equiv (x^{p^k} - x = 0)$ and let

$$\eta_k(x) \equiv (\exists u, t)(x = u + t \wedge \varphi_{\tilde{f}}(u) \wedge \psi_k(t)).$$

As in the proof of Theorem 2.6 we see that $\eta_k(K) \subseteq \mathcal{O}$, and $\eta_k(K) = \mathcal{O}$ if $n = k$. Therefore, with $M = \{k \in \mathbb{N} : m \nmid k \text{ and } p^k \leq c(m)\}$,

$$\varphi(x) \equiv \eta_{\tilde{f}}(x) \vee \bigvee_{k \in M} \eta_k(x)$$

satisfies $\varphi(K) = \mathcal{O}$ for all n with $m \nmid n$. ◻

REMARK 4.2. The condition $m \nmid n$ in Theorem 4.1 is indeed necessary: If a \exists - \emptyset -formula φ defines $\mathbb{F}_{p^n}[[t]]$ in $\mathbb{F}_{p^n}((t))$ for all n in a set M , then there is some $m \in \mathbb{N}$ such that $m \nmid n$ for all $n \in M$: Otherwise, $\bigcup_{n \in M} \mathbb{F}_{p^n}$ would equal the algebraic closure of \mathbb{F}_p , so since every finite extension of $\mathbb{F}_{p^n}((t))$ is isomorphic to $\mathbb{F}_{p^{n'}}((t))$ for some $n|n'$, we would get a definition of a nontrivial valuation ring in the algebraic closure of $\mathbb{F}_p((t))$, which is impossible. For details the reader may consult [5, Theorem 4], where it is shown that no \exists - \emptyset -formula can define the valuation ring uniformly for all finite extensions of a fixed henselian valued field K .

We now turn to uniformity in p . Let \mathbb{P} denote the set of all odd prime numbers. For a subset $P \subseteq \mathbb{P}$, we denote by $\delta(P)$ the Dirichlet-density of P , if it exists. For a formula φ let

$$\mathbb{P}(\varphi) = \{p \in \mathbb{P} : \varphi(\mathbb{Q}_p) = \mathbb{Z}_p\}$$

and let $\mathbb{P}'(\varphi)$ be the set of $p \in \mathbb{P}$ such that $\varphi(K) = \mathcal{O}$ for all henselian valued fields K with valuation ring \mathcal{O} and residue field $F = \mathbb{F}_p$. We have that $\mathbb{P}'(\varphi) \subseteq \mathbb{P}(\varphi)$, and it is known that $\mathbb{P}(\varphi)$ has a Dirichlet-density for every formula φ , cf. [3, Theorem 16], [8, Theorem 20.9.3]. It is also known that $\mathbb{P}(\varphi)$ differs from $\{p \in \mathbb{P} : \varphi(\mathbb{F}_p((t))) = \mathbb{F}_p[[t]]\}$ only by a finite set, see [4, p. 606], so for all results concerning Dirichlet density we could as well use $\mathbb{F}_p((t))$ instead of \mathbb{Q}_p .

THEOREM 4.3. *For every $\varepsilon > 0$ there exists an \exists - \emptyset -formula φ such that $\delta(\mathbb{P}'(\varphi)) > 1 - \varepsilon$.*

PROOF. For $n \in \mathbb{N}$ let $f_n = X^2 - n \in \mathbb{Z}[X]$ and

$$P_n = \left\{ p \in \mathbb{P} : \left(\frac{n}{p} \right) = -1 \right\} = \{ p \in \mathbb{P} : \mathbb{F}_p \not\models (\exists y)(y^2 = n) \}.$$

Note that if K is henselian valued with residue field $F = \mathbb{F}_p$, $p > 2$, then $p \in P_n$ if and only if $K \not\models (\exists y)(y^2 = n)$. If $p \in P_n$ with $p > c(2)$, then $p \in \mathbb{P}'(\eta_{f_n})$ by Proposition 3.3. By the quadratic reciprocity law and Dirichlet's theorem, there exists $N \in \mathbb{N}$ such that for $P = \bigcup_{n=2}^N P_n$ we have $\delta(P) > 1 - \varepsilon$. Let

$$\varphi_n(x) \equiv (\exists y)(y^2 = n) \vee \eta_{f_n}(x)$$

and $\varphi(x) \equiv \bigwedge_{n=2}^N \varphi_n(x)$. Let $p \in \mathbb{P}$ which lies in the open interval $I := (c(2), \infty)$. If $p \in P_n$, then $\varphi_n(K) = \mathcal{O}$, otherwise $\varphi_n(K) = K$. Thus, $\varphi(K) = \bigcap_{n=2}^N \varphi_n(K) = \mathcal{O}$ if $p \in P$, and $\varphi(K) = K$ otherwise. So, if $p \in P$, then $p \in \mathbb{P}'(\varphi) \subseteq \mathbb{P}(\varphi)$, and if $p \notin P$, then $p \notin \mathbb{P}(\varphi)$. Thus, $\mathbb{P}'(\varphi) \cap I = \mathbb{P}(\varphi) \cap I = P \cap I$, and therefore $\delta(\mathbb{P}'(\varphi)) = \delta(\mathbb{P}(\varphi)) = \delta(P) > 1 - \varepsilon$. \dashv

On the other hand, it is well known that there is no such formula that works uniformly for almost all p :

PROPOSITION 4.4. *Let $P \subseteq \mathbb{P}$ be a cofinite set of prime numbers. Then there exists no \exists - \emptyset -formula φ such that $P \subseteq \mathbb{P}(\varphi)$.*

A proof of this can be found in [5, Theorem 5]. In fact, the proof given there shows the following stronger statement:

PROPOSITION 4.5. *Let P be a set of prime numbers with $\delta(P) = 1$. Then there exists no \exists - \emptyset -formula φ such that $P \subseteq \mathbb{P}(\varphi)$.*

This also explains that Theorem 4.3 cannot be strengthened to give a uniform \exists - \emptyset -definition for every set P with $\delta(P) < 1$:

PROPOSITION 4.6. *There exists a set P of prime numbers with $\delta(P) = 0$ for which there exists no \exists - \emptyset -formula φ such that $P \subseteq \mathbb{P}(\varphi)$.*

PROOF. List all \exists - \emptyset -formulas as $\varphi_1, \varphi_2, \dots$ and let $N = \{\ell_1, \ell_2, \dots\} \subseteq \mathbb{P}$ be any infinite set with $\delta(N) = 0$. Proposition 4.4 implies that for each i , $\mathbb{P}(\varphi_i)$ is not cofinite in \mathbb{P} . Therefore, we can choose some $p_i \in \mathbb{P}$ with $p_i > \ell_i$ and $p_i \notin \mathbb{P}(\varphi_i)$. Then $P = \{p_1, p_2, \dots\}$ has $\delta(P) \leq \delta(N) = 0$, but $P \not\subseteq \mathbb{P}(\varphi_i)$ for each i . \dashv

§5. Acknowledgements. The author would like to thank Will Anscombe, Patrick Helbig, Franziska Jahnke, and Jochen Koenigsmann for interesting and productive discussions on this subject, and Moshe Jarden for helpful comments on a previous version of the manuscript.

REFERENCES

[1] WILL ANSCOMBE and JOCHEN KOENIGSMANN, *An existential \emptyset -definition of $\mathbb{F}_q[[t]]$ in $\mathbb{F}_q((t))$* , this JOURNAL, vol. 79 (2014), no. 4, pp. 1336–1343.
 [2] JAMES AX, *On the undecidability of power series fields*. *Proceedings of the American Mathematical Society*, vol. 16 (1965), p. 846.
 [3] ———, *The elementary theory of finite fields*. *Annals of Mathematics*, vol. 88 (1968), no. 2, pp. 239–271.

- [4] JAMES AX and SIMON KOCHEN, *Diophantine problems over local fields I*. *American Journal of Mathematics*, vol. 87 (1965), no. 3, pp. 605–630.
- [5] RAF CLUCKERS, JAMSHID DERAKHSHAN, EVA LEENKNEGT, and ANGUS MACINTYRE, *Uniformly defining valuation rings in henselian valued fields with finite or pseudo-finite residue fields*. *Annals of Pure and Applied Logic*, vol. 164 (2013), pp. 1236–1246.
- [6] IDO EFRAI, *Valuations, Orderings, and Milnor K-Theory*, American Mathematical Society, Providence, 2006.
- [7] ARNO FEHM, *Subfields of ample fields. Rational maps and definability*. *Journal of Algebra*, vol. 323 (2010), pp. 1738–1744.
- [8] MICHAEL D. FRIED and MOSHE JARDEN, *Field Arithmetic*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics, vol. 11, Springer, Berlin, 2008.
- [9] B. HEINEMANN and A. PRESTEL, *Fields regularly closed with respect to finitely many valuations and orderings*. *Canadian Mathematical Society Conference Proceedings*, vol. 4 (1984), pp. 297–336.
- [10] PATRICK HELBIG, *Existentielle Definierbarkeit von Bewertungsringen*, Bachelor thesis, Konstanz, 2013.
- [11] ALEXANDER PRESTEL, *Pseudo real closed fields*, *Set Theory and Model Theory, Proceedings, Bonn 1979* (R. B. Jensen and A. Prestel, editors), Springer, 1981, pp. 127–156.

FACHBEREICH MATHEMATIK UND STATISTIK
UNIVERSITY OF KONSTANZ
78457 KONSTANZ, GERMANY
E-mail: arno.fehm@uni-konstanz.de