

Universität Konstanz, Fakultät für Biologie, Lehrstuhl für Phytopathologie, Konstanz, Germany

## Assessment of Sensitivities to Anilinopyrimidine- and Strobilurin-fungicides in Populations of the Apple Scab Fungus *Venturia inaequalis*

S. KUNZ, B. LUTZ, H. DEISING and K. MENDGEN

Authors' address: Universität Konstanz, Fakultät für Biologie, Lehrstuhl für Phytopathologie, Universitätsstr. 10, D-78434 Konstanz, Germany (correspondence to K. Mendgen)

With 5 figures

Received November 26, 1997; accepted December 1, 1997

### Abstract

The sensitivity of *Venturia inaequalis* populations to the anilinopyrimidine fungicides pyrimethanil and cyprodinil was analysed by microscopic *in vivo* analysis of conidiophore formation. The sensitivity to the strobilurin kresoxim-methyl was analysed using an *in vitro* germination assay and by determination of the diseased leaf area and conidia produced *in vivo*. Baseline sensitivities were determined with *V. inaequalis* populations from control orchards that had never been treated with fungicides. Comparison of the baseline sensitivities with sensitivities of populations obtained from orchards that had received 43 anilinopyrimidine treatments over 4 years, or from an orchard with 54 kresoxim-methyl treatments over 6 years indicated that no resistance to these fungicides has developed at the sites sampled.

### Zusammenfassung

Erfassung der Sensitivitäten gegenüber Anilinopyrimidin- und Strobilurinfungiziden in Populationen des Apfelschorfpilzes (*Venturia inaequalis*)

Die Sensitivität von *Venturia-inaequalis*-Populationen gegenüber den Anilinopyrimidinfungiziden Pyrimethanil und Cyprodinil wurde durch eine mikroskopische *in-vivo*-Analyse der Konidiophorenbildung analysiert. Die Sensitivität gegenüber dem Strobilurin Kresoxim-methyl wurde mittels eines *in-vitro*-Keimtests sowie durch eine Bestimmung der erkrankten Blattfläche und der Konidiobildung *in vivo* erfaßt. Die Basissensitivitäten wurden mit Hilfe von *V-inaequalis*-Populationen aus Kontroll-Äpfelanlagen ermittelt, die noch nie mit Fungiziden behandelt worden waren. Ein Vergleich der Basissensitivitäten mit den Sensitivitäten von Populationen aus Äpfelanlagen, die in 4 Jahren insgesamt 43 Anilinopyrimidinbehandlungen erhalten hatten, sowie aus einer Äpfelanlage, die in 6 Jahren insgesamt 54 Kresoxim-methylbehandlungen erhalten hatte, ergab, daß sich an den untersuchten Standorten keine Resistenz gegen diese Fungizide ausgebildet hatte.

### Introduction

The significance of *Venturia inaequalis* in Germany is indicated by the fact that up to 20 fungicide treatments are performed per season to chemically control apple scab disease (Kollar, 1997). Until 1995, only one group of fungicides with a specific mode of action, the curative sterol demethylation inhibitors (DMIs), was available in addition to nonspecific protective fungicides in Germany. Two years ago the anilinopyrimidine pyrimethanil, and last year the strobilurin kresoxim-methyl were introduced to control *V. inaequalis* in Germany.

Anilinopyrimidine fungicides have both a protective and a curative effect on apple scab, due to the strong inhibition of subcuticular growth of the fungus (Daniels et al., 1994; Knauf-Beiter et al., 1995). While pyrimethanil has been registered for scab control in Germany since 1996, cyprodinil is currently being tested against apple pathogens such as *Venturia*, *Alternaria*, and *Monilinia* (Heye et al., 1994).

The strobilurin kresoxim-methyl inhibits the spore germination of *V. inaequalis* and can thus be regarded as a protective fungicide (Gold et al., 1996). Like DMIs and benzimidazoles, strobilurins have one specific target in fungal metabolism and can thus be regarded as specific fungicides (Becker et al., 1981; Gold et al., 1996).

The fact that resistance to several fungicide classes has occurred in *V. inaequalis* populations treated with the respective fungicides (Fehrmann, 1976; Gilpatrick, 1982; Kiebacher and Hoffmann, 1980; Koller et al., 1997; Kunz et al., 1997), makes an assessment of the level of the sensitivity to new fungicides such as the anilinopyrimidines and strobilurins necessary. To accomplish this, methods are needed that allow the baseline sensitivity for these fungicides in untreated *V. inaequalis* populations to be determined, and shifts in sensitivity levels need to be followed after fungicide applications (Koller et al., 1997; Kunz et al., 1997; Siebels and Mendgen, 1994).

Several methods have been used to analyse fungicide efficiencies. Examples are spore germination assays or

radial growth measurements in the presence of different fungicide concentrations and determination of the minimal inhibitory fungicide concentration on agar plates (Gold et al., 1996; Hildebrand et al., 1988; Smith, 1991; Smith et al., 1991). Such *in vitro* assays may be used to analyse the effects of the protective strobilurins which inhibit spore germination, but they are inappropriate when the mechanism of the fungicide to be tested is related to infection or growth within the plant, e.g. for anilinopyrimidines.

Siebels and Mendgen (1994) have developed a microscopic *in vivo* method to assess the efficiencies of curative fungicides, and Kunz et al. (1997) have demonstrated that this method is appropriate to monitor the development of DMI-fungicide resistances at the population level. Since this method has the advantage that fungicide sensitivities of populations rather than of isolates are seen, and since fungal virulence and susceptibility of the plant tissue contribute to the results of this *in vivo* assay, this method is particularly suited to analyse fungal sensitivity against curative fungicides.

This paper describes the determination by *in vivo* and *in vitro* methods of the baseline sensitivities of *V. inaequalis* populations without fungicide history for the anilinopyrimidines pyrimethanil and cyprodinil, and for the strobilurin kresoxim-methyl. Sensitivities in populations treated with fungicides for up to 6 years were compared with the baseline sensitivity.

## Materials and Methods

### Test plants

Potted apple trees (*Malus domestica*) of the cultivar Golden Delicious, grafted on M 25 rootstocks at the Schweizerischen Forschungsanstalt für Obst und Weinbau, Wädenswil, Switzerland, were kept in a greenhouse at 18–25°C and a 16 h light period. The trees were fertilized weekly with 0.1% Hakaphos red (N:P:K = 8:12:24) and 0.1% Hakaphos blue (N:P:K = 15:10:15) (Compo GmbH, Münster, Germany).

### Fungicides

The anilinopyrimidines pyrimethanil (Scala EC 40; AgrEvo GmbH, Düsseldorf, Germany) and cyprodinil (CGD 20470 F WG 50; Ciba Geigy GmbH, Frankfurt/M., Germany), and the strobilurin kresoxim-methyl (BAS 490 F WG 50; BASF AG, Limburgerhof, Germany) were provided by the manufacturing companies as formulated test preparations. The DMI fungicide flusilazole (Benocap WG 20; DuPont de Nemours GmbH, Bad Homburg, Germany) and the protective fungicide dithianon (Delan SC 750; Cyanamid Agrar, Ingelheim, Germany) were used as commercial formulations.

### Inoculum

The inoculum was collected in six apple orchards at different locations in the Lake Constance area, Baden-Württemberg, Germany, and with different fungicide histories. Orchard 1 (located at Öschingen, with the apple variety Golden Delicious) and orchard 2 (Konstanz, Golden Delicious) served as control orchards that had

never been treated with fungicides. Orchard 3 (Litzelstetten, Golden Delicious) and orchard 4 (Lindau, Jonagold) are managed according to integrated control guidelines. From 1993 to 1995, between 13 and 17 fungicide treatments had been performed per season in these orchards. Until the last sampling in 1995, neither anilinopyrimidines nor strobilurins had been applied in orchards 3 and 4. Orchard 5 (Überlingen, Golden Delicious) is part of a test orchard of the Amtliche Pflanzenschutzberatung, a national advisory service, divided into four plots A, B, C and D with 20 apple trees each. Fungicide treatments performed in this test orchard are given in detail in Table 1. Orchard 6 is a test orchard of BASF AG (Ludwigshafen, located in Bavendorf, Baden-Württemberg) with Golden Delicious trees. Small plots of three trees each have been used to test different kresoxim-methyl concentrations and fungicide formulations since 1990. Leaves with scab lesions were collected in 1996 from control trees, untreated in 1996. However, 10, 8, 8, 10, 10 and 8 kresoxim-methyl treatments were performed in 1990, 1991, 1992, 1993, 1994 and 1995, respectively.

To compare sensitivity levels of *V. inaequalis* populations with different fungicide histories, naturally infected leaves with sporulating scab lesions were collected from the test orchards and stored in plastic bags at –70°C. Conidial suspensions were obtained by shaking the thawed leaves in 100 ml of deionized water. To obtain ascospores, naturally infected leaves were collected in November and overwintered in wire mesh cases on the ground of the orchard. The overwintered leaves were analysed for development of pseudothecia, beginning at the end of February of the following year. Apple leaves with pseudothecia containing 5–10% mature ascospores were collected on March 6, 1995 and April 22, 1996 and stored in plastic bags at –70°C. To obtain ascospore suspensions, the leaves were thawed and incubated in a dark moist chamber for 10 days at 20°C. Leaves containing mature, preconditioned pseudothecia were then washed in deionized water for 10 minutes and placed upside down in a glass Petri dish (14.5 cm), filled with 30 ml deionized water. The Petri dish was illuminated with 1800 lx from beneath for 1 h at +5°C to allow ascospore discharge into the water (Smereka et al., 1987). The conidia or ascospores from one orchard collected during 1 day are referred to as a sample.

### *In vitro* sensitivity test based on spore germination

To avoid fungicide contaminations in conidial suspensions obtained from treated orchards, conidia were propagated in the greenhouse as described (Kunz et al., 1997).

Spore suspensions ( $10^5$  spores/ml) were adjusted to contain different kresoxim-methyl concentrations (0.625 µg–0.625 mg/l) and applied to microscope slides. Duplicates of each treatment were incubated at 100% relative humidity at 20°C. Each fungicide concentration was incubated in a separate chamber to avoid obscuring results, due to vapour diffusion (Gold et al., 1996). The percentage of germinated conidia was determined by counting 200 specimens per microscope slide after 24 h.

Table 1  
Number of fungicide treatments in orchard 5 from 1993 to 1996. Only one fungicide was used per plot and season. Recommended fungicide doses (62.5 mg kresoxim-methyl/l; 300 mg pyrimethanil/l and 150 mg cyprodinil/l) were applied

Plot	1993		1994		1995		1996	
	Fungicide	n <sup>a</sup>	Fungicide	n <sup>a</sup>	Fungicide	n <sup>a</sup>	Fungicide	n <sup>b</sup>
A	Pyrimethanil	12	Cyprodinil	10	Cyprodinil	14	Cyprodinil	7
B	–	–	–	–	Pyrimethanil	14	Pyrimethanil	7
C	–	–	Pyrimethanil	8	–	–	–	–
D	–	–	Kresoxim-methyl	11	Kresoxim-methyl	11	Kresoxim-methyl	5

<sup>a</sup> Number of treatments; <sup>b</sup> Number of treatments until sampling (July 1, 1996).

The means of efficiencies were used to determine dose–response curves.

#### ***In vivo* sensitivity test based on determination of the portion of diseased leaf area and the amount of conidia formed**

Solutions containing different kresoxim-methyl concentrations (0.208–62.5 mg/l) were applied to four shoots until run off. Control plants were sprayed with deionized water. Twenty-four hours after the fungicide or control treatments, the plants were spray inoculated with 10<sup>5</sup> conidia/ml and subsequently incubated at 18°C and 100% relative humidity for 28 h in the dark. The plants were subsequently kept under greenhouse conditions as described above. Sixteen days after inoculation, the proportion of diseased leaf area of the three youngest inoculated leaves was determined (Dahmen and Staub, 1992), and these three leaves were shaken in 30 ml of deionized water for 1 min in order to obtain a conidial suspension. The conidial concentration was determined with a haematocytometer, and conidia formed were based on the leaf fresh weight (Dahmen and Staub, 1992). In total, the diseased leaf area of 36 leaves was assessed per fungicide concentration, and 12 shoots per fungicide concentration were analysed with respect to production of conidia. The resulting fungicide efficiencies were used to plot dose–response curves.

#### ***In vivo* sensitivity test based on microscopic evaluation of conidiophore formation**

Microscopic *in vivo* assays were performed as described by Kunz et al. (1997). Briefly, the plants were spray inoculated (10<sup>5</sup>/ml), and the inoculated plants were incubated in the dark at 18°C and 100% relative humidity for 24 h. Different fungicide concentrations (10–300 mg/l pyrimethanil or 5–150 mg/l cyprodinil) were applied by spraying with a glass atomizer 28 h after inoculation. Three shoots were used per treatment; the controls were sprayed with deionized water.

Six days after inoculation the two youngest inoculated leaves per shoot were glutaraldehyde-fixed, bleached and stained with methyl blue (Kunz et al., 1997). One-hundred infection sites were evaluated at 200 × magnification, and the reduction of conidiophore formation as depending on fungicide concentration was taken as a measure for fungicide sensitivity of *V. inaequalis* populations.

#### **Data analysis**

Different dose–response relationships were statistically compared with a factorial analysis using ranks (Meddis, 1984) with a macro written by Dr W. Nagl, Universität Konstanz, under SAS (Statistical Analysis System, SAS Institute, Inc., Cary, NC, USA). Dose–response relationships were considered significantly different at  $P < 0.01$ . The averages of the efficiencies were logit-transformed (logit efficiency =  $\ln(\text{efficiency}/1.0 - \text{efficiency})$ ) (Neter et al., 1983). Efficiencies of 100% had to be modified before logit-transformation. In microscopic evaluations, modified efficiency equals  $1 - 1/2n$ , in which  $n$  was the number of all infection sites in this treatment (Neter et al., 1983). A linear regression using the regression programme of the 'Statistica for Macintosh' (Statsoft Inc., Tulsa, OK, USA) between the logit-transformed efficiencies and the logarithm of the fungicide concentration revealed the ED<sub>50</sub>- and the ED<sub>90</sub>-value of the sample. The resistance factor was calculated by dividing the ED<sub>50</sub>-value of the sample by the ED<sub>50</sub>-value of the baseline sensitivity.

Different developmental stages of *V. inaequalis* were determined in the percentage of conidia that had germinated on the leaf. The means of a treatment were compared using the Mann–Whitney *U*-test (Non-parametrics of Statistica for Macintosh; Statsoft). The means were considered significantly different at  $P < 0.05$ .

#### **Results**

##### **Inhibition of *V. inaequalis* at different developmental stages by different fungicides**

The effect of the strobilurin kresoxim-methyl and of the anilinopyrimidines pyrimethanil and cyprodinil on spore germination were compared with the inhibitory effects of the contact fungicide dithianon and the DMI fungicide flusilazole (Table 2). Both, kresoxim-methyl and dithianon blocked spore germination in the *in vitro* assays, using the recommended fungicide concentrations, indicating that such assays can thus be used to determine the kresoxim-methyl sensitivity of *V. inaequalis* populations. Both, anilinopyrimidines and the DMI fungicide tested had a negligible effect on spore germination at recommended concentrations.

Microscopic *in vivo* assays were used to investigate the inhibitory effects of the anilinopyrimidines in comparison to the DMI flusilazole on *V. inaequalis* infection structure

**Table 2**  
Efficiency of fungicides determined by *in vitro* germination assays. Conidia of *Venturia inaequalis* were collected from control orchard 1 and allowed to germinate on glass microscope slides in the presence of recommended fungicide concentrations. Germination in deionized water served as control

Fungicide	Concentration (mg/l)	Germination rate (%)	Efficiency
Water		56.4	
Kresoxim-methyl	62.50	0.3	99
Pyrimethanil	300	41.0	27
Cyprodinil	150	57.1	-1
Flusilazole	25	56.5	0
Dithianon	450	0	100

differentiation. Apple leaves had been treated with fungicides either 1 day prior to inoculation, or, to demonstrate the curative effects, 1, 2 or 3 days post-inoculation. Pyrimethanil and cyprodinil only weakly inhibited the germination and penetration hypha differentiation on host leaves, but differentiation of subcuticular structures, i.e. primary subcuticular stroma, runner hyphae and secondary subcuticular stroma, was almost completely blocked in the presence of these fungicides (Fig. 1). When

the fungicides were applied 1 day after inoculation, the development of primary subcuticular stroma and especially of runner hyphae were inhibited (Fig. 1).


These results indicate that microscopic *in vivo* assays based on conidiophore formation (Kunz et al., 1997) are appropriate to test for shifts in the anilinopyrimidine sensitivity of *V. inaequalis* populations.

#### Sensitivity of *V. inaequalis* populations to the strobilurin fungicide kresoxim-methyl

Baseline sensitivity was determined in *in vitro* germination assays, using eight samples from four orchards without strobilurin history (Table 3). The ED<sub>50</sub>-values of these samples varied from 0.0005 to 0.0125 mg/l; statistical differences of dose-response relationships of these samples have not been detected ( $P = 0.72$ ), and the curves were thus unified to yield the baseline sensitivity (Fig. 2). After logit transformation, an ED<sub>50</sub>-value of 0.0065 mg/l kresoxim-methyl was determined for baseline sensitivity (Fig. 2, insert).

Since *in vitro* germination assays are performed by incubating the spores directly in fungicide suspensions, the field concentrations yielding the same effects must be higher. Determination of shifts in fungicide sensitivity *in vitro* does not allow a prediction of the magnitudes of the sensitivity shifts at which fungicide efficiencies would be insufficient in the field. Therefore, comparative *in vivo* studies with *V. inaequalis* conidia, collected in the untreated control orchard 2 in 1995 were performed. The results show that the ED<sub>50</sub>-value determined by measurement of diseased leaf area is 137-fold, and that determined by counting the conidia formed is 104-fold higher than the ED<sub>50</sub>-value determined by *in vitro* germination assays (Fig. 3). The magnitude of a sensitivity shift in fungal populations necessary to affect fungicide efficiencies in the field has to exceed the safety margin, which was defined as the quotient of the recommended field concentration and the ED<sub>90</sub>-value of the baseline sensitivity. Thus, a safety margin of 41 (measurement of diseased leaf area) or 80 (counting conidia formed) was found in the case of kresoxim-methyl at the recommended concentration of 62.5 mg/l.

Twenty-seven kresoxim-methyl applications had been performed over 3 years in plot D of orchard 5, and 54 applications of this fungicide had been performed over 6 years in orchard 6, and no significant difference to baseline sensitivity was found using germination assays



**Fig. 1** Development of *Venturia inaequalis* on apple leaves treated with different fungicides. Flusilazole (25 mg/l), cyprodinil (150 mg/l) or pyrimethanil (300 mg/l) were applied to the leaves 1 day before (1 d a.i.), 1, 2 or 3 days after inoculation (1, 2 or 3 d p.i.) with conidia collected from control orchard 1. Penetration pores (□), primary stroma (■), runner hyphae (■), secondary stroma (■), and conidiophores (■) were counted 6 days after inoculation. The columns represent means of 10 leaves; bars represent standard errors. Different letters over the columns indicate significant differences in the respective developmental stage ( $P < 0.05$ ) according to the Mann-Whitney *U*-test

Table 3  
ED<sub>50</sub>- and ED<sub>90</sub>-values of *Venturia inaequalis* populations from control orchards for kresoxim-methyl. All values except those for multiplied conidia of orchard 2 were determined by *in vitro* germination assays. Baseline sensitivity represents the combination of all samples tested *in vitro*. Resistance factors are based on ED<sub>50</sub>-values

Orchard	Sample <sup>a</sup>	ED <sub>50</sub> (mg/l)	ED <sub>90</sub> (mg/l)	Resistance-factor (ED <sub>50</sub> )
Orchard 1	conidia 1994 <sup>b</sup>	0.0015	0.011	0.24
	multiplied conidia 1994 <sup>b</sup>	0.0035	0.013	0.53
	ascospores 1995 <sup>b</sup>	0.0005	0.011	0.07
	multiplied conidia 1995 <sup>b</sup>	0.0125	0.042	1.93
Orchard 2	multiplied conidia 1995 <sup>b</sup>	0.0045	0.031	0.69
	multiplied conidia 1995 <sup>c</sup>	0.62	1.51	
	multiplied conidia 1995 <sup>d</sup>	0.47	0.78	
Orchard 3	multiplied conidia 1994 <sup>b</sup>	0.0100	0.059	1.54
Orchard 4	multiplied conidia 1994 <sup>b</sup>	0.0135	0.104	2.08
	ascospores 1995 <sup>b</sup>	0.0076	0.023	1.17
Baseline <sup>b</sup>		0.0065	0.042	1.00

<sup>a</sup> Spore type and year of sampling; <sup>b</sup> *in vitro* germination assay; <sup>c</sup> *in vivo* determination of portion of diseased leaf area; <sup>d</sup> *in vivo* determination of the number of conidia produced.




Fig. 2 Baseline sensitivity of *Venturia inaequalis* populations for kresoxim-methyl. *In vitro* germination assays were performed with eight samples from control orchards. Means of 32 microscope slides and standard deviations are given. The insert shows the logit transformed efficiency curve ( $R = 0.989$ ); the point of intersection indicates an ED<sub>50</sub> of 0.0065 mg kresoxim-methyl/l




Fig. 3 Comparison of *in vitro* and *in vivo* sensitivity assays. Multiplied conidia of *Venturia inaequalis* collected from control orchard 2 in 1995 were used for *in vitro* germination assays (●), and to determine diseased leaf area (■) or the number of conidia produced (▲) in the presence of kresoxim-methyl. The arrow indicates the recommended kresoxim-methyl concentration

(Table 4). Conidia of orchard 6, which had been exposed to the most drastic selection pressure, were analysed with respect to diseased leaf area and the number of conidia formed. Again, no significant differences to baseline sensitivity were found.

#### Sensitivity of *V. inaequalis* populations to anilinopyrimidine fungicides

To determine the baseline sensitivity of *V. inaequalis* populations to pyrimethanil and cyprodinil, *in vivo* sensitivity tests were performed with conidia from orchard 1, collected in 1994 and 1995, and with conidia from orchard 2 of 1995. Dose-response relationships of these three samples showed no significant statistical difference for pyrimethanil ( $P = 0.90$ ), and for cyprodinil ( $P = 0.66$ ). The curves were thus combined to determine the baseline sensitivity (Figs 4, 5). Logit-transformation allowed the calculation of ED<sub>50</sub>-values of 35.5 mg/l pyrimethanil and of 14.5 mg/l cyprodinil (inserts of Figs 4, 5).

With the recommended field concentrations of 300 mg/l pyrimethanil and 150 mg/l cyprodinil, fungicide efficiencies of 100% and 99% were found; the safety margin was 4.6 for pyrimethanil and 3.6 for cyprodinil.

Samples were collected from orchard 5 to evaluate shifts in the anilinopyrimidine sensitivities in *V. inaequalis* populations. In the pyrimethanil-treated plots B and C, samples were taken in October 1994 (plot C) and in June and September 1995 (plot B). Dose-response relationships of these populations were not significantly different from the baseline sensitivity. Resistance factors for pyrimethanil were between 0.92 and 1.69 (Table 5).


In plot A, samples were taken in October 1994, June 1995 and July 1996 to analyse the cyprodinil sensitivity. Again, dose-response relationships indicated no significant differences as compared with the baseline sensitivity (Table 5), and after logit-transformation and determination of ED<sub>50</sub>-values, the resistance factors between 0.69 and 1.13 were calculated for this fungicide (Table 5).

Taken together, after 43 anilinopyrimidine treatments in plot A from 1993 to 1996, no shift in anilinopyrimidine sensitivity in *V. inaequalis* populations has been detected.


**Table 4**  
ED<sub>50</sub>-values and resistance factors of *Venturia inaequalis* populations from kresoxim-methyl-treated orchards. All values were determined by *in vitro* germination assays

Sample <sup>a</sup>	ED <sub>50</sub> (mg/l)	Resistance-factor	P-value <sup>b</sup>
Baseline sensitivity	0.0065		
Orchard 5 part D			
multiplied conidia 1994	0.0117	1.8	0.148
ascospores 1995	0.0064	1.0	0.398
multiplied conidia 1995	0.0288	4.5	0.013
ascospores 1996	0.0240	3.7	0.102
multiplied conidia 1996	0.0081	1.3	0.790
Orchard 6:			
conidia 1996	0.0036	0.6	0.926

<sup>a</sup> Orchard, spore type and year of sampling; <sup>b</sup> P-value after comparison of the dose-response relationship of the sample with the baseline sensitivity using factorial analysis of ranks.



**Fig. 4** Baseline sensitivity of *Venturia inaequalis* populations for pyrimethanil. Conidia collected from three different nontreated control orchards were used to determine fungicide sensitivity by microscopic *in vivo* analysis of conidiophores. A total of 24 leaves per fungicide concentration was examined. The arrow indicates the recommended pyrimethanil concentration. The insert shows the logit transformed efficiency curve ( $R = 0.966$ ); the point of intersection indicates an ED<sub>50</sub> of 35.5 mg pyrimethanil/l



**Fig. 5** Baseline sensitivity of *Venturia inaequalis* populations for cyprodinil. Conidia collected from three different nontreated control orchards were used to determine fungicide sensitivity by microscopic *in vivo* analysis of conidiophores. A total of 24 leaves per fungicide concentration was examined. The arrow indicates the recommended cyprodinil concentration. The insert shows the logit transformed efficiency curve ( $R = 0.991$ ); the point of intersection indicates an ED<sub>50</sub> of 14.5 mg cyprodinil/l

## Discussion

Assays to analyse the sensitivity shifts in fungal populations must depend on the mode of fungicide action. Anilinopyrimidines act through inhibition of secretion of enzymes such as cutinases, lipases, invertases, and of cell wall-degrading enzymes. This group of fungicides thus exhibits a novel fungicidal mechanism (Milling and Richardson, 1995; Miura et al., 1994). Extracellular enzymes have been discussed to contribute to fungal virulence (Mendgen et al., 1996; Schäfer, 1994). Cutinases, pectic and cellulolytic enzymes have been demonstrated in *V. inaequalis* (Kollar, 1994; Köller et al., 1991; Valsangiacomo and Gessler, 1992), and these enzymes could play a critical role in the penetration and/or nutrient supply of the fungus (Kollar, 1997; MacHardy, 1996). Thus, the interference of anilinopyrimidines with enzyme secretion is consistent with the results obtained from microscopic analyses of fungal infection sites on fungicide-treated leaves (Fig. 1). While the formation of a penetration peg can often be observed, extended subcuticular development requiring enzyme secretion is completely blocked (Daniels et al., 1994; Knauf-Beiter et al., 1995). However, another suspected mode of action of these fungicides, i.e. the inhibition of methionine biosynthesis (Fritz et al., 1997; Masner et al., 1994) could also lead to the inhibition of subcuticular growth, as suggested in an early paper by Kline et al. (1957). These studies showed that mutants of *V. inaequalis*, deficient in methionine biosynthesis, were arrested after penetration of the cuticle and were not able to differentiate conidiophores and conidia. Exogenously applied methionine restored both subcuticular growth and conidiation (Kline et al., 1957). These results indicate that anilinopyrimidines affect the fungus-plant interaction, rather than saprophytic growth.

It has recently been shown that *Colletotrichum* and *Fusarium* species exhibit a targeted secretion of cutinolytic enzymes only during infection structure differentiation (Podila et al., 1995). However, *V. inaequalis* does not differentiate infection structures during growth on agar plates (Smith, 1991). Probably, an infection-related secretion of enzymes is not needed during growth on agar. Therefore, a possible inhibition of enzyme secretion may not inhibit growth on agar, as has been shown with *Botryotinia fuckeliana*. With this fungus, ani-

Table 5  
ED<sub>50</sub>-values and resistance factors of *Venturia inaequalis* populations for anilinopyrimidines. All values were determined by microscopic determination of conidiophores *in vivo*

Sample <sup>a</sup>	Number of treatments <sup>b</sup>	ED <sub>50</sub> (mg/l)	Resistance-factor	P-value <sup>c</sup>
Pyrimethanil				
Baseline sensitivity	0	35.5		
Orchard 5 plot C				
October 1994	8	60.0	1.69	0.92
Orchard 5 plot B				
June 1995	9	33.9	0.95	0.77
September 1995	14	32.7	0.92	0.97
Cyprodinil				
Baseline sensitivity	0	14.5	–	
Orchard 5 plot A				
October 1994	22	14.2	0.98	0.21
June 1995	30	16.4	1.13	0.31
July 1996	43	10.0	0.69	0.33

<sup>a</sup> Orchard and month of sampling. Only conidia were evaluated; <sup>b</sup> Number of pyrimidinamine treatments (pyrimethanil or cyprodinil) in the orchard since 1993; <sup>c</sup> P-value after comparison of the dose–response relationships of the sample with the baseline using factorial analysis of ranks.

linopyrimidines only slightly reduced growth on nutrient-rich media (Hilber and Schüepp, 1996; Masner et al., 1994). Obviously, *in vitro* assays may not reflect the situation in nature. Sensitive isolates could thus be falsely classified as resistant in *in vitro* assays, where infection structures are not differentiated (Birchmore et al., 1996). Therefore, the recent report of significant differences in the sensitivities of *V. inaequalis* isolates (Schnabel and Parisi, 1997) may not be valid under field conditions. The *in vitro* tests performed by these authors were done with spores growing on complex media which have been shown to lead to extended variations in fungicide sensitivity (Birchmore et al., 1996).

As a consequence, *in vivo* tests involving the host plants (Kunz et al., 1997; Siebels and Mendgen, 1994) are an absolute necessity to correctly investigate sensitivity shifts in *V. inaequalis* populations to anilinopyrimidine fungicides.

The results of the present study, based on microscopic *in vivo* analyses, indicate that, under the experimental conditions used, no reduced fungicide sensitivities were found in *V. inaequalis* populations, even after a total of 43 anilinopyrimidine applications in 4 years.

Birchmore et al. (1996) have reported that after 15 anilinopyrimidine applications in grapes over 5 years, no sensitivity shifts have been found in *B. fuckeliana*. However, after 8 years of cyprodinil application, reduced fungicide efficiencies to this fungus have been detected by Hilber and Schüepp (1996).

The strobilurin kresoxim-methyl specifically inhibits germination of *V. inaequalis* conidia (Gold et al., 1996). This inhibition is due to blocking of the electron transport at the cytochrome bc<sub>1</sub>-complex, and kresoxim-methyl can thus be regarded as an inhibitor of mitochondrial respiration (Becker et al., 1981; Mizutani et al., 1995). This high specificity and the fact that strobilurin A-producing fungi such as *Strobilurus tenacellus* and *Mycena galopoda* are insensitive to the drug, due to a single amino acid exchange in their cytochrome b, make the occurrence of fungicide resistance likely (Kraiczky et al., 1996). Also in the yeasts *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*, resistance has been shown to be

mediated by amino acid exchange in the cytochrome molecule (DiRago et al., 1989; Kraiczky et al., 1996).

To measure the inhibition of germination the host plant is not required, and sensitivity tests can therefore be performed on microscope slides (Table 2) or agar plates (Smith, 1991). A disadvantage of this method is that *in vitro* the inhibitory concentrations are smaller than those determined in the field (Fig. 3), and from such experiments no conclusions concerning the magnitude of sensitivity shifts necessary to reduce the fungicide efficiency in the field can be drawn. It is therefore important to determine the safety margin by *in vivo* sensitivity tests on the plant. Measurements of the diseased leaf area or production of conidia would be appropriate *in vivo* assays.

In the present studies, the highest selection pressure has been applied in orchard 6. Fifty-four kresoxim-methyl treatments have been performed in 6 years, but no significant difference from baseline sensitivity was found in germination assays and *in vivo* assays based on diseased leaf area or production of conidia. This indicates that a shift in sensitivity did not occur. To reduce efficiency in the field, a sensitivity shift by a factor of 40 would be needed. The highest resistance factor found in the field, however, was 4.5.

So far, resistance in *V. inaequalis* populations to strobilurins or anilinopyrimidines has not been described. In the present investigations, these fungicides have been tested in comparatively small apple scab populations, and treatment with these fungicides was carried out in a limited time frame of 4 and 6 years, respectively. This, as indicated by the studies by Birchmore et al. (1996) and Hilber and Schüepp (1996), may not be sufficient for resistance to develop in a fungal population. Under conditions of agricultural practice, however, the fungal population would be significantly larger, and fungicide application would occur over a longer time than in the present investigations. These factors may drastically increase the probability of selection of isolates with reduced sensitivity to strobilurins or anilinopyrimidines (Scheinpflug, 1988; Veverka, 1996).

Based on these considerations, it cannot be concluded

that resistance to the new fungicides tested in this paper is unlikely to occur. However, these fungicides registered for *V. inaequalis* control will certainly be of value, especially in combinations with other fungicides. Alternate application of fungicides with different modes of inhibition will help to reduce the risk of resistance occurring. In addition, since the anilinoimidazole derivatives are curative fungicides, they could be used in combination with DMIs in integrated disease control and would thus help to reduce the overall amount of chemicals applied in the field.

#### Acknowledgements

The authors thank BASF AG, Limburgerhof, for donation of kresoxim-methyl (BAS 490 F) and apple leaf samples from the test orchard in Bavendorf. Ciba-Geigy GmbH, Frankfurt, and AgrEvo GmbH, Düsseldorf, which provided test preparations of cyprodinil (CGD 20470 F) and pyrimethanil (SCH 32050 F). They especially thank E. Lange, Übergebieliche Pflanzenschutzberatung Überlingen, for the supply of apple leaf samples from test orchards, and for stimulating discussions.

#### Literature

- Becker, W. F., G. von Jagow, T. Anke, W. Steglich (1981): Oudemansin, strobilurin A, strobilurin B and myxothiazol: new inhibitors of the bc<sub>1</sub> segment of the respiratory chain with an  $\alpha$ - $\beta$ -methoxyacrylate system as common structural element. *FEBS Lett.* **132**, 329–333.
- Birchmore, R. J., R. J. Williams, R. J. Milling, P. E. Russell (1996): Development of methods for monitoring populations of *Botrytis cinerea* for sensitivity to pyrimethanil. In: Lyr, H., P. E. Russel and H. D. Sisler (eds), *Modern Fungicides and Antifungal Compounds*, pp. 377–385. Intercept Ltd, Andover, UK.
- Dahmen, H., T. Staub (1992): Protective, curative, and eradicator activity of difenoconazole against *Venturia inaequalis*, *Cercospora arachidicola*, and *Alternaria solani*. *Plant Dis.* **76**, 774–777.
- Daniels, A., R. J. Birchmore, E. H. Winter (1994): Activity of pyrimethanil on *Venturia inaequalis*. *Proc. Br. Crop. Prot. Conf. – Pests Dis.* **4**, 525–532.
- DiRago, J.-P., J.-Y. Coppée, A.-M. Colson (1989): Molecular basis for resistance to myxothiazol, mucidin (strobilurin A), and stigmatellin. *J. Biol. Chem.* **264**, 14543–14548.
- Fehrmann, H. (1976): Systemische Fungizide-ein Überblick II. Fungizidresistenz phytopathogener Pilze. *J. Phytopathol.* **86**, 144–185.
- Fritz, R., C. Lanen, V. Colas, P. Leroux (1997): Inhibition of methionine biosynthesis in *Botrytis cinerea* by the anilinoimidazole fungicide pyrimethanil. *Pestic. Sci.* **49**, 40–46.
- Gilpatrick, J. D. (1982): Case study 2: *Venturia* of pome fruits and *Monilinia* of stone fruits. In: Dekker J. and S. G. Georgopoulos (eds), *Fungicide Resistance in Crop Protection*, pp. 195–206. Centre for Agricultural Publishing and Documentation, Wageningen.
- Gold, R. E., E. Ammermann, H. Köhle, G. M. E. Leinhos, G. Lorenz, J. B. Speakman, M. Stark-Urnau, H. Sauter (1996): The synthetic strobilurin BAS 490 F: profile of a modern fungicide. In: Lyr, H., P. E. Russel and H. D. Sisler (eds), *Modern Fungicides and Antifungal Compounds*, pp. 79–92. Intercept Ltd, Andover, UK.
- Heye, U. J., J. Speich, H. Siegle, R. Wohlhauser, A. Hubele (1994): CGA 219417-A novel broad-spectrum fungicide. *Proc. Br. Crop. Prot. Conf. – Pests Dis.* **4**, 501–508.
- Hilber, U. W., H. Schüpp (1996): A reliable method for testing the sensitivity of *Botryotinia fuckeliana* to anilinoimidazole derivatives. *Pestic. Sci.* **47**, 241–247.
- Hildebrand, P. D., C. L. Lockhart, R. J. Newberry, R. G. Ross (1988): Resistance of *Venturia inaequalis* to bitertanol and other demethylation-inhibiting fungicides. *Can. J. Plant Pathol.* **10**, 311–316.
- Kiebacher, J., G. M. Hoffmann (1980): Qualitative und quantitative Untersuchungen zur Resistenz von *Venturia inaequalis* gegen Benzimidazol-Fungizide. *Z. PflKrankh. PflSchutz* **87**, 705–716.
- Kline, D. M., D. M. Boone, G. W. Keitt (1957): *Venturia inaequalis* (Cke.) Wint. XIV. Nutritional control of pathogenicity of certain induced biochemical mutants. *Am. J. Bot.* **44**, 797–803.
- Knauf-Beiter, G., H. Dahmen, U. Heye, T. Staub (1995): Activity of cyprodinil: optimal treatment timing and site of action. *Plant Dis.* **79**, 1089–1103.
- Kollar, A. (1994): Characterization of specific induction, activity, and isozyme polymorphism of extracellular cellulases from *Venturia inaequalis* detected *in vitro* and on the host plant. *Mol. Plant-Microbe Interact.* **7**, 603–611.
- Kollar, A. (1997): Aktuelle Forschung an dem bedeutendsten Erreger im Apfelanbau, dem Apfelschorfpilz *Venturia inaequalis*. *Nachrichtenbl. Deut. Pflanzenschutzd.* **49**, 131–136.
- Köller, W., D. M. Parker, C. M. Becker (1991): Role of cutinase in the penetration of apple leaves by *Venturia inaequalis*. *Phytopathology* **81**, 1375–1379.
- Köller, W., W. F. Wilcox, J. Barnard, A. L. Jones, P. G. Braun (1997): Detection and quantification of resistance of *Venturia inaequalis* populations to sterol demethylation inhibitors. *Phytopathology* **87**, 184–190.
- Kraiczky, P., U. Haase, S. Gencic, S. Flindt, T. Anke, U. Brandt, G. von Jagow (1996): The molecular basis for the natural resistance of the cytochrome bc<sub>1</sub> complex from strobilurin-producing basidiomycetes to center Q<sub>o</sub> inhibitors. *Eur. J. Biochem.* **235**, 54–63.
- Kunz, S., H. Deising, K. Mendgen (1997): Acquisition of resistance to sterol demethylation inhibitors by populations of *Venturia inaequalis*. *Phytopathology* **87**, 1272–1278.
- MacHardy, W. E. (1996): *Apple Scab*, pp. 161–162, APS Press, St. Paul, MN.
- Masner, P., P. Muster, J. Schmid (1994): Possible methionine biosynthesis inhibition by anilinoimidazole fungicides. *Pestic. Sci.* **42**, 163–166.
- Meddis, R. (1984): *Statistics Using Ranks: A Unified Approach*, pp. 299–345, Basil Blackwell Publisher Ltd., Oxford.
- Mendgen, K., M. Hahn, H. Deising (1996): Morphogenesis and mechanisms of penetration by plant pathogenic fungi. *Ann. Rev. Phytopathol.* **34**, 367–386.
- Milling, R. J., C. J. Richardson (1995): Mode of action of the anilinoimidazole fungicide Pyrimethanil. 2. Effects on enzyme secretion in *Botrytis cinerea*. *Pestic. Sci.* **45**, 43–48.
- Miura, I., T. Kamakura, S. Maeno, S. Hayashi, I. Yamaguchi (1994): Inhibition of enzyme secretion in plant pathogens by Mepanipyrim, a novel fungicide. *Pestic. Biochem. Physiol.* **48**, 222–229.
- Mizutani, A., H. Yukioka, H. Tamura, N. M. Iki, M. Masuko, R. Takeda (1995): Respiratory characteristics in *Pyricularia oryzae* exposed to a novel alkoxyiminoacetamide fungicide. *Phytopathology* **85**, 306–311.
- Neter, J., W. Wassermann, M. H. Kutner (1983): *Applied Linear Regression Models*, pp. 362–367, Richard D. Irwin, Inc, Homewood, IL.
- Podila, G. K., E. Rosen, M. J. D. S. Francisco, P. E. Kollatukudy (1995): Targeted secretion of cutinase in *Fusarium solani* f. sp. *pisi* and *Colletotrichum gloeosporioides*. *Phytopathology* **85**, 238–242.
- Schäfer, W. (1994): Molecular mechanisms of fungal pathogenicity to plants. *Ann. Rev. Phytopathol.* **32**, 461–477.
- Scheinflug, H. (1988): History of DMI fungicides and monitoring for resistance. In: Delp C. J. (ed.), *Fungicide Resistance in North America*, pp. 77–78. American Phytopathological Society, St. Paul, MN.
- Schnabel, G., L. Parisi (1997): Sensitivity of *Venturia inaequalis* to five DMI fungicides, including the new triazole fluquinconazole, and to pyrimethanil. *Z. PflKrankh. PflSchutz* **104**, 36–46.
- Siebels, C., K. Mendgen (1994): A microscopic evaluation of the sensitivity of *Venturia inaequalis* populations to sterol demethylation inhibitors. *Mycol. Res.* **98**, 619–624.
- Smereka, K. J., W. E. MacHardy, A. P. Kausch (1987): Cellular differentiation in *Venturia inaequalis* ascospores during germination and penetration of apple leaves. *Can. J. Bot.* **65**, 2549–2561.
- Smith, C. M. (1991): *In vitro* methods for monitoring flusilazole sensitivity in populations of *Venturia inaequalis* (apple scab) and *Cercosporidium personatum* (peanut late leaf spot). *EPPO-Bulletin* **21**, 323–328.
- Smith, F. D., D. M. Parker, W. Köller (1991): Sensitivity distribution of *Venturia inaequalis* to the sterol demethylation inhibitor flusilazole: baseline sensitivity and implications for resistance monitoring. *Phytopathology* **81**, 392–396.
- Valsangiacomo, C., C. Gessler (1992): Purification and characterization of an exo-polygalacturonase produced by *Venturia inaequalis*, the causal agent of apple scab. *Physiol. Molec. Plant Pathol.* **40**, 63–77.
- Veverka, K. (1996): General concepts of fungicide resistance. In: Lyr, H., P. E. Russel and H. D. Sisler (eds), *Modern Fungicides and Antifungal Compounds*, pp. 283–292. Intercept Ltd, Andover, UK.